

# *Bibliografia completa*

- AGOSTA, F., SALA, S., VALSASINA, P., MEANI, A., CANU, E., MAGNANI, G., y cols. (2013). Brain network connectivity assessed using graph theory in frontotemporal dementia. *Neurology*, 81(2), 134-143.
- ALAM, S. S., JUNAID, S., y AHMED, S. M. (2016). Evaluation of Technetium-99m glucoheptonate single photon emission computed tomography for brain tumor grading. *Asian Journal of Neurosurgery*, 11(2), 118-128.
- ALBRECHT, D. S., KAREKEN, D. A., CHRISTIAN, B. T., DZEMIDZIC, M., y YODER, K. K. (2014). Cortical dopamine release during a behavioral response inhibition task. *Synapse*, 68(6), 266-274.
- ALLMAN, J. M., TETREALT, N. A., HAKEEM, A. Y., MANAYE, K. F., SEMENDEFERI, K., ERWIN, J. M., y cols. (2011). The von Economo neurons in the frontoinsular and anterior cingulate cortex. *Annals of the New York Academy of Sciences*, 1225, 59-71.
- ANDERSON, B. A., KUWABARA, H., WONG, D. F., GEAN, E. G., RAHMIM, A., BRAŠIĆ, J. R., y cols. (2016). The Role of Dopamine in Value-Based Attentional Orienting. *Current Biology: CB*, 26(4), 550-555.
- ANDREASEN, N. C., REZAI, K., ALLIGER, R., SWAYZE, V. W., FLAUM, M., KIRCHNER, P., y cols. (1992). Hypofrontality in neuroleptic-naive patients and in patients with chronic schizophrenia. Assessment with xenon 133 single-photon emission computed tomography and the Tower of London. *Archives of General Psychiatry*, 49(12), 943-958.
- BALLANGER, B., BARADUC, P., BROUSSOLLE, E., LE BARS, D., DESMURGET, M., y THOBOIS, S. (2008). Motor urgency is mediated by the contralateral cerebellum in Parkinson's disease. *Journal of Neurology, Neurosurgery, and Psychiatry*, 79(10), 1110-1116.
- Bassett, D. S., y Bullmore, E. (2006) Small-world brain networks. *The Neuroscientist: A Review Journal Bringing Neurobiology, Neurology and Psychiatry*, 12(6), 512-523.
- BECKMANN, C. F., DeLUCA, M., DEVLIN, J. T., y SMITH, S. M. (2005). Investigations into resting-state connectivity using independent component analysis. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, 360(1457), 1001-1013.
- BIANCARDI, M., FUKUNAGA, M., VAN GELDEREN, P., HOROVITZ, S. G., DE ZWART, J. A., SHMUELI, K., y DUYN, J. H. (2009). Sources of functional magnetic resonance imaging signal fluctuations in the human brain at rest: a 7 T study. *Magnetic Resonance Imaging*, 27(8), 1019-1029.
- BIRN, R. M., MOLLOY, E. K., PATRIAT, R., PARKER, T., MEIER, T. B., KIRK, G. R., y cols. (2013). The effect of scan length on the reliability of resting-state fMRI connectivity estimates. *NeuroImage*, 83, 550-558.

- BISWAL, B. B., MENNES, M., ZUO, X.-N., GOHEL, S., KELLY, C., SMITH, S. M., y cols. (2010) Toward discovery science of human brain function. *Proceedings of the National Academy of Sciences of the United States of America*, 107(10), 4734-4739.
- BISWAL, B., YETKIN, F. Z., HAUGHTON, V. M., y HYDE, J. S. (1995). Functional connectivity in the motor cortex of resting human brain using echo-planar MRI. *Magnetic Resonance in Medicine*, 34(4), 537-541.
- BLAMIRE, A. M., OGAWA, S., UGURBIL, K., ROTHMAN, D., MCCARTHY, G., ELLERMANN, J. M., y cols. (1992). Dynamic mapping of the human visual cortex by high-speed magnetic resonance imaging. *Proceedings of the National Academy of Sciences of the United States of America*, 89(22), 11069-11073.
- BREMNER, J. D., INNIS, R. B., WHITE, T., FUJITA, M., SILBERSWEIG, D., GODDARD, A. W., y cols. (2000). SPECT [I-123]iomazenil measurement of the benzodiazepine receptor in panic disorder. *Biological Psychiatry*, 47(2), 96-106.
- BRITTON, J. W., FREY, L. C., HOPP, J. L., KORB, P., KOUBEISSI, M. Z., LIEVENS, W. E., y cols. En St. LOUIS, E. K. (ed), FREY L. C. (ed), (2016). *Electroencephalography (EEG): An Introductory Text and Atlas of Normal and Abnormal Findings in Adults, Children, and Infants*. Chicago, EUA: American Epilepsy Society.
- BUCKNER, R. L. (2010). Human functional connectivity: new tools, unresolved questions. *Proceedings of the National Academy of Sciences of the United States of America*, 107(24), 10769-10770.
- BUCKNER, R. L., SEPULCRE, J., TALUKDAR, T., KRIENEN, F. M., LIU, H., HEDDEN, T., y cols. (2009). Cortical hubs revealed by intrinsic functional connectivity: mapping, assessment of stability, and relation to Alzheimer's disease. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 29(6), 1860-1873.
- BULLMORE, E. T., y BASSETT, D. S. (2011). Brain graphs: graphical models of the human brain connectome. *Annual Review of Clinical Psychology*, 7, 113-140.
- BUTTS, C. T. (2009). Revisiting the foundations of network analysis. *Science*, 325(5939), 414-416.
- CATAFAU, A. M., DANUS, M., BULLICH, S., NUCCI, G., LLOP, J., ABANADES, S., y cols. (2006). Characterization of the SPECT 5-HT<sub>2A</sub> receptor ligand 123I-R91150 in healthy volunteers: part 2--ketanserin displacement. *Journal of Nuclear Medicine: Official Publication, Society of Nuclear Medicine*, 47(6), 929-937.
- CHAO-GAN, Y., y YU-FENG, Z. (2010). DPARSF: A MATLAB Toolbox for "Pipeline" Data Analysis of Resting-State fMRI. *Frontiers in Systems Neuroscience*, 4, 13.
- CHENG, Y., XU, J., NIE, B., LUO, C., YANG, T., LI, H., y cols. (2013). Abnormal resting-state activities and functional connectivities of the anterior and the posterior cortexes in medication-naïve patients with obsessive-compulsive disorder. *PLoS One*, 8(6), e67478.
- CIRIC, R., WOLF, D. H., POWER, J. D., ROALF, D. R., BAUM, G. L., RUPAREL, K., y cols. (2017). Benchmarking of participant-level confound regression strategies for the control of motion artifact in studies of functional connectivity. *NeuroImage*, 154, 174-187.
- COLE, D. M., SMITH, S. M., y BECKMANN, C. F. (2010). Advances and pitfalls in the analysis and interpretation of resting-state FMRI data. *Frontiers in Systems Neuroscience*, 4, 8.

## *Bibliografía completa*

- COLLOBY, S. J., MCKEITH, I. G., BURN, D. J., WYPER, D. J., O'BRIEN, J. T., y TAYLOR, J. P. (2016). Cholinergic and perfusion brain networks in Parkinson disease dementia. *Neurology*, 87(2), 178-185.
- COLLOBY, S. J., MCKEITH, I. G., WYPER, D. J., O'BRIEN, J. T., y TAYLOR, J. P. (2015). Regional covariance of muscarinic acetylcholine receptors in Alzheimer's disease using (R, R) [(123) I]-QNB SPECT. *Journal of Neurology*, 262(9), 2144-2153.
- CONTRERAS-RODRÍGUEZ, O., PUJOL, J., BATALLA, I., HARRISON, B. J., BOSQUE, J., IBERN-REGÀS, I., y cols. (2014). Disrupted neural processing of emotional faces in psychopathy. *Social Cognitive and Affective Neuroscience*, 9(4), 505-512.
- CORDES, D., HAUGHTON, V., CAREW, J. D., ARFANAKIS, K., y MARAVILLA, K. (2002). Hierarchical clustering to measure connectivity in fMRI resting-state data. *Magnetic Resonance Imaging*, 20(4), 305-317.
- CORRIPIO, I., FERREIRA, A., PORTELLA, M. J., PÉREZ, V., ESCARTÍ, M. J., DEL VALLE CAMACHO, M., y cols. (2012). The role of striatal dopamine D2 receptors in the occurrence of extrapyramidal side effects: iodine-123-iodobenzamide single photon emission computed tomography study. *Psychiatry Research*, 201(1), 73-77.
- DAMOISEAUX, J. S., y GREICIUS, M. D. (2009). Greater than the sum of its parts: a review of studies combining structural connectivity and resting-state functional connectivity. *Brain Structure y Function*, 213(6), 525-533.
- DANIELSEN, E. R., y ROSS, B. (1999). *Magnetic Resonance Spectroscopy Diagnosis of neurological Diseases* (pp. 1-43). New York: Marcel Dekker.
- DE LUCA, MARILENA, SMITH, S., DE STEFANO, N., FEDERICO, A., y MATTHEWS, P. M. (2005). Blood oxygenation level dependent contrast resting state networks are relevant to functional activity in the neocortical sensorimotor system. *Experimental Brain Research*, 167(4), 587-594.
- DECO, G., y CORBETTA, M. (2011). The dynamical balance of the brain at rest. *The Neuroscientist: A Review Journal Bringing Neurobiology, Neurology and Psychiatry*, 17(1), 107-123.
- DI MARTINO, A., SCHERES, A., MARGULIES, D. S., KELLY, A. M. C., UDDIN, L. Q., SHEHZAD, Z., y cols. (2008). Functional Connectivity of Human Striatum: A Resting State fMRI Study. *Cerebral Cortex*, 18(12), 2735-2747.
- Dickie, D. A., SHENKIN, S. D., ANBLAGAN, D., LEE, J., BLESÁ CABEZ, M., RODRIGUEZ, D., y cols. (2017) Whole Brain Magnetic Resonance Image Atlases: A Systematic Review of Existing Atlases and Caveats for Use in Population Imaging. *Frontiers in Neuroinformatics*, 11, 1.
- DIRNBERGER, G., FULLER, R., FRITH, C., y JAHANSHAH, M. (2014). Neural correlates of executive dysfunction in schizophrenia: failure to modulate brain activity with task demands. *Neuroreport*, 25(16), 1308-1315.
- DOSENBACH, N. U. F., FAIR, D. A., MIEZIN, F. M., COHEN, A. L., WENGER, K. K., DOSENBACH, R. A. T., y cols. (2007). Distinct brain networks for adaptive and stable task control in humans. *Proceedings of the National Academy of Sciences of the United States of America*, 104(26), 11073-11078.
- DUNCAN, N. W., y NORTHOFF, G. (2013). Overview of potential procedural and participant-related confounds for neuroimaging of the resting state. *Journal of Psychiatry y Neuroscience: JPN*, 38(2), 84-96.

- EVANS, A. C., COLLINS, D. L., MILLS, S. R., BROWN, E. D., KELLY, R. L., y PETERS, T. M. (1993). 3D statistical neuroanatomical models from 305 MRI volumes. En 1993 IEEE Conference Record Nuclear Science Symposium and Medical Imaging Conference (pp. 1813-1817). San Francisco, EUA: IEEE Conference Record.
- EVANS, A. C., JANKE, A. L., COLLINS, D. L., y BAILLET, S. (2012). Brain templates and atlases. *NeuroImage*, 62(2), 911-922.
- FARRÁS-PERMANYER, L., GUÀRDIA-OLMOS, J., y PERÓ-CEBOLLERO, M. (2015). Efecto del tamaño kernel en el suavizado de señal BOLD en paradigmas funcionales (RMf). *Escritos de Psicología* 8(1): 21-29.
- FOX, M. D., y RAICHEL, M. E. (2007). Spontaneous fluctuations in brain activity observed with functional magnetic resonance imaging. *Nature Reviews. Neuroscience*, 8(9), 700-711.
- FOX, M. D., SNYDER, A. Z., VINCENT, J. L., y RAICHEL, M. E. (2007). Intrinsic fluctuations within cortical systems account for intertrial variability in human behavior. *Neuron*, 56(1), 171-184.
- FRANKLE, W. G., y LARUELLE, M. (2002). Neuroreceptor imaging in psychiatric disorders. *Annals of Nuclear Medicine*, 16(7), 437-446.
- FRISTON, K. J. Introduction Experimental design and Statistical Parametric Mapping. En Frackowiak, W. P. Richard (ed), Friston, K. (ed), Frith, C. (ed), Dolan, R. (ed), Price, C. (ed), Zeki, S. (ed), Ashburner, J. (ed.) (2003). *Human Brain Function* (2nd Edition). San Diego, CA, EUA: Elsevier Science.
- FRISTON, K. J., BUECHEL, C., FINK, G. R., MORRIS, J., ROLLS, E., y DOLAN, R. J. (1997). Psychophysiological and modulatory interactions in neuroimaging. *NeuroImage*, 6(3), 218-229.
- FUKUNAGA, M., HOROVITZ, S. G., VAN GELDEREN, P., DE ZWART, J. A., JANSMA, J. M., IKONOMIDOU, V. N., y cols. (2006). Large-amplitude, spatially correlated fluctuations in BOLD fMRI signals during extended rest and early sleep stages. *Magnetic Resonance Imaging*, 24(8), 979-992.
- GARCÍA-GARCÍA, I., JURADO, M. Á., GAROLERA, M., SEGURA, B., SALA-LLONCH, R., MARQUÉS-ITURRIA, I., y cols. (2013). Alterations of the salience network in obesity: a resting-state fMRI study. *Human Brain Mapping*, 34(11), 2786-2797.
- GIMÉNEZ, M., GUINEA-IZQUIERDO, A., VILLALTA-GIL, V., MARTÍNEZ-ZALACAÍN, I., SEGALÀS, C., SUBIRÀ, M., y cols. (2017). Brain alterations in low-frequency fluctuations across multiple bands in obsessive compulsive disorder. *Brain Imaging and Behavior*, 11(6), 1690-1706.
- GITELMAN, D. R., PENNY, W. D., ASHBURNER, J., y FRISTON, K. J. (2003). Modeling regional and psychophysiological interactions in fMRI: the importance of hemodynamic deconvolution. *NeuroImage*, 19(1), 200-207.
- GONG, G., ROSA-NETO, P., CARBONELL, F., CHEN, Z. J., HE, Y., y EVANS, A. C. (2009). Age-and gender-related differences in the cortical anatomical network. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 29(50), 15684-15693.
- GREICIUS, M. (2008) Resting-state functional connectivity in neuropsychiatric disorders. *Current Opinion in Neurology*, 21(4), 424-430.
- GREICIUS, M. D., KIVINIEMI, V., TERVONEN, O., VAINIONPÄÄ, V., ALAHUHTA, S., REISS, A. L., y MENON, V. (2008). Persistent default-mode network connectivity during light sedation. *Human Brain Mapping*, 29(7), 839-847.

## *Bibliografía completa*

- GREICIUS, M. D., SRIVASTAVA, G., REISS, A. L., y MENON, V. (2004). Default-mode network activity distinguishes Alzheimer's disease from healthy aging: evidence from functional MRI. *Proceedings of the National Academy of Sciences of the United States of America*, 101(13), 4637-4642.
- GUERRA-CARRILLO, B., MACKEY, A. P., y BUNGE, S. A. (2014). Resting-state fMRI: a window into human brain plasticity. *The Neuroscientist: A Review Journal Bringing Neurobiology, Neurology and Psychiatry*, 20(5), 522-533.
- GUJAR, S. K., MAHESHWARI, S., BJÖRKMANN-BURTSCHER, I., y SUNDGREN, P. C. (2005). Magnetic resonance spectroscopy. *Journal of Neuro-Ophthalmology: The Official Journal of the North American Neuro-Ophthalmology Society*, 25(3), 217-226.
- HABAS, C., KAMDAR, N., NGUYEN, D., PRATER, K., BECKMANN, C. F., MENON, V., y GREICIUS, M. D. (2009). Distinct cerebellar contributions to intrinsic connectivity networks. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 29(26), 8586-8594.
- HANDWERKER, D. A., GONZALEZ-CASTILLO, J., D'ESPOSITO, M., y Bandettini, P. A. (2012). The continuing challenge of understanding and modeling hemodynamic variation in fMRI. *NeuroImage*, 62(2), 1017-1023.
- HARRISON, B. J., PUJOL, J., CONTRERAS-RODRÍGUEZ, O., SORIANO-MAS, C., LÓPEZ-SOLÀ, M., DEUS, J., y cols. (2011). Task-induced deactivation from rest extends beyond the default mode brain network. *PloS One*, 6(7), e22964.
- HAXBY, J. V., HOFFMAN, E. A., y GOBBINI, M. I. (2000). The distributed human neural system for face perception. *Trends in Cognitive Sciences*, 4(6), 223-233.
- HE, Y., WANG, J., WANG, L., CHEN, Z. J., YAN, C., YANG, H., y cols. (2009). Uncovering intrinsic modular organization of spontaneous brain activity in humans. *PloS One*, 4(4), e5226.
- HE, Y., WANG, L., ZANG, Y., TIAN, L., ZHANG, X., LI, K., y JIANG, T. (2007). Regional coherence changes in the early stages of Alzheimer's disease: a combined structural and resting-state functional MRI study. *NeuroImage*, 35(2), 488-500.
- HERRERA, M., NOTARIO, B., BARRIO, M. C., METSCHER, B. D., y MURILLO GONZALEZ, J. (2018). X-ray micro-computed tomography of postmortem brain tissue using potassium dichromate as a contrast agent. *Archives Italiennes De Biologie*, 156(1-2), 48-53.
- HOPTMAN, M. J., ZUO, X.-N., BUTLER, P. D., JAVITT, D. C., D'ANGELO, D., MAURO, C. J., y MILHAM, M. P. (2010). Amplitude of low-frequency oscillations in schizophrenia: a resting state fMRI study. *Schizophrenia Research*, 117(1), 13-20.
- HOU, J., WU, W., LIN, Y., WANG, J., ZHOU, D., GUO, J., y cols. (2012). Localization of cerebral functional deficits in patients with obsessive-compulsive disorder: a resting-state fMRI study. *Journal of Affective Disorders*, 138(3), 313-321.
- HOWE, F. A., BARTON, S. J., CUDLIP, S. A., STUBBS, M., SAUNDERS, D. E., MURPHY, M., y cols. (2003). Metabolic profiles of human brain tumors using quantitative in vivo <sup>1</sup>H magnetic resonance spectroscopy. *Magnetic Resonance in Medicine*, 49(2), 223-232.
- HUMPHRIES, M. D., y GURNEY, K. (2008). Network «small-world-ness»: a quantitative method for determining canonical network equivalence. *PloS One*, 3(4), e0002051.
- JASPER, H. H. (1958). Report of the committee on methods of clinical examination in electroencephalography. *Electroencephalography and Clinical Neurophysiology*, 10(2): 370-375.

- JEZZARD, P., y CLARE, S. (1999). Sources of distortion in functional MRI data. *Human Brain Mapping*, 8(2-3), 80-85.
- JIANG, L., y Zuo, X.-N. (2016). Regional Homogeneity: A Multimodal, Multiscale Neuroimaging Marker of the Human Connectome. *The Neuroscientist: A Review Journal Bringing Neurobiology, Neurology and Psychiatry*, 22(5), 486-505.
- KAHNT, T., CHANG, L. J., PARK, S. Q., HEINZLE, J., y HAYNES, J.-D. (2012). Connectivity-based parcellation of the human orbitofrontal cortex. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 32(18), 6240-6250.
- KARBOWSKI, J. (2007). Global and regional brain metabolic scaling and its functional consequences. *BMC Biology*, 5, 18.
- (2011). Scaling of brain metabolism and blood flow in relation to capillary and neural scaling. *PloS One*, 6(10), e26709.
- KASSENBRÖCK, A., VASDEV, N., y LIANG, S. H. (2016). Selected PET Radioligands for Ion Channel Linked Neuroreceptor Imaging: Focus on GABA, NMDA and nACh Receptors. *Current Topics in Medicinal Chemistry*, 16(16), 1830-1842.
- KINOSHITA, Y., KAJIWARA, H., YOKOTA, A., y KOGA, Y. (1994). Proton magnetic resonance spectroscopy of brain tumors: an in vitro study. *Neurosurgery*, 35(4), 606-613; discussion 613-614.
- KONDO, A., ISHII, H., AOKI, S., SUZUKI, M., NAGASAWA, H., KUBOTA, K., y cols. (2016). Phase IIa clinical study of [<sup>18</sup>F]fluciclovine: efficacy and safety of a new PET tracer for brain tumors. *Annals of Nuclear Medicine*, 30(9), 608-618.
- KÖNIGS, M., VAN HEURN, L. W. E., BAKX, R., VERMEULEN, R. J., GOSLINGS, J. C., POLL-THE, B. T., y cols. (2017). The structural connectome of children with traumatic brain injury. *Human Brain Mapping*.
- KOYAMA, M. S., KELLY, C., SHEHZAD, Z., PENESETTI, D., CASTELLANOS, F. X., y MILHAM, M. P. (2010). Reading networks at rest. *Cerebral Cortex*, 20(11), 2549-2559.
- KÜBLBÖCK, M., WOLETZ, M., HÖFLICH, A., SLADKY, R., KRANZ, G. S., HOFFMANN, A., y cols. (2014). Stability of low-frequency fluctuation amplitudes in prolonged resting-state fMRI. *NeuroImage*, 103, 249-257.
- LASCANO, A. M., LALIVE, P. H., HARDMEIER, M., FUHR, P., y SEECK, M. (2017). Clinical evoked potentials in neurology: a review of techniques and indications. *Journal of Neurology, Neurosurgery and Psychiatry*, 88(8):688-696.
- LEWIS, C. M., BALDASSARRE, A., COMMITTERI, G., ROMANI, G. L., y CORBETTA, M. (2009). Learning sculpts the spontaneous activity of the resting human brain. *Proceedings of the National Academy of Sciences of the United States of America*, 106(41), 17558-17563.
- LI, Y., LIU, Y., LI, J., QIN, W., LI, K., YU, C., y JIANG, T. (2009). Brain anatomical network and intelligence. *PLoS Computational Biology*, 5(5), e1000395.
- LIU, C.-H., MA, X., WU, X., LI, F., ZHANG, Y., ZHOU, F.-C., y cols. (2012). Resting-state abnormal baseline brain activity in unipolar and bipolar depression. *Neuroscience Letters*, 516(2), 202-206.
- LONG, X.-Y., ZUO, X.-N., KIVINIEMI, V., YANG, Y., ZOU, Q.-H., ZHU, C.-Z., y cols. (2008). Default mode network as revealed with multiple methods for resting-state functional MRI analysis. *Journal of Neuroscience Methods*, 171(2), 349-355.

## *Bibliografía completa*

- MANDAL, P. K., MAHAJAN, R., y DINOVA, I. D. (2012). Structural brain atlases: design, rationale, and applications in normal and pathological cohorts. *Journal of Alzheimer's Disease: JAD*, 31 Suppl 3, S169-188.
- MARGULIES, D. S., BÖTTGER, J., LONG, X., LV, Y., KELLY, C., SCHÄFER, A., y cols. (2010). Resting developments: a review of fMRI post-processing methodologies for spontaneous brain activity. *Magma*, 23(5-6), 289-307.
- MATTHEWS, P. M., RABINER, E. A., PASSCHIER, J., y GUNN, R. N. (2012). Positron emission tomography molecular imaging for drug development. *British Journal of Clinical Pharmacology*, 73(2), 175-186.
- MEARS, D., y POLLARD, H. B. (2016). Network science and the human brain: Using graph theory to understand the brain and one of its hubs, the amygdala, in health and disease. *Journal of Neuroscience Research*, 94(6), 590-605.
- MEUNIER, D., LAMBIOTTE, R., FORNITO, A., ERSCHKE, K. D., y BULLMORE, E. T. (2009). Hierarchical modularity in human brain functional networks. *Frontiers in Neuroinformatics*, 3, 37.
- MICHELOYANNIS, S., VOURKAS, M., TSIRKA, V., KARAKONSTANTAKI, E., KANATSOULI, K., y STAM, C. J. (2009). The influence of ageing on complex brain networks: a graph theoretical analysis. *Human Brain Mapping*, 30(1), 200-208.
- MONTÉ-RUBIO, G. C., FALCÓN, C., POMAROL-CLOTET, E., ASHBURNER, J. (2018). A comparison of various MRI feature types for characterizing whole brain anatomical differences using linear pattern recognition methods. *Neuroimage*, 178:753-768.
- MOURÃO-MIRANDA, J., HARDOON, D.R., HAHN, T., MARQUAND, A.F., WILLIAMS, S.C., SHAWE-TAYLOR, J. y cols. (2011). Patient classification as an outlier detection problem: an application of the One-Class Support Vector Machine. *Neuroimage*, 58(3):793-804.
- MURPHY, M., LOOSEMORE, A., CLIFTON, A. G., HOWE, F. A., TATE, A. R., CUDLIP, S. A., y cols. (2002). The contribution of proton magnetic resonance spectroscopy (1H MRS) to clinical brain tumour diagnosis. *British Journal of Neurosurgery*, 16(4), 329-334.
- NUGENT, A. C., MARTINEZ, A., D'ALFONSO, A., ZARATE, C. A., y THEODORE, W. H. (2015). The relationship between glucose metabolism, resting-state fMRI BOLD signal, and GABAA-binding potential: a preliminary study in healthy subjects and those with temporal lobe epilepsy. *Journal of Cerebral Blood Flow and Metabolism: Official Journal of the International Society of Cerebral Blood Flow and Metabolism*, 35(4), 583-591.
- PARKES, L., FULCHER, B., YÜCEL, M., y FORNITO, A. (2018). An evaluation of the efficacy, reliability, and sensitivity of motion correction strategies for resting-state functional MRI. *NeuroImage*, 171, 415-436.
- PARRADO-HERNÁNDEZ, E., GÓMEZ-VERDEJO, V., MARTÍNEZ-RAMÓN, M., SHAWE-TAYLOR, J., ALONSO, P., PUJOL, J. y cols. (2014). Discovering brain regions relevant to obsessive-compulsive disorder identification through bagging and transduction. *Med Image Anal*, 18(3):435-448.
- PASTOR, J., y SOLA, R. G. (2002). Biophysical foundations of magnetoencephalography. *Revista de Neurología*, 34, 843-850.
- PETERSEN, S. E., y DUBIS, J. W. (2012). The mixed block/event-related design. *NeuroImage*, 62(2), 1177-1184.
- POLDRACK, R. A. (2007). Region of interest analysis for fMRI. *Social Cognitive and Affective Neuroscience*, 2(1), 67-70.

- POWER, J. D., MITRA, A., LAUMANN, T. O., SNYDER, A. Z., SCHLAGGAR, B. L., y PETERSEN, S. E. (2014) Methods to detect, characterize, and remove motion artifact in resting state fMRI. *NeuroImage*, 84, 320-341.
- PTAK, R. (2012) The frontoparietal attention network of the human brain: action, saliency, and a priority map of the environment. *The Neuroscientist: A Review Journal Bringing Neurobiology, Neurology and Psychiatry*, 18(5), 502-515.
- PUJOL, J., BATALLA, I., CONTRERAS-RODRÍGUEZ, O., HARRISON, B. J., PERA, V., HERNÁNDEZ-RIBAS, R., y cols. (2012). Breakdown in the brain network subserving moral judgment in criminal psychopathy. *Social Cognitive and Affective Neuroscience*, 7(8), 917-923.
- PUJOL, JESÚS, SORIANO-MAS, C., ALONSO, P., CARDONER, N., MENCHÓN, J. M., DEUS, J., y VALLEJO, J. (2004) Mapping structural brain alterations in obsessive-compulsive disorder. *Archives of General Psychiatry*, 61(7), 720-730.
- RADEMACHER, J., MOROSAN, P., SCHORMANN, T., SCHLEICHER, A., WERNER, C., FREUND, H. J., y ZILLES, K. (2001). Probabilistic mapping and volume measurement of human primary auditory cortex. *NeuroImage*, 13(4), 669-683.
- RAICHLER, M. E. (2015). The brain's default mode network. *Annual Review of Neuroscience*, 38, 433-447.
- RAICHLER, M. E., y MINTUN, M. A. (2006). Brain work and brain imaging. *Annual Review of Neuroscience*, 29, 449-476.
- RECHTSCHAFFEN, A. (ed), KALES, A. (ed) (1968). A manual of standardized terminology, techniques and scoring system of sleep stages in human subjects. Los Angeles, EUA: Brain Information Service/Brain Research Institute, University of California.
- ROSAZZA, C., y MINATI, L. (2011). Resting-state brain networks: literature review and clinical applications. *Neurological Sciences: Official Journal of the Italian Neurological Society and of the Italian Society of Clinical Neurophysiology*, 32(5), 773-785.
- ROWLAND, L. M., SHADMEHR, R., KRAVITZ, D., y HOLCOMB, H. H. (2008). Sequential neural changes during motor learning in schizophrenia. *Psychiatry Research*, 163(1), 1-12.
- RUBINOV, M., y SPORNS, O. (2010). Complex network measures of brain connectivity: uses and interpretations. *NeuroImage*, 52(3), 1059-1069.
- SALA-LLONCH, R., BARTRÉS-FAZ, D., y JUNQUÉ, C. (2015). Reorganization of brain networks in aging: a review of functional connectivity studies. *Frontiers in Psychology*, 6, 663.
- SALIBI, N., y BROW, M. (1998). *Clinical MR Spectroscopy- First principles* (pp 62-99, 143-146). New York: Wiley-Liss.
- SALVADOR, R., RADUA, J., CANALES-RODRÍGUEZ, E. J., SOLANES, A., SARRÓ, S., GOIKOLEA, J. M., y cols. (2017). Evaluation of machine learning algorithms and structural features for optimal MRI-based diagnostic prediction in psychosis. *PLoS One*, 12(4):e0175683.
- SAVOY, R. L., BANDETTINI, P. A., O'CRIVEN, K. M., KWONG, K. K., DAVIS, T. L., BAKER, J. R., y cols. (1995). Pushing the temporal resolution of fMRI: studies of very brief visual stimuli, onset variability and asynchrony, and stimulus-correlated changes in noise. *Proceedings of the Society for Magnetic Resonance Medicine*, 3:450.
- SCHMITHORST, V. J., y HOLLAND, S. K. (2004). Comparison of three methods for generating group statistical inferences from independent component analysis of functional magnetic resonance imaging data. *Journal of Magnetic Resonance Imaging: JMIR*, 19(3), 365-368.



*Bibliografía completa*

- SCHMITT, J. E., LENROOT, R. K., WALLACE, G. L., ORDAZ, S., TAYLOR, K. N., KABANI, N., y cols. (2008). Identification of genetically mediated cortical networks: a multivariate study of pediatric twins and siblings. *Cerebral Cortex*, 18(8), 1737-1747.
- SEELEY, W. W., MENON, V., SCHATZBERG, A. F., KELLER, J., GLOVER, G. H., KENNA, H., y cols. (2007). Dissociable intrinsic connectivity networks for salience processing and executive control. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 27(9), 2349-2356.
- SETO, E., SELA, G., MCLROY, W. E., BLACK, S. E., STAINES, W. R., BRONSKILL, M. J., y cols. (2001). Quantifying head motion associated with motor tasks used in fMRI. *NeuroImage*, 14(2), 284-297.
- SEVEL, L.S., BOISSONEAULT, J., LETZEN, J.E., ROBINSON, M.E., STAUD, R. (2018). Structural brain changes versus self-report: machine-learning classification of chronic fatigue syndrome patients. *Exp Brain Res*, 236(8):2245-2253.
- SHANNON, B. J., DOSENBACH, R. A., SU, Y., VLASSENKO, A. G., LARSON-PRIOR, L. J., NOLAN, T. S., y cols. (2013). Morning-evening variation in human brain metabolism and memory circuits. *Journal of Neurophysiology*, 109(5), 1444-1456.
- SHIRAHATA, N., HENRIKSEN, L., VORSTRUP, S., HOLM, S., LAURITZEN, M., PAULSON, O. B., y LASSEN, N. A. (1985) Regional cerebral blood flow assessed by <sup>133</sup>Xe inhalation and emission tomography: normal values. *Journal of Computer Assisted Tomography*, 9(5), 861-866.
- SHMUEL, A., y LEOPOLD, D. A. (2008). Neuronal correlates of spontaneous fluctuations in fMRI signals in monkey visual cortex: Implications for functional connectivity at rest. *Human Brain Mapping*, 29(7), 751-761.
- SHUKE, N., HASHIZUME, K., KIRIYAMA, K., OKIZAKI, A., YAMAMOTO, W., ZHAO, C., y cols. (2004). Correct localization of epileptogenic focus with I-123 iomazenil cerebral benzodiazepine receptor imaging: a case report of temporal lobe epilepsy with discordant ictal cerebral blood flow SPECT. *Annals of Nuclear Medicine*, 18(6), 541-545.
- SLAGTER, H. A., TOMER, R., CHRISTIAN, B. T., FOX, A. S., COLZATO, L. S., KING, C. R., y cols. (2012). PET evidence for a role for striatal dopamine in the attentional blink: functional implications. *Journal of Cognitive Neuroscience*, 24(9), 1932-1940.
- SMITH, A. M., LEWIS, B. K., RUTTIMANN, U. E., YE, F. Q., SINNWELL, T. M., YANG, Y., y cols. (1999). Investigation of low frequency drift in fMRI signal. *NeuroImage*, 9(5), 526-533.
- SMITH, S. M., MILLER, K. L., SALIMI-KHORSHIDI, G., WEBSTER, M., BECKMANN, C. F., NICHOLS, T. E., y cols. (2011). Network modelling methods for FMRI. *NeuroImage*, 54(2), 875-891.
- SOHN, W. S., YOO, K., LEE, Y.-B., SEO, S. W., NA, D. L., y JEONG, Y. (2015). Influence of ROI selection on resting state functional connectivity: an individualized approach for resting state fMRI analysis. *Frontiers in Neuroscience*, 9, 280.
- SONG, X., ZHANG, Y., y LIU, Y. (2014). Frequency specificity of regional homogeneity in the resting-state human brain. *PloS One*, 9(1), e86818.
- SONG, X.-W., DONG, Z.-Y., LONG, X.-Y., LI, S.-F., ZUO, X.-N., ZHU, C.-Z., y cols. (2011). REST: a toolkit for resting-state functional magnetic resonance imaging data processing. *PloS One*, 6(9), e25031.
- SORIANO-MAS, C., PUJOL, J., ALONSO, P., CARDONER, N., MENCHÓN, J.M., HARRISON, B.J., y cols. (2007) Identifying patients with obsessive-compulsive disorder using whole-brain anatomy. *Neuroimage*, 35(3):1028-1037.

- STEJSKAL, E. O., y TANNER, J. E. (1965). Spin diffusion measurements: spin echoes in the presence of a time-dependent field gradient. *The Journal of Chemical Physics*, 42, 288-292.
- STONE, J. V. (2002). Independent component analysis: an introduction. *Trends in Cognitive Sciences*, 6(2), 59-64.
- TALAIRACH, J., TOURNOUX, P. (1988). *Co-planar stereotaxic atlas of the human brain*. (1st ed.). New York, USA: Thieme.
- TANG, Y., GE, J., LIU, F., WU, P., GUO, S., LIU, Z., y cols. (2016). Cerebral Metabolic Differences Associated with Cognitive Impairment in Parkinson's Disease. *PLoS One*, 11(4), e0152716.
- TASCHLER, B., GE, T., BENDFELDT, K., MÜLLER-LENKE, N., JOHNSON, TD., NICHOLS, TE. (2014). Spatial modeling of multiple sclerosis for disease subtype prediction.. *Med Image Comput Comput Assist Interv*, 17(Pt 2):797-804.
- TOHKA, J., MORADI, E., HUTTUNEN, H., Alzheimer's Disease Neuroimaging Initiative (2016). Comparison of Feature Selection Techniques in Machine Learning for Anatomical Brain MRI in Dementia. *Neuroinformatics*, 14(3):279-296.
- VAIDYA, C. J., y GORDON, E. M. (2013). Phenotypic variability in resting-state functional connectivity: current status. *Brain Connectivity*, 3(2), 99-120.
- VAN DE GIESSEN, E., CELIK, F., SCHWEITZER, D. H., VAN DEN BRINK, W., y BOOU, J. (2014). Dopamine D2/3 receptor availability and amphetamine-induced dopamine release in obesity. *Journal of Psychopharmacology (Oxford, England)*, 28(9), 866-873.
- VAN DIJK, K. R. A., SABUNCU, M. R., y BUCKNER, R. L. (2012). The influence of head motion on intrinsic functional connectivity MRI. *NeuroImage*, 59(1), 431-438.
- VISSCHER, K. M., MIEZIN, F. M., KELLY, J. E., BUCKNER, R. L., DONALDSON, D. I., McAVOY, M. P., y cols. (2003). Mixed blocked/event-related designs separate transient and sustained activity in fMRI. *NeuroImage*, 19(4), 1694-1708.
- VIVIANI, R., GRÖN, G., y SPITZER, M. (2005). Functional principal component analysis of fMRI data. *Human Brain Mapping*, 24(2), 109-129.
- WAGNER, H. N. (1998). A brief history of positron emission tomography (PET). *Seminars in Nuclear Medicine*, 28(3), 213-220.
- WICHMANN, E. H. (1996). *Física cuántica* (pp 324-326). Barcelona: Reverté.
- WULFF, S., PINBORG, L. H., SVARER, C., JENSEN, L. T., NIELSEN, M. Ø., ALLERUP, P., y cols. (2015). Striatal D(2/3) Binding Potential Values in Drug-Naïve First-Episode Schizophrenia Patients Correlate With Treatment Outcome. *Schizophrenia Bulletin*, 41(5), 1143-1152.
- YAN, C., LIU, D., HE, Y., ZOU, Q., ZHU, C., ZUO, X., y cols. (2009). Spontaneous brain activity in the default mode network is sensitive to different resting-state conditions with limited cognitive load. *PLoS One*, 4(5), e5743.
- YUAN, R., DI, X., KIM, E. H., BARIK, S., RYPMA, B., y BISWAL, B. B. (2013). Regional homogeneity of resting-state fMRI contributes to both neurovascular and task activation variations. *Magnetic Resonance Imaging*, 31(9), 1492-1500.
- ZALESKY, A., FORNITO, A., HARDING, I. H., COCCHI, L., YÜCEL, M., PANTELIS, C., y BULLMORE, E. T. (2010). Whole-brain anatomical networks: does the choice of nodes matter? *NeuroImage*, 50(3), 970-983.
- ZANG, Y., JIANG, T., LU, Y., HE, Y., y TIAN, L. (2004) Regional homogeneity approach to fMRI data analysis. *NeuroImage*, 22(1), 394-400.

*Bibliografía completa*

- ZANG, Y.-F., HE, Y., ZHU, C.-Z., CAO, Q.-J., SUI, M.-Q., LIANG, M., y cols. (2007) Altered baseline brain activity in children with ADHD revealed by resting-state functional MRI. *Brain y Development*, 29(2), 83-91.
- ZHANG, D., SNYDER, A. Z., FOX, M. D., SANSBURY, M. W., SHIMONY, J. S., y RAICHLER, M. E. (2008). Intrinsic functional relations between human cerebral cortex and thalamus. *Journal of Neurophysiology*, 100(4), 1740-1748.
- ZHONG, Y., WANG, H., LU, G., ZHANG, Z., JIAO, Q., y LIU, Y. (2009). Detecting functional connectivity in fMRI using PCA and regression analysis. *Brain Topography*, 22(2), 134-144.
- ZOU, Q., WU, C. W., STEIN, E. A., ZANG, Y., y YANG, Y. (2009). Static and dynamic characteristics of cerebral blood flow during the resting state. *NeuroImage*, 48(3), 515-524.
- ZOU, Q.-H., ZHU, C.-Z., YANG, Y., ZUO, X.-N., LONG, X.-Y., CAO, Q.-J., y cols. (2008). An improved approach to detection of amplitude of low-frequency fluctuation (ALFF) for resting-state fMRI: fractional ALFF. *Journal of Neuroscience Methods*, 172(1), 137-141.
- ZUO, X.-N., DI MARTINO, A., KELLY, C., SHEHZAD, Z. E., GEE, D. G., KLEIN, D. F., y cols. (2010). The oscillating brain: complex and reliable. *NeuroImage*, 49(2), 1432-1445.