

Bibliografía web

- ACHIRON A., CHAPMAN J., MAGALASHVILI D., DOLEV M., LAVIE M., BERCOVICH E., POLLIACK M., DONIGER G. M., STERN Y., KHILKEVICH O., MENASCU S., HARARAI G., GUREVICH M., BARAK Y. Modeling of cognitive impairment by disease duration in multiple sclerosis: a cross-sectional study. *PLoSOne*. 2013 Aug 1; 8(8): e71058.
- AGOSTA F., FILIPPI M. MRI of spinal cord in multiple sclerosis. *J Neuroimaging* 2007; Suppl 1: S46-49.
- AGOSTA F., PAGANI E., CAPUTO D., FILIPPI M. Associations between cervical cord gray matter damage and disability in patients with multiple sclerosis. *Arch Neurol*. 2007; 64(9): 1302-5.
- AKBAR N., HONARMAND K., KOU N., LEVINE B., RECTOR N., FEINSTEIN A. Validity of an Internet version of the Multiple Sclerosis Neuropsychological Questionnaire. *Mult Scler*. 2010 Dec; 16(12):1500-6. doi: 10.1177/1352458510379615. Epub 2010 Sep 2.
- AKHONDZADEH S., TAHMACEBI-POUR N., NOORBALA A. A., AMINI H., FALLAH-POUR H., JAMSHIDI A. H., *et al*. Crocus sativus L. in the treatment of mild to moderate depression: A double-blind, randomized and placebo-controlled trial. *Phytother Res*. 2005; 19:148-51.
- ALONSO J., PRIETO L., FERRER M., BROQUETAS J. M., ROCA J., SERRA-BATLLES J., *et al*. Testing the Spanish version of the SF-36 Health Survey among male patients with chronic obstructive pulmonary disease. *J Clin Epidemiol*. 1998; 46:373-81.
- ALONSO J., REGIDOR E., BARRIO G., PRIETO L., RODRÍGUEZ C. y DE LA FUENTE L. Valores poblacionales de referencia de la versión española del Cuestionario de Salud SF-36. *Med Clin Barc*. 1998; 111:410-416.
- ALONSO J., ANTO J. M., MORENO C. Spanish version of the Nottingham Health Profile: translation and preliminary validity. *Am J Public Health*. 1990 Jun; 80(6):704-8.
- ALTENMALLER E. O., MÜNTE T. F., GERLOFF C. Neurocognitive functions and the EEG. En: NIEDERMEYER, E., LOPES DA SILVA, F. H., eds. *Electroencephalography: Basic principles, clinical applications, and related fields* (5th ed.). Philadelphia: Lippincott Williams & Wilkins, 2005.
- AMANN M., DÖSSEGGER L. S., PENNER I. K., HIRSCH J. G., RASELLI C., CALABRESE P., WEIER K., RADÜ E. W., KAPPOS L., GASS A. Altered functional adaptation to attention and working memory tasks with increasing complexity in relapsing-remitting multiple sclerosis patients. *Hum Brain Mapp* 2011; 32(10):1704-19.

- AMATO M. P., BARTOLOZZI M. L., ZIPOLI V., PORTACCIO E., MORTILLA M., GUIDI L., SIRACUSA G., SORBI S., FEDERICO A., DE STEFANO N. Neocortical volume decrease in relapsing-remitting MS patients with mild cognitive impairment. *Neurology* 2004; 63:89-93.
- AMATO M. P., GORETTI B., GHEZZI A., HAKIKI B., NICCOLAI C., LORI S., MOIOLA L., FALAUTANO M., VITERBO R. G., PATTI F., CILIA S., POZZILLI C., BIANCHI V., ROSCIO M., MARTINELLI V., COMI G., PORTACCIO E., TROJANO M.; MS Study Group of the Italian Neurological Society. Neuropsychological features in childhood and juvenile multiple sclerosis: five-year follow-up. *Neurology* (2014b) Oct 14; 83(16):1432-8.
- AMATO M. P., GORETTI B., VITERBO R. G., PORTACCIO E., NICCOLAI C., HAKIKI B., *et al.* Computer assisted rehabilitation of attention in patients with multiple sclerosis: results of a randomized, double-blind trial. *Mult Scler* 2014a; 20(1):91-8.
- AMATO M. P., LANGDON D., MONTALBAN X., BENEDICT R. H., DE LUCA J., KRUPP L. B., THOMPSON A. J., COMI G. Treatment of cognitive impairment in multiple sclerosis: position paper. *J Neurol.* 2013 Jun; 260(6):1452-68.
- AMATO M. P., PONZIANI G., SIRACUSA G., SORBI S. Cognitive dysfunction in early-onset multiple sclerosis: a reappraisal after 10 years. *Arch Neurol.* 2001; 58(10):1602-6.
- AMATO M. P., RAZZOLINI L., GORETTI B., STROMILLO M. L., ROSSI F., GIORGIO A., HAKIKI B., GIANNINI M., PASTÒ L., PORTACCIO E., DE STEFANO N. Cognitive reserve and cortical atrophy in multiple sclerosis: a longitudinal study. *Neurology* 2013 May 7; 80(19):1728-33.
- AMINOFF J. C., GOODIN D. S. Long-latency cerebral event-related potentials in multiple sclerosis. *J Clin Neurophysiol.* 2001; 18(4): 372-7.
- ANDREU-CATALÀ M., PASCUAL-LOZANO A. M. Afectación de las funciones cognitivas en la esclerosis múltiple secundaria progresiva. *Rev Neurol.* 2008; 46(11):664-666
- ARANGO-LASPRILLA J. C., DELUCA J., CHIARAVALLOTI N. El perfil neuropsicológico en la esclerosis múltiple. *Psicothema*, 2007;19:1-6.
- ARCHIBALD C. J., FISK J. D. Information processing efficiency in patients with multiple sclerosis. *J Clin Exp Neuropsychol.* 2000; 22:686-701.
- ARNETT P. Neuropsicología en esclerosis múltiple. En: FERNANDEZ O., ed. Cuadernos de esclerosis múltiple. Madrid: Alcandora Publicidad S. L., 2004.
- ARNETT P. A., BARWICK F. H., BEENEY J. E. Depression in multiple sclerosis: review and theoretical proposal. *J Int Neuropsychol Soc.* 2008; 14:691-724.
- ARNETT P. A., RAO S. M., BERNARDIN L., GRAFMAN J., YETKIN F. Z., LOBECK L. Relationship between frontal lobe lesions and Wisconsin Card Sorting Test performance in patients with multiple sclerosis. *Neurology* 1994; 44:420-5.
- ARNETT P., FRÍAS C. F. Evaluación neuropsicológica en la esclerosis múltiple. *Revista de neurología* 2007; 44(3):166-172.
- ARNOLD D. L., MATTHEWS P. M., FRANCIS G. S., *et al.*, Proton magnetic resonance spectroscopic imaging for metabolic characterization of demyelinating plaques. *Ann Neurol.* 1992; 31:235-41.
- ASCHERIO A., MUNGER K. L. Environmental risk factors for multiple sclerosis. Part I: the role of infection. *Ann Neurol.* 2007; 61(4):288-99.
- AUDOIN B., IBARROLA D., RANJEVA J. P., CONFORT-GOUNY S., MALIKOVA I., ALI-CHÉRIF A., PELLETIER J., COZZONE P. Compensatory cortical activation observed by fMRI during a cognitive task at the earliest stage of MS. *Hum Brain Mapp.* 2003; 20(2):51-8.

- AUDOIN B., MY VAN A. D., RANJEVA J. P., IBARROLA D., MALIKOVA I., CONFORT GOUNY S., SOULIER E., VIOUT P., ALI-CHERIF A., PELLETIER J., COZZONE P. J., *et al.*, Magnetic resonance study of the influence of tissue damage and cortical reorganization on PASAT performance at the earliest stage of multiple sclerosis. *Hum Brain Mapp* 2005; 24:216-28.
- AUPPERLE R. L., BEATTY W. W., DE NAP S. F., GONTKOVSKY S. T. Three screening batteries to detect cognitive impairment in multiple sclerosis. *Mult Scler.* 2002; 8:382-9. doi: 10.1191/1352458502ms832oa.
- AYMERICH M., GUILLAMÓN I., PERKAL H., NOS C., PORCEL J., BERRA S., RAJMIL L., MONTALBÁN X. Spanish adaptation of the disease-specific questionnaire MSQOL-54 in multiple sclerosis patients. *Neurologia* 2006 May; 21(4):181-7.
- BACKUS D., MANELLA C., BENDER A., SWEATMAN M. Impact of Massage Therapy on Fatigue, Pain, and Spasticity in People with Multiple Sclerosis: Pilot Study. *IntTher Massage Bodywork* 2016 Dec 9; 9(4):4-13.
- BADDELEY A. D., BADDELEY H. A., BUCKS R. S., WILCOCK G. K. Attentional control in Alzheimer's disease. *Brain* 2001; 124:1492-1508.
- BADÍA X., ALONSO J. La medida de la salud. Guía de escalas de medición en español. 4.^a edición. Barcelona: Lilly, 2007.
- BADÍA X., LIZÁN L. Estudios de calidad de vida. Cuadernos de Atención Primaria. Barcelona: Ser-vet Catalá de Salut, 2005.
- BAGNATO F., JEFFRIES N., RICHERT N. D., STONE R. D., OHAYON J. M., MCFARLAND H. F., FRANK J. A. Evolution of T1 black holes in patients with multiple sclerosis imaged monthly for 4 years. *Brain* 2003; 26(Pt 8):1782-9.
- BAKSHI R., DANDAMUDI V. S., NEEMA M., DE C., BERMEL R. A. Measurement of brain and spinal cord atrophy by magnetic resonance imaging as a tool to monitor multiple sclerosis. *J Neuro-imaging* 2005; 15(4 Suppl):S30-45.
- BAKSHI R., DMOCHOWSKI J., SHAIKH Z. A., JACOBS L. Gray matter T2 hypointensity is related to plaques and atrophy in the brains of multiple sclerosis patients. *J Neurol Sci.* 2001; 185(1):19-26.
- BAKSHI R., HUTTON G. J., MILLER J. R., RADUE E. W. The use of magnetic resonance imaging in the diagnosis and long-term management of multiple sclerosis. *Neurology* 2004; 63(11 Suppl 5):S3-S11.
- BALTRUSCHAT S. A., VENTURA-CAMPOS N., CRUZ-GÓMEZ Á. J., BELENGUER A., FORN C. Gray matter atrophy is associated with functional connectivity reorganization during the Paced Auditory Serial Addition Test (PASAT) execution in Multiple Sclerosis. *J Neuroradiol.* 2015; 42(3):141-9.
- BARAK Y., ACHIRON A. Effect of interferon-beta-1b on cognitive functions in multiple sclerosis. *Eur Neurol* 2002;47:11-14.
- BARCO A. N., LÓPEZ M. P. S., RIBAL J. B., PÉREZ T. O., PÉREZ M. Á. H. Alteraciones cognitivas en etapas iniciales de la esclerosis múltiple y su relación con el estado de ánimo, variables demográficas y clínicas. *Psicothema* 2008; 20:583-588.
- BARKHOF F., VAN WALDERVEEN M. Characterization of tissue damage in multiple sclerosis by nuclear magnetic resonance. *Philos Trans R Soc Lond B Biol Sci.* 1999; 354(1390):1675-86.

- BASSO M. R., LOWERY N., GHORMLEY C., COMBS D., JOHNSON J. Self-generated learning in people with multiple sclerosis. *J Int Neuropsychol Soc.* 2006; 12:640-8.
- BAUMSTARCK K., BOYER., BOUCEKINE M., MICHEL P., PELLETIER J. measuring the quality of life in patients with multiple sclerosis in clinical practice: A necessary challenge. *Mult Scler Int.* 2013; 2013:524894.
- BAUMSTARCK-BARRAU K., SIMEONI M. C., REUTER F. Cognitive function and quality of life in multiple sclerosis patients: a cross-sectional study. *BMC Neurol.* 2011; 11:17.
- BEATTY W. W., PAUL R. H., WILBANKS S. L., HAMES K. A., BLANCO C. R., GOODKIN D. E. Identifying multiple sclerosis patients with mild or global cognitive impairment using the Screening Examination for Cognitive Impairment (SEFCI). *Neurology* 1995 Apr; 45(4):718-23
- BEIER M., BOMBARDIER C. H., HARTOONIAN N., MOTL R. W., KRAFT G. H. Improved physical fitness correlates with improved cognition in multiple sclerosis. *Arch Phys Med Rehabil.* 2014 Jul; 95(7):1328-34.
- BENEDICT R. H., AMATO M. P., BORINGA J., BROCHET B., FOLEY F., FREDRIKSON S., HAMALAINEN P., HARTUNG H., KRUPP L., PENNER I., REDER A. T., LANGDON D. Brief International Cognitive Assessment for MS (BICAMS): international standards for validation. *BMC Neurol.* 2012 Jul; 16:12:55. doi: 10.1186/1471-2377-12-55.
- BENEDICT R. H., BAKSHI R., SIMON J. H., PRIORE R., MILLER C., MUNSCHAUER F. Frontal cortex atrophy predicts cognitive impairment in multiple sclerosis. *J Neuropsychiatry Clin Neurosci.* 2002; 14(1):44-51.
- BENEDICT R. H., BRUCE J. M., DWYER M. G., ABDELRAHMAN N., HUSSEIN S., WEINSTOCK-GUTTMAN B., GARG N., MUNSCHAUER F., ZIVADINOV R. Neocortical atrophy, third ventricular width, and cognitive dysfunction in multiple sclerosis. *Arch Neurol.* 2006; 63(9):1301-6.
- BENEDICT R. H., COX D., THOMPSON L. L., FOLEY F., WEIN-STOCK-GUTTMAN B., MUNSCHAUER F. Reliable screen- ing for neuropsychological impairment in multiple sclerosis. *Mult Scler.* 2004 Dec; 10(6):675-678.
- BENEDICT R. H., MORROW S. A., WEINSTOCK GUTTMAN B., *et al.*, Cognitive reserve moderates decline in information processing speed in multiple sclerosis patients. *J Int Neuropsychol Soc.* 2010; 16:829-835.
- BENEDICT R. H., MUNSCHAUER F., LINN R., MILLER C., MURPHY E., FOLEY F., JACOBS L. Screening for multiple sclerosis cognitive impairment using a self-administered 15-item questionnaire. *Mult Scler.* 2003 Feb; 9(1):95-101.
- BENEDICT R. H., MUNSCHAUER F., ZAREVICS P., ERLANGER D., ROWE V., FEASTER T., CARPENTER R. L. Effects of l-amphetamine sulfate on cognitive function in multiple sclerosis patients. *J Neurol.* 2008 Jun; 255(6):848-52.
- BENEDICT R. H., SMERBECK A., PARIKH R., *et al.*, Reliability and equivalence of alternate forms for the Symbol Digit Modalities Test: Implications for multiple sclerosis clinical trials. *Mult Scler.* 2012; 18(9):1320-1325.
- BENEDICT R. H., WEINSTOCK-GUTTMAN B., FISHMAN I., SHARMA J., TJOA C. W., BAKSHI R. Prediction of neuropsychological impairment in multiple sclerosis: comparison of conventional magnetic resonance imaging measures of atrophy and lesion burden. *Arch Neurol.* 2004; 61:226-30.

- BENEDICT R. H., ZIVADINOV R., CARONE D. A., WEINSTOCK-GUTTMAN B., GAINES J., MAGGIORE C., SHARMA J., TOMASSI M. A., BAKSHI R. Regional lobar atrophy predicts memory impairment in multiple sclerosis. *AJNR Am J Neuroradiol.* 2005; 26(7):1824-31.
- BENEDICT R. H. B., BRUCE J. M., DWYER M. G., ABDELRAHMAN N., HUSSEIN S., WEINSTOCK-GUTTMAN B., *et al.* Neocortical atrophy, third ventricular width, and cognitive dysfunction in multiple sclerosis. *Arch Neurol.* 2006; 63:1301-6.
- BENEDICT R. H. B., DELUCA J., PHILLIPS G., LARocca N. Validity of the Symbol Digit Modalities Test as a cognition performance outcome measure for multiple sclerosis. *Mult Scler.* 2017 Apr; 23(5):721-733.
- BENEDICT R. H. B., FISCHER J. S., ARCHIBALD C. J., ARNETT P. A., BEATTY W. W., BOBHOLZ J., *et al.* Minimal neuropsychological assessment of MS patients: a consensus approach. *Clin Neuropsychol.* Agosto 2002; 16(3):381-97.
- BENITO-LEÓN J., MITCHELL A. J., RIVERA-NAVARRO J., MORALES-GONZÁLEZ J. M. Impaired health-related quality of life predicts progression of disability in multiple sclerosis. *Eur J Neurol.* 2013; 20:79-86.
- BENITO-LEÓN J., MORALES J. M., RIVERA-NAVARRO J. Calidad de vida relacionada con la salud y su relación con el funcionamiento cognitivo y emocional en pacientes con esclerosis múltiple. *Revista Europea de Neurología* 2002; 9(5):497-502.
- BERG D., MAURER M., WARMUTH-METZ M., RIECKMANN P., BECKER G. The correlation between ventricular diameter measured by transcranial sonography and clinical disability and cognitive dysfunction in patients with multiple sclerosis. *Arch Neurol* 2000; 57:1289-92.
- BERGENDAL G¹, FREDRIKSON S., ALMKVIST O. Selective decline in information processing in subgroups of multiple sclerosis: an 8-year longitudinal study. *European Neurology*, 57(4): 193-202.
- BERMEL R. A., BAKSHI R. The measurement and clinical relevance of brain atrophy in multiple sclerosis. *Lancet Neurology* 2006; 5(2):158-70.
- BETHOUX F., BENNETT S. Evaluating walking in patients with multiple sclerosis. which assessment tools are useful in clinical practice? *Int J MS Care.* 2011 Spring; 13(1):4-14.
- BIBERACHER V., BOUCARD C. C., SCHMIDT P., ENGL C., BUCK D., BERTHELE A., HOSHI M. M., ZIMMER C., HEMMER B., MÜHLAU M. Atrophy and structural variability of the upper cervical cord in early multiple sclerosis. *Mult Scler.* 2015; 21(7):875-84.
- BLINKENBERG M., RUNE K., JENSEN C. V., RAVNBORG M., KYLLINGSBAEK S., HOLM S., *et al.*, Cortical cerebral metabolism correlates with MRI lesion load and cognitive dysfunction in MS. *Neurology* 2000; 54:558-64.
- BONAVITA S., GALLO A., SACCO R., CORTE M. D., BISECCO A., DOCIMO R., LAVORGNA L., CORBO D., COSTANZO A. D., TORTORA F., CIRILLO M., ESPOSITO F., TEDESCHI G. Distributed changes in default-mode resting-state connectivity in multiple sclerosis. *Mult Scler.* 2011; 17(4):411-22.
- BONAVITA S., SACCO R., DELLA CORTE M., ESPOSITO S., SPARACO M., D'AMBROSIO A., *et al.*, Computer-aided cognitive rehabilitation improves cognitive performances and induces brain functional connectivity changes in relapsing remitting multiple sclerosis patients: an exploratory study. *J Neurol.* 2015; 262(1):91-100.
- BORCHI M., CAVALLO M., CARLETTO S., OSTACOLI L., ZUFFRANIERI M *et al.* Presence and significant determinants of cognitive impairment in a large simple of patients with multiple sclerosis. *Plos One.* 2013; 8(7):e69820.

- BORGHİ M., CARLETTO S., OSTACOLI L., SCAVELLI F., PIA L., PAGANI M., BERTOLOTTO A., MALUCCHI S., SIGNORI A., CAVALLO M. Decline of Neuropsychological Abilities in a Large Sample of Patients with Multiple Sclerosis: A Two-Year Longitudinal Study. *Front Hum Neurosci.* 2016 Jun 8;10:282. doi: 10.3389/fnhum.2016.00282. eCollection 2016.
- BORINGA J. B., LAZERON R. H. C., REULING J. E. W., ADÈR., H. J., PFENNINGS LEMA., LINDEBOOM J., *et al.*, The Brief Repeatable Battery of normative values allow application in multiple sclerosis clinical practice. *Mult Scler.* 2001; 7:263-8.
- BOSMA L., KRAGT J. J., POLMAN C. H., UITDEHAAG B. M. J. Walking speed, rather than Expanded Disability Status Scale, relates to long-term patient-reported impact in progressive MS. *Mult Scler J.* 2013; 19:326-33.
- BOWLING A. C., IBRAHİM R., STEWART T. M. Alternative medicine and multiple sclerosis: An objective review from an American perspective. *Int J MS Care.* 2000; 2:15-28.
- BRADSHAW, J. Cognition, depression and fatigue in multiple sclerosis. *ACNR 2008 September/October*; 8(4).
- BRANGER P., PARIENTI J. J., SORMANI M. P., DEFER G. The effect of disease-modifying drugs on brain atrophy in relapsing-remitting multiple sclerosis: a meta-analysis. *PLoS One* 2016; 11(3):e0149685.
- BRASS S. D., BENEDICT R. H., WEINSTOCK-GUTTMAN B., MUNSCHAUER F., BAKSHI R. Cognitive impairment is associated with subcortical magnetic resonance imaging grey matter T2 hypointensity in multiple sclerosis. *Mult Scler.* 2006; 12(4): 437-44.
- BRAZIER J. E., HARPER R. USHERWOOD T *et al.* Validating the SF36 health survey questionnaire: new outcome measure for primary care. *Br Med J.* 1992; 305:160-4.
- BRENK A., LAUN K., HAASE C. G. Short-term cognitive training improves mental efficiency and mood in patients with multiple sclerosis. *Eur Neurol.* 2008; 60(6):304-9.
- BRİKEN S., GOLD S. M., PATRA S., VETTORAZZI E., HARBS D., TALLNER A., KETELS G., SCHULZ K. SH., HEESEN C. Effects of exercise on fitness and cognition in progressive MS: a randomized, controlled pilot trial. *Mult Scler.* 2014 Mar; 20(3):382-90.
- BRİSSART H., LEROY M., MORELE E., BAUMANN C., SPITZ E., DEBOUVERIE M. Cognitive rehabilitation in multiple sclerosis. *Neurocase* 2013;19(6): 553-65.
- BROCHET B., DOUSSET V. Pathological correlates of magnetization transfer imaging abnormalities in animal models and humans with multiple sclerosis. *Neurology* 1999; 53(5 suppl 3):S12-17.
- BRODSKY M., *et al.*, The Optic Neuritis Study Group. Multiple sclerosis risk after optic neuritis: final optic neuritis treatment trial follow-up. *Arch Neurol* 2008; 65(6):727-32.
- BROOKS J. B. B., GIRAUD V. O., SALEH Y. J., RODRIGUES S. J., DAIA L. A., FRAGOSO Y. D. Paced auditory serial addition test (PASAT): a very difficult test even for individuals with high intellectual capability. *Arq Neuropsiquiatr.* Junio 2011; 69(3):482-4.
- BRÜCK W., LUCCHINETTI C., LASSMANN H. The pathology of primary progressive multiple sclerosis. *Mult Scler* 2002; 8(2):93-7.
- CABRERA M. E., AGOSTINI M. T., GARCÍA VINIEGRAS., V. (1998). En: Centro de investigaciones Psicológicas y Sociológicas, ed. Calidad de vida y trabajo comunitario integrado. Sistema de indicadores para la medición de la calidad de vida. Reporte de investigación. <http://www.cips.w/index.php/resultados de investigacion del grupo de estudios psicossociales>.

- CADER S., CIFELLI A., ABU-OMAR Y., PALACE J., MATTHEWS P. M. Reduced brain functional reserve and altered functional connectivity in patients with multiple sclerosis. *Brain* 2006; 129(Pt 2):527-37.
- CALABRESE M., PORETTO V., FAVARETTO A., ALESSIO S., BERNARDI V., ROMUALDI C., RINALDI F., PERINI P., GALLO P. Cortical lesion load associates with progression of disability in multiple sclerosis. *Brain* 2012; 135(Pt 10):2952-61.
- CALABRESE M., REYNOLDS R., MAGLIOZZI R., CASTELLARO M., MORRA A., SCALFARI A., FARINA G., ROMUALDI C., GAJOFATTO A., PITTERI M., BENEDETTI M. D., MONACO S. Regional Distribution and Evolution of Gray Matter Damage in Different Populations of Multiple Sclerosis Patients. *PLoS One* 2015; 10(8):e0135428.
- CALABRESE M., RINALDI F., MATTISI I., GROSSI P., FAVARETTO A., ATZORI M., BERNARDI V., BARACHINO L., ROMUALDI C., RINALDI L., PERINI P., GALLO P. Widespread cortical thinning characterizes patients with MS with mild cognitive impairment. *Neurology* 2010; 74:321-28.
- CARRETIÉ-ARANGÜENA L. *Psicofisiología*. Madrid: Ed Pirámide, 2009.
- CARROLL M., GATES R., ROLDAN F. Memory impairment in multiple sclerosis. *Neuropsychologia*. 1984; 22(3):297-302.
- CARTER A. M., DALEY A. J., KESTERTON S. W., WOODROOFE N. M., SAXTON J. M., SHARRACK B. Pragmatic exercise intervention in people with mild to moderate multiple sclerosis: a randomised controlled feasibility study. *Contemp Clin Trials*. 2013 Jul; 35(2):40-7.
- CASANOVA B., MARTÍNEZ-BISBAL M. C., VALERO C., CELDA B., MARTÍ-BONMATÍ L., PASCUAL A., LANDENTE L., CORET F. Evidence of Wallerian degeneration in normal appearing white matter in the early stages of relapsing-remitting multiple sclerosis: a HMRS study. *J Neurol* 2003; 250:22-28.
- CASANOVA-ESTRUCH B., CORET-FERRER F., LANDETE-PASCUAL L. Estudio de diversas escalas de fatiga e impacto en la calidad de vida de los pacientes afectos de esclerosis múltiple. *Rev Neurol* 2000; 30:1235-1241
- CASTRO-BORRERO W., GRAVES D., FROHMAN T. C., FLORES A. B., HARDEMAN P., LOGAN D *et al.*, Current and emerging therapies in multiple sclerosis: a systematic review. *Ther Adv Neurol Disorder*. 2012; 5(4):205-20.
- CAZA S. M., MCKENNA S. P., MCEWEN J., WILLIAMS J., PAPP E. El perfil de salud de Nottingham: Estado de salud subjetivo y consultas médicas. *Soc Sci Med A*. 1981 Mayo; 15 (3 Pt 1):221-9.
- CECCARELLI A., ROCCA M. A., PAGANI E., COLOMBO B., MARTINELLI V., COMI G., FILIPPI M. A voxel-based morphometry study of grey matter loss in MS patients with different clinical phenotypes. *Neuroimage* 2008; 42(1):315-22.
- CELLA D. F., DINNEN K., AMASON B., REDER A., WEBSTER K. A., KARABATROS G. Validation of the functional assessment of multiple sclerosis quality of life instrument. *Neurology* 1996; 47:129-39.
- CERASA A., GIOIA M. C., VALENTINO P., NISTICO R., CHIRIACO C., PIRRITANO D., *et al.* Computer-assisted cognitive rehabilitation of attention deficits for multiplesclerosis: a randomized trial with fMRI correlates. *Neurorehabil Neural Repair* 2013; 27(4):284-95.
- CERCIGNANI M., BOZZALI M., IANNUCCI G., *et al.*, Magnetisation transfer ratio and mean diffusivity of normal-appearing white and gray matter from patients with multiple sclerosis. *J Neurol Neurosurg Psychiatry* 2001; 70: 311-17.

- CEREZO GARCÍA M., MARTÍN PLASENCIA P., ALADRO BENITO Y. Perfil de alteración de las funciones ejecutivas en la esclerosis múltiple. *Acta Neurol Scand.* Mayo 2015; 131(5):313-20. doi: 10.1111 / ane.12345. Epub 2015 8 de febrero.
- CHARIL A., DAGHER A., LERCH J. P., ZIJDENBOS A. P., WORSLEY K. J., EVANS A. C. Focal cortical atrophy in multiple sclerosis: relation to lesion load and disability. *Neuroimage* 2007; 34(2):509-17.
- CHEN J. T., COLLINS D. L., ATKINS H. L., FREEDMAN M. S., ARNOLD DL; Canadian MS/BMT Study Group. Magnetization transfer ratio evolution with demyelination and remyelination in multiple sclerosis lesions. *Ann Neurol.* 2008; 63(2):254-62.
- CHIARAVALLI N., HILLARY F., RICKER J., CHRISTODOULOU C., KALNIN A., LIU W. C., STEFFENER J., DELUCA J. Cerebral activation patterns during working memory performance in multiple sclerosis using fMRI. *J Clin Exp Neuropsychol.* 2005; 27(1):33-54.
- CHIARAVALLI N. D., DE LUCA J., MOORE N. B., RICKER J. H. Treating learning impairments improves memory performance in multiple sclerosis: a randomized clinical trial. *Mult Scler.* 2005;11:58-68.
- CHIARAVALLI N. D., DELUCA J. Cognitive impairment in multiple sclerosis. *Lancet Neurol.* 2008; 7(12):1139-51.
- CHIARAVALLI N. D., DEMAREE H., GAUDINO E., DE LUCA J. Can the repetition effect maximize learning in multiple sclerosis? *Clin Rehabil* 2003; 17:58-68.
- CHIARAVALLI N. D., GENOVA H. M., DE LUCA J. Cognitive rehabilitation in multiple sclerosis: the role of plasticity. *Front Neurol.* 2015 Apr 2; 6:67.
- CHIARAVALLI N. D., WYLIE G., LEAVITT V., DELUCA J. Increased cerebral activation after behavioral treatment for memory deficits in MS. *J Neurol* 2012; 259(7):1337-46.
- CHRISTODOULOU C., KRUPP L. B., LIANG Z., HUANG W., MELVILLE P., ROQUE C., SCHERL W. F., MORGAN T., MACALLISTER W. S., LI L., TUDORICA L. A., LI X., ROCHE P., PEYSTER R. Cognitive performance and MR markers of cerebral injury in cognitively impaired MS patients. *Neurology* 2003; 60(11):1793-8.
- CHWASTIAK L. A., GIBBONS L. E., EHDE DM *et al.*, (2005) Fatigue and psychiatric illness in a large community sample of persons with multiple sclerosis. *Journal of psychosomatic research* 59 (5):291-298
- CIFELLI A., MATTHEWS P. M. Cerebral plasticity in multiple sclerosis: insights from fMRI. *Mult Scler.* 2002; 8(3):193-99.
- COHEN J. A., BARKHOF F., COMI G. *et al.* Oral fingolimod or intramuscular interferon for relapsing multiple sclerosis. *N Engl J Med* 2010; 362(5):402-415.
- COHEN M., BROCHET B., CLAVELOU P., LE PAGE E., VERMERSCH P., TOURBAH A., MOREAU T., JOLY H., SAKAROVITCH C., LEBRUN C. Cognition and quality of life in clinically isolated syndrome patients starting a disease modifying therapy in the QUALICIS study may not predict treatment response at one year. *J Neurol Sci.* 2017 Nov 15; 382:73-78.
- COLORADO RVA., SHUKLA K., ZHOU Y., WOLINSKY J. SS., NARAYANA P. A. Multi-task functional MRI in multiple sclerosis patients without clinical disability. *Neuroimage* 2012; 59(1):573-81.
- COMI G., FILIPI M., MARTINELLI V., CAMPI A., RODGHER M., ALBERTONI M., *et al.* Brain MRI correlates of cognitive impairment in primary and secondary progressive multiple sclerosis. *J Neurol Sci.* 1995; 132:222-7.

- COMI G., FILIPPI M., ROVARIS M., LEOCANI L., MEDAGLINI S., LOCATELLI T. Clinical, neurophysiological, and magnetic resonance imaging correlations in multiple sclerosis. *J Neurol Neurosurg Psychiatr.* 1998; 4(Suppl.):s21-5.
- COMI G., ROVARIS M., FALAUTANO M., SANTUCCIO G., MARTINELLI V., ROCCA M. A., *et al.* A multiparametric MRI study of frontal lobe dementia in multiple sclerosis. *J Neurol Sci.* 1999; 171:135-44.
- COMPSTON A., COLES A. Multiple sclerosis. *Lancet* 2008 Oct 25; 372(9648):1502-17. doi: 10.1016/S0140-6736(08)61620-7.
- COO H., HOPMAN WM *et al.* The Paced Auditory Serial Addition Test: to what extent is it performed as instructed, and is it associated with disease course? *Mult Scler.* 2005 Feb; 11(1):85-9.
- CORES E., VANOTTI S., MOYANO P. Resolution strategies of the PASAT in Multiple Sclerosis patients and viability of a short version of the test. *Rev Chil Neuropsicol.* 2011; 6(2):81-85.
- COSTA S. L., GENOVA H. M., DELUCA J., CHIARAVALLOTI N. D. Information processing speed in multiple sclerosis: Past, present, and future. *Multiple Sclerosis* 2017; 23(6):772-789.
- COVEY T. J., SHUCARD J. L., SHUCARD D. W. Event-related brain potential indices of cognitive function and brain resource reallocation during working memory in patients with Multiple Sclerosis. *Clin Neurophysiol.* 2017; 128(4):604-21.
- CRUCE R., VOSOUGHI R., FREEDMAN M. S. Cognitive impact of anticholinergic medication in MS: Adding insult to injury? *Mult Scler Relat Disord.* 2012 Oct; 1(4):156-61.
- CRUZ A., BELENGUER A., MARTÍNEZ B., FITTIPALDI M., FROM C. Cambios estructurales y funcionales del hipocampo en pacientes con esclerosis múltiple y su relación con procesos de memoria. *Rev Neurol.* 2016; 62(1):6-12.
- CRUZ-GOMEZ A. J., VENTURA-CAMPOS N., BELENGUER A., AVILA C., FORN C. The link between resting-state functional connectivity and cognition in MS patients. *Mult Scler.* 2014; 20(3):338-48.
- CUTTER G. R., BAIER M. L., RUDICK R. A., COOKFAIR D. L., FISCHER J. S., PETKAU J., *et al.* Development of a multiple sclerosis functional composite as a clinical trial outcome measure. *Brain J Neurol.* Mayo 1999; 122(Pt5):871-82.
- D'ALISA S., MISCIO G., BAUDO S., SIMONE A., TESIO L., MAURO A. Depression is the main determinant of quality of life in multiple sclerosis: a classification-regression (CART) study. *Disabil Rehabil.* 2006; 28:307-14.
- DACKOVIC J., PEKMEZOVIC T., MESAROS S., *et al.*, The Rao's Brief Repeatable Battery in the study of cognition in different multiple sclerosis phenotypes: application of normative data in a Serbian population. *Neurol Sci.* 2016 Sep; 37(9):1475-81. doi: 10.1007/s10072-016-2610-1. Epub 2016 May 20.
- DAGENAIS E., ROULEAU I., TREMBLAY A., DEMERS M., ROGER É., JOBIN C., DUQUETTE P. Memoria prospectiva en la esclerosis múltiple : el impacto de la característica distintiva y el funcionamiento ejecutivo. *Brain Cogn.* 2016 Nov; 109:66-74. doi: 10.1016 / j.bandc.2016.07.011. Epub 2016 Sep 17.
- DAHNIKE R., YOTTER R. A., GASER C. Cortical thickness and central surface estimation. *Neuroimage* 2013; 65:336-48.

- DALTON C. M., CHARD D. T., DAVIES G. R., *et al.*, Early development of multiple sclerosis is associated with progressive grey matter atrophy in patients presenting with clinically isolated syndromes. *Brain* 2004; 127:1101-07.
- DASGUPTA R., FOWLER C. J., BLADDER., bowel and sexual dysfunction in multiple sclerosis: management strategies. *Drugs* 2003; 63:153-66.
- DAVIE C. A., HAWKINS C. P., BARKER G. SJ., *et al.*, Serial proton magnetic resonance spectroscopy in acute multiple sclerosis lesions. *Brain* 1994; 117:49-58.
- DAVIES M. B., WILLIAMS R., HAQ N., PELOSI L., HAWKINS C. P. MRI of optic nerve and postchiasmal visual pathways and visual evoked potentials in secondary progressive multiple sclerosis. *Neuroradiology* 1998; 40:765-70.
- DE GIGLIO L., DE LUCA F., PROSPERINI L., BORRIELLO G., BIANCHI V., PANTANO P., POZZILLI C. A low-cost cognitive rehabilitation with a commercial video game improves sustained attention and executive functions in multiple sclerosis: a pilot study. *Neurorehabil Neural Repair*. 2015 Jun; 29(5):453-61.
- DE GIGLIO L., MARINELLI F., BARLETTA V. T., PAGANO V. A., DE ANGELIS F., FANELLI F., PETSAS N., PANTANO P., TOMASSINI V., POZZILLI C. Effect on Cognition of Estroprogestins Combined with Interferon Beta in Multiple Sclerosis: Analysis of Secondary Outcomes from a Randomised Controlled Trial. *CNS Drugs* 2017 Feb; 31(2):161-168.
- DE GIGLIO L., TONA F., DE LUCA F., PETSAS N., PROSPERINI L., BIANCHI V., POZZILLI C., PANTANO P. Multiple sclerosis: Changes in thalamic resting-state functional connectivity induced by a home-based cognitive rehabilitation program. *Radiology*. 2016 Mar; 8:150710.
- DE SONNEVILLE L. M. J., BORINGA J. B., REULING I. E. W., LAZARON R. H. C., ADER H. J., POLMAN C. H. Information processing characteristics in subtypes of multiple sclerosis. *Neuropsychologia* 2002; 40:1751-65.
- DE STEFANO N., BARTOLOZZI M. L., GUIDI L., STROMILLO M. SL., FEDERICO A. Magnetic resonance spectroscopy as a measure of brain damage in multiple sclerosis. *J Neurol Sci*. 2005; 233(1-2):203-8.
- DE STEFANO N., FILIPPI M. MR spectroscopy in multiple sclerosis. *J Neuroimaging*. 2007; 17 (Suppl 1):S31-S35.
- DE STEFANO N., MATTHEWS P. M., FILIPPI M., AGOSTA F., DE LUCA M., BARTOLOZZI ML *et al.* Evidence of early cortical atrophy in MS: relevance to white matter changes and disability. *Neurology* 2003; 60:1157-62.
- DEFER G., DANIEL F. MARIÉ R. Study of episodic memory in multiple sclerosis using the California Verbal Learning Test: the data favour altered encoding. *Rev Neurol*. 2006; 162:852-7.
- DEHMESKI J., SILVER N. SC., LEARY S. M., *et al.*, Magnetisation transfer ratio histogram analysis of primary progressive and other multiple sclerosis subgroups. *J Neurol Sci*. 2001; 185:11-17.
- DELGADO-MENDILÍVAR J. M., CADENAS-DÍAZ J. C., FERNÁNDEZ-TORRICO J. M., NAVARRO-MASCARELL G., IZQUIERDO G. Estudio de la calidad de vida en la esclerosis múltiple. *Rev Neurol*. 2005; 41:257-62.
- DELOIRE M., RUET A., HAMEL D., BONNET M., BROCHET B. Early cognitive impairment in multiple sclerosis predicts disability outcome several years later. *Mult Scler*. 2010;16(5):581-7.
- DELOIRE M. S., SALORT E., BONNET M., ARIMONE Y., BOUDINEAU M., AMIEVA H., BARROSO B., OUALLET J. C., PACHAI C., GALLIAUD E., PETRY K. G., DOUSSET V., FABRIGOULE C., BROCHET

- B. Cognitive impairment as marker of diffuse brain abnormalities in early relapsing remitting multiple sclerosis. *J Neurol Neurosurg Psychiatry* 2005; 76(4):519-26.
- DELUCA J., GAUDINO E. A., DIAMOND B. J., *et al.* Acquisition and storage deficits in multiple sclerosis. *J Clin Exp Neuropsychol* 1998; 20:376-390.
- DEMAREE H. A., GAUDINO E., DELUCA J. The relationship between depressive symptoms and cognitive dysfunction in multiple sclerosis. *Cogn Neuropsychiatry* 2003; 8:161- 171.
- D'ESPOSITO M. Functional neuroimaging of cognition. *Semin Neurol.* 2000; 20(4):487-98.
- DOBRYAKOVA E., WYLIE G. R., DE LUCA J., CHIARAVALLOTI N. D. A pilot study examining functional brain activity 6 months after memory retraining in MS: the MEMREHAB trial. *Brain Imaging Behav.* 2014 Sep; 8(3):403-6.
- DOWARD L. C., MCKENNA S. P., MEADS D. M., TWISS J., ECKERT B. J. The development of patient-reported outcome indices for multiple sclerosis (PRIMUS). *Mult Scler.* 2009; 15:1092-102.
- DRAKE A. S., WEINSTOCK-GUTTMAN B., MORROW S. A., *et al.* Psychometrics and normative data for the Multiple Sclerosis Functional Composite: Replacing the PASAT with the Symbol Digit Modalities Test. *Mult Scler.* 2010; 16: 228-237.
- DUNCAN-JOHNSON C. C., DONCHIN E. The P300 component of the event-related brain potential as an index of information processing. *Biol Psychol.* 1982; 14: 1-52.
- DUQUE B., SEPULCRE J., BEJARANO B., SAMARANCH L., PASTOR P., VILLOSLADA P. Memory decline evolves independently of disease activity in MS. *Mult Scler Houndmills Basingstoke Engl.* 2008 Agosto; 14(7):947-53.
- DUQUE P., OLTRA-CUCARELLA O., FERNÁNDEZ O., SEPULCRE J. Bateria neuropsicológica breve en esclerosis múltiple. Baremación normativa estratificada por edad y nivel educativo. *Revista Neuropsicológica* 2017; 64:97-104.
- ELLGER T., BETHKE F., FRESE A., LUETTMANN R. J., BUCHHEISTER A., RINGELSTEIN E. B., EVERS S. Event-related potentials in different subtypes of multiple sclerosis: a cross-sectional study. *J Neurol Sci.* 2002; 205(1): 35-40.
- ENGLUND A., FREEMAN T. P., MURRAY R. M., MCGUIRE P. Can we make cannabis safer? *Lancet Psychiatry.* 2017 Mar 1. pii: S2215-0366(17)30075-5.
- ENZINGER C., BARKHOF F., CICCARELLI O., FILIPPI M., KAPPOS L., ROCCA M. A., ROPELE S., ROVIRA À., SCHNEIDER T., DE STEFANO N., VRENKEN H., WHEELER-KINGSHOTT C., WUERFEL J., FAZEKAS F; MAGNIMS study group. Non-conventionalMRI and microstructuralcerebralchanges in multiplesclerosis. *Nat Rev Neurol.* 2015; 11(12):676-86.
- ENZINGER C., FAZEKAS F. Measuring gray matter and white matter damage in MS: Why this is not enough. *Front Neurol.* 2015; 17:6:56.
- ENZINGER C., PINTER D., ROCCA M. A., DE LUCA J., SASTRE-GARRIGA J., AUDOIN B., FILIPPI M. Longitudinal fMRI studies: Exploring brain plasticity and repair in MS. *Mult Scler.* 2016; 22(3):269-78.
- ERIKSSON M., ANDERSEN O., RUNMARKER B. Long-term follow up of patients with clinically isolated syndromes, relapsing-remitting and secondary progressive multiple sclerosis. *Mult Scler.* 2003; 9(3):260-74.
- ERNST A., BOTZUNG A., GOUNOT D., SELLAL F., BLANC F., DE SEZE J., *et al.*, Induced brain plasticity after a facilitation programme for autobiographical memory in multiple sclerosis: a preliminary study. *Mult Scler Int* 2012:820240.

- FAIVRE A., RICO A., ZAARAOUI W., CRESPIY L., REUTER F., WYBRECHT D., *et al.*, Assessing brain connectivity at rest is clinically relevant in early multiple sclerosis. *Mult Scler.* 2012; 18(9):1251-8.
- FAN J., McCANDLISS B. D., SOMMER T., RAZ A., POSNER M. I. Testing the efficiency and independence of attentional networks. *J Cogn Neurosci.* 2002; 14(3):340-347.
- FEINSTEIN A., LAPSHIN H., O'CONNOR P., LANCTÔT K. L. Sub-threshold cognitive impairment in multiple sclerosis: the association with cognitive reserve. *J Neurol.* 2013 Sep;260(9):2256-61.
- FERNÁNDEZ O., FERNÁNDEZ V., BAUMSTARCK-BARRAU K., MUÑOZ L., GONZÁLEZ ALVAREZ MDEL M., ARRABAL J. C., LEÓN A., ALONSO A., LÓPEZ-MADRONA J. C., BUSTAMANTE R., LUQUE G., GUERRERO M., DI CANTOGNO E. V., AUQUIER P; MusiQoL grupo de estudio de España. Validación de la versión en español del cuestionario de Calidad de Vida Internacional de esclerosis múltiple (MusiQoL). *BMC Neurol.* 2011 18 Octubre; 11:127. doi: 10.1186 / 1471-2377-11-127.
- FERRUCCI R., VERGARI M., COGIAMANIAN F., BOCCI T., CIOCCA M., TOMASINI E., DE RIZ M., SCARPINI E., PRIORI A. Transcranial direct current stimulation (tDCS) for fatigue in multiple sclerosis. *NeuroRehabilitation.* 2014; 34(1):121-7.
- FEUILLET L., REUTER F., AUDOIN B., MALIKOVA I., BARRAU K., CHERIF A. A., *et al.*, Early cognitive impairment in patients with clinically isolated syndrome suggestive of multiple sclerosis. *Mult Scler.* 2007; 13(1):124-7.
- FILIPPI M., ABSINTA M., ROCCA M. A. Future MRI tools in multiple sclerosis. *J Neurol Sci.* 2013a; 331(1-2):14-8.
- FILIPPI M., AGOSTA F. Imaging biomarkers in multiple sclerosis. *J Magn Reson Imaging* 2010; 31(4):770-88.
- FILIPPI M., AGOSTA F. Magnetic resonance techniques to quantify tissue damage, tissue repair, and functional cortical reorganization in multiple sclerosis. *Prog Brain Res.* 2009; 175:465-82.
- FILIPPI M., HORSFIELD M. A., ADÈR H. J., BARKHOF F., BRUZZI P., EVANS A., FRANK J. A., GROSSMAN R. I., MCFARLAND H. F., MOLYNEUX P., PATY D. W., SIMON J., TOFTS P. S., WOLINSKY J. S., MILLER D. H. Guidelines for using quantitative measures of brain magnetic resonance imaging abnormalities in monitoring the treatment of multiple sclerosis. *Ann Neurol.* 1998; 43(4):499-506.
- FILIPPI M., PREZIOSA P., COPETTI M., RICCITELLI G., HORSFIELD M. A., MARTINELLI V., COMI G., ROCCA M. A. Gray matter damage predicts the accumulation of disability 13 years later in MS. *Neurology* 2013b; 81(20):1759-67.
- FILIPPI M., RICCITELLI G., MATTIOLI F., CAPRA R., STAMPATORI C., PAGANI E., *et al.*, Multiple sclerosis: effects of cognitive rehabilitation on structural and functional MR imaging measures-an explorative study. *Radiology* 2012; 262(3):932-40.
- FILIPPI M., ROCCA M. A., CICCARELLI O., DE STEFANO N., EVANGELOU N., KAPPOS L., ROVIRA A., SASTRE-GARRIGA J., TINTORÈ M., FREDERIKSEN J. L., GASPERINI C., PALACE J., REICH D. S., BANWELL B., MONTALBAN X., BARKHOF F; MAGNIMS Study Group. MRI criteria for the diagnosis of multiple sclerosis: MAGNIMS consensus guidelines. *Lancet Neurol.* 2016; 15(3):292-303.
- FILIPPI M., ROCCA M. A., DE STEFANO N., ENZINGER C., FISHER E., HORSFIELD M. A., INGLESE M., PELLETIER D., COMI G. Magnetic resonance techniques in multiple sclerosis: the present and the future. *Ann Neurol.* 2011; 68(12):1514-2011.

- FILIPPI M., ROCCA M. A. Conventional MRI in multiple sclerosis. *J Neuroimaging* 2007; 17(Suppl 1):3S-9S.
- FILIPPI M., ROCCA M. A. Functional MR imaging in multiple sclerosis. *Neuroimaging Clin N Am.* 2009; 19(1):59-70.
- FILIPPI M., ROCCA M. A. Magnetization transfer magnetic resonance imaging of the brain, spinal cord, and optic nerve. *Neurotherapeutics* 2007b; 4(3):401-13.
- FILIPPI M., ROCCA M. A. Present and future of fMRI in multiple sclerosis. *Expert Rev Neurother.* 2013; 13(12 Suppl):27-31.
- FINK F., RISCHKAU E., BUTT M., KLEIN J., ELING P., HILDEBRANDT H. Efficacy of an executive function intervention programme in MS: A placebo-controlled and pseudorandomized trial. *Mult Scler.* 2010; 16(9):1148-51.
- FISCHER J. S., FOLEY F. W., AIKENS J. E., ERICSON G. D., RAO S. M., SHINDELL S. J. What do we really know about cognitive dysfunction, affective disorders, and stress in multiple sclerosis? A practitioner's guide. *Neurol Rehabil.* 1994; 8:151-64. b;15(2):244-50. doi:10.1177/1352458508097924. Epub 2008
- FISCHER J. S., LARocca N. G., MILLER D. M., RITVO P. G., ANDREWS H., PATY D. Recent developments in the assessment of quality of life in multiple sclerosis. *Mult Scler.* 1999; 5:251-9.
- FISCHER J. S., PRIORE R. L., JACOBS L. D *et al.* Neuropsychological effects of interferon beta-1a in relapsing multiple sclerosis. Multiple Sclerosis Collaborative Research Group. *Ann Neurol* 2000; 48:885-892.
- FISHER E., LEE J. C., NAKAMURA K., RUDICK R. A. Gray matter atrophy in multiple sclerosis: A longitudinal study. *Ann Neurol.* 2008; 64(3): 255-65.
- FISNIKU L. K., BREX P. A., ALTMANN D. R., MISZKIEL K. A., BENTON C. E., LANYON R., THOMPSON A. J., MILLER D. H. Disability and T2 MRI lesions: a 20-year follow-up of patients with relapse onset of multiple sclerosis. *Brain* 2008; 131(Pt 3):808-17.
- FITTIPALDI-MÁRQUEZ M. S., CRUZ-GÓMEZ Á. J., SANCHIS-SEGURA C., BELENGUER A., ÁVILA C., FORN C. Exploring neural efficiency in multiple sclerosis patients during the symbol digit modalities test: a functional magnetic resonance imaging study. *Neurodegener Dis.* 2017; 17(4-5):199-207.
- FOLEY F. W., LARocca N. G., SANDERS A. Z., ZEMON V. Rehabilitation of intimacy and sexual dysfunction in couples with multiple sclerosis. *Mult Scler.* 2001; 7:417-21.
- FOONG J., ROZEWICZ L., CHONG W. K., THOMPSON A. J., MILLER D. H., RON M. A. A comparison of neuropsychological deficits in primary and secondary progressive multiple sclerosis. *J Neurol* 2000; 247(2):97-101.
- FOONG J., ROZEWICZ L., QUAGHEBEUR G., DAVIE C. A., KARTSOUNIS L. D., THOMPSON A. J., *et al.* Executive functions in multiple sclerosis. The role of frontal lobe pathology. *Brain* 1997; 120:15-26.
- FORN C., BARROS-LOSCERTALES A., ESCUDERO J., BELLOCH V., CAMPOS S., PARCET M. A., AVILA C. Cortical reorganization during PASAT task in MS patients with preserved working memory functions. *Neuroimage* 2006; 31(2):686-91.

- FORN C., BARROS-LOSCERTALES A., ESCUDERO J., BENLLOCH V., CAMPOS S., ANTÒNIA PARCET M., AVILA C. Compensatory activations in patients with multiple sclerosis during preserved performance on the auditory N-back task. *Hum Brain Mapp* 2007; 28(5):424-30.
- FORN C., ROCCA M. A., VALSASINA P., BOSCA I., CASANOVA B., SANJUAN A., ÀVILA C., FILIPPI M. Functional magnetic resonance imaging correlates of cognitive performance in patients with a clinically isolated syndrome suggestive of multiple sclerosis at presentation: an activation and connectivity study. *Mult Scler*. 2012; 18(2):153-63.
- FRANKLIN G. M., NELSON L. M., FILLEY C. M., HEATON R. K. Cognitive loss in multiple sclerosis. Case reports and review of the literature. *Arch Neurol*. 1989; 46:162-167.
- FREITAS S., CALLEGARO D., POMPÉIA S., ORLANDO F. Working memory impairment in multiple sclerosis relapsing-remitting patients with episodic memory deficits. *Arquivos de Neuropsiquiatria* 2010; 68:205-11.
- GADEA M., MARTÍNEZ-BISBAL M. C., MARTI-BONMATÍ L., ESPERT R., CASANOVA B., CORET F., CELDA B. Spectroscopic axonal damage of the right locus coeruleus relates to selective attention impairment in early stage relapsing-remitting multiple sclerosis. *Brain* 2004; 127(Pt 1):89-98.
- GALVAO-CARMONA A., GONZÁLEZ-ROSA J. J., HIDALGO-MUÑOZ A. R., PÁRAMO D., BENÍTEZ M. L., IZQUIERDO G., VÁZQUEZ-MARRUFO M. Disentangling the attention network test: behavioral, event related potentials, and neural source analyses. *Front Hum Neurosci*. 2014; 13; 8: 813.
- GARCÍA-LORENZO D., FRANCIS S., NARAYANAN S., ARNOLD D. L., COLLINS D. L. Review of automatic segmentation methods of multiple sclerosis white matter lesions on conventional magnetic resonance imaging. *Med Image Anal*. 2013; 17(1):1-18.
- GARRETT, M., HOGAN, N., LARKIN, A., SAUNDERS, J., JAKEMAN, P., COOTE, S. Exercise in the community for people with minimal gait impairment due to MS: An assessor-blind randomized controlled trial. *Multiple Sclerosis* 2013; 19:782-789.
- GARRIDO M. I., KILNER J. M., STEPHAN K. E., FRISTON K. J. The mismatch negativity: A review of underlying mechanisms. *Clin Neurophysiol*. 2009; 120:453-63.
- GASS A., ROCCA M. A., AGOSTA F., CICCARELLI O., CHARD D., VALSASINA P., BROOKS J. C., BISCHOF A., EISELE P., KAPPOS L., BARKHOF F., FILIPPI M; MAGNIMS Study Group. MRI monitoring of pathological changes in the spinal cord in patients with multiple sclerosis. *Lancet Neurol*. 2015; 14(4):443-54.
- GAUDINO E. A., CHIARAVALLOTI N. P., DELUCA J., DIAMOND B. J. A comparison of memory performance in relapsing-remitting, primary progressive and secondary progressive, multiple sclerosis. *Neuropsychiatry Neuro- psychol Behav Neurol*. 2001; 14:32-44.
- GE Y., GROSSMAN R. I., UDUPA J. K., WEI L., MANNON L. J., POLANSKY M., KOLSON D. L. Brain atrophy in relapsing-remitting multiple sclerosis and secondary progressive multiple sclerosis: longitudinal quantitative analysis. *Radiology* 2000; 214(3):665-70.
- GEISLER M. W., SLIWINSKI M., COYLE P. K., MASUR D. M., DOSCHER C., KRUPP L. B. The effects of amantadine and pemoline on cognitive functioning in multiple sclerosis. *Arch Neurol*. 1996 Feb; 53(2):185-8.
- GEISSELER O., PFLUGSHAUPT T., BEZZOLA L., REUTER K., WELLER D., SCHUKNECHT B., BRUGGER P., LINNEBANK M. The relevance of cortical lesions in patients with multiple sclerosis. *BMC Neurol*. 2016; 16(1):204.

- GENOVA H. M., HILLARY F. G., WYLIE G., RYPMA B., DELUCA J. Examination of processing speed deficits in multiple sclerosis using functional magnetic resonance imaging. *J Int Neuropsychol Soc.* 2009; 15(3):383-93.
- GENOVA., H. M., LENGENFELDER J., CHIARAVALLOTI N. D., MOORE N. B., DELUCA J. Processing Speed Versus Working Memory: Contributions to an Information-Processing Task in Multiple Sclerosis. *Applied Neuropsychology: Adult* 2012; 19(2):132-140. doi:10.1080/09084282.2011.643951
- GICH J., FREIXANET J., GARCÍA R., VILANOVA J. C., GENÍS D., SILVA Y., MONTALBAN X., RAMIÓ-TORRENTÀ L. A randomized, controlled, single-blind, 6-month pilot study to evaluate the efficacy of MS-Line!: a cognitive rehabilitation programme for patients with multiple sclerosis. *Mult Scler.* 2015b Sep; 21(10):1332-43.
- GICH J., FREIXANET J., GARCIA R., VILANOVA J. C., GENÍS D., SILVA Y., MONTALBAN X., RAMIÓ-TORRENTÀ L. A new cognitive rehabilitation programme for patients with multiple sclerosis: the 'MS-line! Project'. *Mult Scler.* 2015a Sep; 21(10):1344-8.
- GIESSER B. S., SCHROEDER M. M., LARocca N. G., KURTZBERG D., RITTER W., VAUGHAN H. G., SCHEINBERG L. C. Endogenous event-related potentials as indices of dementia in multiple sclerosis patients. *Electroencephalogr Clin Neurophysiol.* 1992; 82(5):320-9.
- GIL R., ZAI L., NEAU J. P., JONVEAUX T., AGBO C., ROSOLACCI T., BURBAUD P., INGRAND P. Event-related auditory evoked potentials and multiple sclerosis. *Electroencephalogr Clin Neurophysiol.* 1993; 88(3):182-7.
- GILSON B. S., GILSON J. S., BERGNER M., BOBBIT R. A., KRESSEL S., POLLARD W. E., VESSELAGO M. The sickness impact profile. Development of an outcome measure of health care. *Am J Public Health.* 1975; 65:1304-10.
- GIORDANO A., PUCCI E., NALDI P., *et al.* Responsiveness of patient reported outcome measures in multiple sclerosis relapses: the REMS study. *Journal of Neurology, Neurosurgery and Psychiatry* 2009; 80(9):1023-1028.
- GIORGIO A., DE STEFANO N. Cognition in multiple sclerosis: relevance of lesions., brain atrophy and proton MR spectroscopy. *Neurol Sci.* 2010; 31(Suppl 2):S245-48.
- GLADE M. J. Vitamin D: Health panacea or false prophet? *Nutrition.* 2013 Jan; 29(1):37-41.
- GLANZ B. I., HEALY B. C., RINTELL D. J., JAFFIN S. K., BAKSHI R., WEINER H. L. The association between cognitive impairment and quality of life in patients with early multiple sclerosis. *Journal of the Neurological Sciences* 2010; 290(1-2):75-79.
- GLANZ B. I., HOLLAND C. M., GAUTHIER S. A., AMUNWA E. L., LIPTAK Z., HOUTCHENS M. K., *et al.* Cognitive dysfunction in patients with clinically isolated syndromes or newly diagnosed multiple sclerosis. *Mult Scler.* 2007; 13(8):1004-10.
- GOLD S. M., HEESEN C., SCHULZ H., *et al.* Disease specific quality of life instruments in multiple sclerosis: validation of the Hamburg Quality of Life Questionnaire in Multiple Sclerosis (HAQUAMS). *Mult Scler.* 2001 7:119-130.
- GOLD S. M., SCHULZ H., MÖNCH A., SCHULZ K. H., HEESEN C. Cognitive impairment in multiple sclerosis does not affect reliability and validity of self-report health measures. *Mult Scler.* 2003; 9(4):404-410.

- GONZÁLEZ TORRE J. A., CRUZ-GÓMEZ A. J., BELENGUER A., SANCHIS-SEGURA C., ÁVILA C., FORN C. Hippocampal dysfunction is associated with memory impairment in multiple sclerosis: A volumetric and functional connectivity study. *Mult Scler.* 2017; 23(14):1854-63.
- GONZÁLEZ-GARCÍA C., TUDELA P., RUZ M. Resonancia magnética funcional: análisis crítico de sus implicaciones técnicas, estadísticas y teóricas en neurociencia humana. *Rev Neurol.* 2014; 58(7):318-25.
- GONZALEZ-ROSA J. J., VAZQUEZ-MARRUFO M., VAQUERO E., DUQUE P., BORGES M., GAMERO M. A., GOMEZ C. M., IZQUIERDO G. Differential cognitive impairment for diverse forms of multiple sclerosis. *BMC Neurosci.* 2006; 7:39.
- GONZALEZ-ROSA J. J., VAZQUEZ-MARRUFO M., VAQUERO E., DUQUE P., BORGES M., GOMEZ-GONZALEZ C. M., IZQUIERDO G. Cluster analysis of behavioural and event-related potentials during a contingent negative variation paradigm in remitting-relapsing and benign forms of multiple sclerosis. *BMC Neurol.* 2011;11:64.
- GOVEROVER Y., CHIARAVALLOTI N., DELUCA J. The relationship between self-awareness of neuro-behavioral symptoms, cognitive functioning, and emotional symptoms in multiple sclerosis. *Multiple Sclerosis* 2005; 11(2):203-212.
- GOVEROVER Y., CHIARAVALLOTI N., GENOVA H., DELUCA J. A randomized controlled trial to treat impaired learning and memory in multiple sclerosis: The self-GEN trial. *Mult Scler.* 2017 May 1:1352458517709955. doi: 10.1177/1352458517709955.
- GROMISCH E. S., ZEMN V., HOLTZER R., CHIARAVALLOTI N. D., DELUCA J *et al.*, Assessing the criterion validity of for highly abbreviated measures from the Minimal Assessment of Cognitive Function in Multiple Sclerosis (MACFIMS). *Clin Neuropsychol.* 2016 Oct; 30(7):1032-49.
- HABEK M., HOJSAK I., BRINAR V. V. Nutrition in multiple sclerosis. *Clin Neurol Neurosurg.* 2010 Sep; 112(7):616-20
- HALPER J., PERRIN ROSS A. Challenges in the Treatment of Mobility Loss and Walking Impairment in Multiple Sclerosis. *Int J MS Care* 2010; 12:13-16.
- HANKOMÄKI E., MULTANEN J., KINNUNEN E., HÄMÄLÄINEN P. The progress of cognitive decline in newly diagnosed MS patients. *Acta Neurol Scand.* 2014 Marzo; 129(3):184-91.
- HANSEN S., MUENSSINGER J., KRONHOFMANN S., LAUTENBACHER S., OSCHMANN P., KEUNE P. M. Cognitive screening in Multiple Sclerosis: the Five-Point Test as a substitute for the PASAT in measuring executive function. *Clin Neuropsychol.* 2017 Jan; 31(1):179-192.
- HAO J. J., CHENG W., LIU M., LI H., LÜ X., SUN Z. Treatment of Multiple sclerosis With Chinese scalp Acupuncture. *Glob Adv Health Med.* 2013; 2:8-13.
- HARDMEIER M., SCHOONHEIM M. M., GEURTS J. J., HILLEBRAND A., POLMAN C. H., BARKHOF F., STAM C. J. Cognitive dysfunction in early multiple sclerosis: altered centrality derived from resting-state functional connectivity using magneto-encephalography. *PLoS One* 2012; 7(7):e42087.
- HAREL Y., APPLEBOIM N., LAVIE M., ACHIRON A. Single dose of methylphenidate improves cognitive performance in multiple sclerosis patients with impaired attention process. *J Neurol Sci.* 2009 Jan 15; 276(1-2):38-40.
- HARRIS INTERACTIVE. Experiences with MS: Perspectives of people with MS and MS care partners. 2008 March 25.

- HE D., ZHANG Y., DONG S., WANG D., GAO X., ZHOU H. Pharmacological treatment for memory disorder in multiple sclerosis. *Cochrane Database Syst Rev.* 2013 Dec 17; (12):CD008876.
- HEEGER D. J., RESS D. What does fMRI tell us about neuronal activity? *Nat Rev Neurosci.* 2002; 3(2): 142-51.
- HEESEN C., BOHM J., REICH C., KASPER J., GOEBEL M., GOLD S. M. Patient perception of bodily functions in multiple sclerosis: gait and visual function are the most valuable. *Mult Scler.* 2008; 14(7):988-991.
- HERNÁNDEZ M. A., MORA S. Evaluación de la calidad de vida mediante cuestionario PRIMUS en la población española de pacientes con esclerosis múltiple. *Neurología* 2013; 28(6):340-347.
- HICKMAN S. J. Optic Nerve Imaging in Multiple Sclerosis. *J Neuroimaging* 2007; 17:S42-45.
- HILLARY F. G., CHIARAVALLOTI N. D., RICKER J. H., STEFFENER J., BLY B. M., LANGE G., LIU W. C., KALNIN A. J., DELUCA J. An investigation of working memory rehearsal in multiple sclerosis using fMRI. *J Clin Experimental Neuropsychol.* 2003; 25(7):965-78.
- HOBART J., LAMPING D., FITZPATRICK R., RIAZI, THOMPSON A. The Multiple Sclerosis Impact Scale (MSIS-29): A new patient-based outcome measure. *Brain* 2001 May; 124(5):962-973. <https://doi.org/10.1093/brain/124.5.962>.
- HOBART J. C., RIAZI A., LAMPING D. L., *et al.*, How responsive is the Multiple Sclerosis Impact Scale (MSIS-29)? A comparison with some other self report scales. *J Neurol Neurosurg Psychiatry.* 2005 Nov; 76(11):1539-1543.
- HOLLINGER K. R., ALT J., RIEHM A. M., SLUSHER B. S., KAPLIN A. I. Dose-dependent inhibition of GCPII to prevent and treat cognitive impairment in the EAE model of multiple sclerosis. *Brain Res.* 2016 Mar 15; 1635:105-12.
- HONARMAND K., AKBAR N., KOU N., FEINSTEIN A. Predicting employment status in multiple sclerosis patients: the utility of the MS functional composite. *J Neurol.* 2011 Feb; 258(2):244-9. doi: 10.1007/s00415-010-5736-8. Epub 2010 Sep 12.
- HONIG L. S., RAMSAY R. E., SHEREMATA W. A. Event-related potential P300 in multiple sclerosis. Relation to magnetic resonance imaging and cognitive impairment. *Arch Neurol.* 1992; 49(1):44-50.
- HORAKOVA D., DWYER M. G., HAVRDOVA E., COX J. L., DOLEZAL O., BERGLAND N., RIMES B., SEIDL Z., VANECKOVA M., ZIVADINOV R. Gray matter atrophy and disability progression in patients with early relapsing-remitting multiple sclerosis: a 5-year longitudinal study. *J Neurol Sci.* 2009; 282(1-2):112-9.
- HORSFIELD M. A. Magnetization transfer imaging in multiple sclerosis. *J Neuroimaging.* 2005; 15(4 Suppl):S58-67.
- HOUTCHENS M. K., BENEDICT R. H., KILLIANY R., SHARMA J., JAISANI Z., SINGH B., WEINSTOCK-GUTTMAN B., GUTTMANN C. R., BAKSHI R. Thalamic atrophy and cognition in multiple sclerosis. *Neurology* 2007; 69(12):1213-23.
- HOWARTH A. L. Will aromatherapy be a useful treatment strategy for people with multiple sclerosis who experience pain? *Complement Ther Nurs Midwifery.* 2002; 8:138-41.
- HUIJBREGTS, S. C. J., KALKERS, N. F., DE SONNEVILLE, L. M. J., DE GROOT, V. Differences in cognitive impairment of relapsing remitting, secondary, and primary progressive MS. *Neurology* 2004 Jul 27; 63(2):335-9.

- HUISMAN T. A. Diffusion-weighted and diffusion tensor imaging of the brain, made easy. *Cancer Imaging* 2010; 10 Spec no A: S163-71.
- IAFFALDANO P., RUGGIERI M., VITERBO R. G., MASTRAPASQUA M., TROJANO M. The improvement of cognitive functions is associated with a decrease of plasma Osteopontin levels in Natalizumab treated relapsing multiple sclerosis. *Brain Behav Immun.* 2014 Jan; 35:176-81.
- IAFFALDANO P., VITERBO R. G., PAOLICELLI D., LUCCHESI G., PORTACCIO E., GORETTI B., DIRENZO V., D'ONGHIA M., ZOCCOLELLA S., AMATO M. P., TROJANO M. Impact of natalizumab on cognitive performances and fatigue in relapsing multiple sclerosis: a prospective, open-label, two years observational study. *PLoS One.* 2012; 7(4):e35843.
- IENTILE R., CURRO' M., FERLAZZO N., CONDELLO S., CACCAMO D., PISANI F. Homocysteine, vitamin determinants and neurological diseases. *Front Biosci (Schol Ed).* 2010 Jan 1; 2:359-72.
- INGLESE M., GROSSMAN R. I., FILIPPI M. Magnetic resonance imaging monitoring of multiple sclerosis lesion evolution. *J Neuroimaging.* 2005; 15(4 Suppl):S22-9.
- INTROZZI I., CANET L., ANDRÉS M. L., RICHARD'S M. Análisis de procesos de codificación y recuperación en pacientes con Esclerosis Múltiple. *Revista Chilena de Neuropsicología* 2007; 2:34-43.
- IRITI M., VITALINI S., FICO G., FAORO F. Neuroprotective herbs and foods from different traditional medicines and diets. *Molecules.* 2010; 15:3517-55.
- IZQUIERDO G., RUIZ-PEÑA J. L. Evaluación clínica de la esclerosis múltiple: cuantificación mediante la utilización de escalas. *Rev Neurol.* 2003; 36(2):145-152.
- JANARDHAN V., BAKSHI R. Quality of life in patients with multiple sclerosis: the impact of fatigue and depression. *J Neurol Sci.* 2002; 205:51-8.
- JOHNSON S. K., FREDERICK J., KAUFMAN M., MOUNTJOY B. A controlled investigation of bodywork in multiple sclerosis. *J Altern Complement Med.* 1999 Jun; 5(3):237-43.
- JORGE J., VAN DER ZWAAG W., FIGUEIREDO P. EEG-fMRI integration for the study of human brain function. *Neuroimage* 2014; 102(Pt 1):24-34.
- JUNG J., MORLET D., MERCIER B., CONFAVREUX C., FISCHER C. Mismatch negativity (MMN) in multiple sclerosis: An event-related potentials study in 46 patients. *Clin Neurophysiol.* 2006; 117(1): 85-93.
- KAIL R. Speed of information processing in patients with multiple sclerosis. *J Clin Exp Neuropsychol.* 1998; 20 (1):98-106.
- KANTARCI O., WINGERCHUK D. Epidemiology and natural history of multiple sclerosis: new insights. *Curr Opin Neurol.* 2006 Jun; 19(3):248-54. PubMed PMID: 16702830. Epub 2006/05/17. eng.
- KANTARCI O. H., WEINSHENKER B. G. Natural history of multiple sclerosis. *Neurol Clin.* 2005; 23(1):17-38.
- KAPELLER P., BREX P. A., CHARD D., *et al.* Quantitative ¹H MRS imaging 14 years after presenting with a clinically isolated syndrome suggestive of multiple sclerosis. *Mult Scler.* 2002; 8:207-10.
- KAPPOS L., MOERI D., RADUE E. W., SCHOETZAU A., SCHWEIKERT K., BARKHOF F., MILLER D., GUTTMANN C. R., WEINER H. L., GASPERINI C., *et al.*, Predictive value of gadolinium-enhanced magnetic resonance imaging for relapse rate and changes in disability or impairment in multiple sclerosis: a meta-analysis. *Gadolinium MRI Meta-analysis Group. Lancet* 1999; 353(9157):964-69.

- KAUNZNER U. W., GAUTHIER S. A. MRI in the assessment and monitoring of multiple sclerosis: an update on best practice. *Ther Adv Neurol Disord.* 2017; 10(6):247-61.
- KERN K. C., GOLD S. M., LEE B., MONTAG M., HORSFALL J., O'CONNOR M. F., SCOTTE N. L. Thalamic-hippocampal-prefrontal disruption in relapsing-remitting multiple sclerosis. *Neuroimage Clin.* 2015; 8:440-7.
- KESER I., KIRDI N., MERIC A., KURNE A. T., KARABUDAK R. Comparing routine neurorehabilitation program with trunk exercises based on Bobath concept in multiple sclerosis: pilot study. *J Rehabil Res Dev.* 2013;50(1):133-40.
- KHALIL M., ENZINGER C., LANGKAMMER C., PETROVIC K., LOITFELDER M., TSCHERNER M., JEHNA M., BACHMAIER G., WALLNER-BLAZEK M., ROPELE S., SCHMIDT R., FUCHS S., FAZEKAS F. Cognitive impairment in relation to MRI metrics in patients with clinically isolated syndrome. *Mult Scler.* 2011; 17(2):173-80.
- KHAN O., SHEN Y., CAON C., BAO F., CHING W., REZGAR M., BUCHEISTER A., HU J., LATIF Z., TSELIS A., LISAK R. Axonal metabolic recovery and potential neuroprotective effect of glatiramer acetate in relapsing-remitting multiple sclerosis. *Mult Scler.* 2005; 11:646-51.
- KIDD D., BARKHOF F., MCCONNELL R., ALGRA P. R., ALLEN I. V., REVESZ T. Cortical lesions in multiple sclerosis. *Brain* 1999; 122:17-26.
- KIDD P. M. Multiple sclerosis, an autoimmune inflammatory disease: prospects for its integrative management. *Altern Med Rev.* 2001 Dec; 6(6):540-66.
- KIISKI H., REILLY R. B., LONERGAN R., KELLY S., O'BRIEN M., KINSELLA K., BRAMHAM J., BURKE T., O'DONNCHADHA S., NOLAN H., HUTCHINSON M., TUBRIDY N., WHELAN R. Change in PASAT performance correlates with change in P3 ERP amplitude over a 12-month period in multiple sclerosis patients. *J Neurol Sci.* 2011a; 305(1-2):45-52.
- KIISKI H., WHELAN R., LONERGAN R., NOLAN H., KINSELLA K., HUTCHINSON M., TUBRIDY N., REILLY R. B. Preliminary evidence for correlation between PASAT performance and P3a and P3b amplitudes in progressive multiple sclerosis. *Eur J Neurol.* 2011b; 18:792-95.
- KIMISKIDIS V. K., PAPALIAGKAS V., SOTIRAKOGLU K., KOUVATSOU Z. K., KAPINA V. K., PAPADAKI E., TSIMOURTOU V., MASOURA E., KAZIS D. A., PAPAYIANNPOULOS S., GEROUKIS T., BOSTANJOPOULOU S. Cognitive event-related potentials in multiple sclerosis: Correlation with MRI and neuropsychological findings. *Mult Scler Relat Disord.* 2016; 10:192-97.
- KINSINGER S. W., LATTIE E., MOHR D. C. Relationship between depression, fatigue, subjective cognitive impairment, and objective neuropsychological functioning in patients with multiple sclerosis. *Neuropsychology.* 2010 Sep; 24(5):573-80. doi: 10.1037/a0019222.
- KLEIJNEN J., KNIPSCHILD P. Hyperbaric oxygen for multiple sclerosis. *Acta Neurol Scand.* 1995; 91:330-334.
- KNIPPENBERG S1., DAMOISEAUX J., BOL Y., HUPPERTS R., TAYLOR B. V., PONSONBY A. L., DWYER T., SIMPSON S., VAN DER MEI I. A. Higher levels of reported sun exposure, and not vitamin D status, are associated with less depressive symptoms and fatigue in multiple sclerosis. *Acta Neurol Scand.* 2014 Feb;129(2):123-31.
- KOCER B., UNAL T., NAZLIEL B., BIYIKLI Z., YESILBUDAK Z., KARAKAS S., IRKEC C. Evaluating sub-clinical cognitive dysfunction and event-related potentials (P300) in clinically isolated syndrome. *Neurol Sci.* 2008; 29(6):435-44.

- KOCH M., MOSTERT J., HEERINGS M., DE KEYSER J. Fatigue, depression and disability accumulation in multiple sclerosis: a cross-sectional study. *European Journal of Neurology* 2009, 16(3):348-352.
- KRAUS J. A., SCHUTZE C., BROKATE B., KROGER B., SCHWENDEMANN G., HILDEBRANDT H. Discriminant analysis of the cognitive performance profile of MS patients differentiates their clinical course. *J Neurol.* 2005; 252:808-13.
- KRUPP L. B., CHRISTODOULOU C., MELVILLE P., SCHERL W. F., PAI L. Y., MUENZ L. R., HE D., BENEDICT R. H., GOODMAN A., RIZVI S., SCHWID S. R., WEINSTOCK-GUTTMAN B., WESTERVELT H. J., WISHART H. Multicenter randomized clinical trial of donepezil for memory impairment in multiple sclerosis. *Neurology.* 2011 Apr 26;76(17):1500-7.
- KSIAZEK-WINIAREK D. J., SZPAKOWSKI P., GLABINSKI A. Neural Plasticity in Multiple Sclerosis: The Functional and Molecular Background. *Neural Plast* 2015; 2015:307175.
- KUHLE J., DISANTO G., DOBSON R., ADIUTORI R., BIANCHI L., TOPPING J., *et al.*, Conversion from clinically isolated syndrome to multiple sclerosis: A large multicentre study. *Mult Scler J.* [Internet]. 2015.
- KVAVILASHVILI L., ELLIS J. Varieties of intention: some distinction and classifications. En BRANDIMONTE M., EINSTEIN G. O., MCDANIEL M. A., eds. *Prospective memory: theory and applications.* Hillsdale, NJ: Erlbaum Associates, 1996.
- LABIANO-FONTCUBERTA A., MATO-ABAD V., ÁLVAREZ-LINERA J., HERNÁNDEZ-TAMAMES J. A., MARTÍNEZ-GINÉS M. L., ALADRO Y., AYUSO L., DOMINGO-SANTOS A., BENITO-LEÓN J. Normal-appearing brain tissue analysis in radiologically isolated síndrome using 3 T MRI. *Mult Scler.* 2016; 22(2):250-3.
- LAFOSSE J. M., MITCHELL S. M., CORBOY J. R., FILLEY C. M. The nature of verbal memory impairment in multiple sclerosis: a list-learning and meta-analytic study. *J Int Neuropsychol Soc.* 2013 Oct;19(9):995-1008. doi: 10.1017/S1355617713000957.
- LANDA L., KRPOUN Z., KOLAROVA M., KASPAREK T. Event-related Potentials and Their Applications. *Act Nerv Super* 2014; 56:17-23.
- LANDETE L., CASANOVA B. Deterioro cognitivo, formas clínicas y progresión en esclerosis múltiple. *Revista de Neurología* 2001; 32 (9):884-7.
- LANDRØ N. I., CELIUS E. G., SLETVOLD H. Depressive symptoms account for deficient information processing speed but not for impaired working memory in early phase multiple sclerosis (MS). *Journal of the Neurological Sciences* 2004; 217:211-216.
- LANGDON D. W., AMATO M. P., BORINGA J., BROCHET B., FOLEY F., FREDRIKSON S., HÄMÄLÄINEN P., HARTUNG H. P., KRUPP L., PENNER I. K., REDER A. T., BENEDICT R. H. Recommendations for a Brief International Cognitive Assessment for Multiple Sclerosis(BICAMS). *Mult Scler.* 2012 Jun;18(6):891-8. doi: 10.1177/1352458511431076. Epub 2011 Dec 21.
- LANGE R., VOLKMER M., HEESSEN C., LIEPERT J. Modafinil effects in multiple sclerosis patients with fatigue. *J Neurol.* 2009 Apr; 256(4):645-50.
- LANKHORST G. J., JELLES F., SMITS R. C., POLMAN C. H., KUIK D. J., PFENNINGS L. E., COHEN L., VAN DER PLOEG H. M., KETELAER P., VLEUGELS L. Quality of life in multiple sclerosis: the disability and impact profile (DIP). *J Neurol.* 1996 Jun; 243(6):469-74.

- LAZERON R. H., ROMBOUTS S. A., SCHELTENS P., POLMAN C. H., BARKHOF F. An fMRI study of planning-related brain activity in patients with moderately advanced multiple sclerosis. *Mult Scler.* 2004; 10(5):549-55.
- LEAVITT V. M., TOSTO G., RILEY C. S. Cognitive phenotypes in multiple Sclerosis. *Journal of Neurology* 2018 Jan; 1-5.
- LEBRUN C *et al.* "Cognitive function in radiologically isolated síndrome" *Mult Scler.* 2010; 16(8):919-25.
- LEE M., REDDY H., JOHANSEN-BERG H., PENDLEBURY S., JENKINSON M., SMITH S., *et al.*, The motor cortex shows adaptive functional changes to brain injury from multiple sclerosis. *Ann Neurol.* 2000; 47:606-13.
- LENZI D., CONTE A., MAINERO C., FRASCA V., FUBELLI F., TOTARO P., CARAMIA F., INGHILLERI M., POZZILLI C., PANTANO P. Effect of corpus callosum damage on ipsilateral motor activation in patients with multiple sclerosis: a functional and anatomical study. *Hum Brain Mapp.* 2007; 28(7):636-44.
- LEVIN N., ORLOV T., DOTAN S., ZOHARY E. Normal and abnormal fMRI activation patterns in the visual cortex after recovery from optic neuritis. *Neuroimage* 2006; 33:1161-68.
- LI Y., CHIARAVALLOTI N. D., HILLARY F. G., DELUCA J., LIU W. C., KALNIN A. J., RICKER J. Differential cerebellar activation on functional magnetic resonance imaging during working memory performance in persons with multiple sclerosis. *Arch Phys Med Rehabil.* 2004; 85(4):635-9.
- LINCOLN N. B., DENT A., HARDING J., *et al.*, Evaluation of cognitive assessment and cognitive intervention for people with multiple sclerosis. *J Neurol Neurosurg Psychiatry* 2002; 72:93-8.
- LOBENTANZ I. S., ASENBAUM S., VASS K., SAUTER C., KLÖSCH G., KOLLEGER H., KRISTOFERITSCH W., ZEITLHOFFER J. Factors influencing quality of life in multiple sclerosis patients: disability, depressive mood, fatigue and sleep quality. *Acta Neurol Scand.* 2004 Jul; 110(1):6-13.
- LOITFELDER M., FAZEKAS F., PETROVIC K., FUCHS S., ROPELE S., WALLNER-BLAZEK M., JEHNA M., ASPECK E., KHALIL M., SCHMIDT R., NEUPER C., ENZINGER C. Reorganization in cognitive networks with progression of multiple sclerosis: insights from fMRI. *Neurology* 2011; 76(6):526-33.
- LÓPEZ-GÓNGORA M., ESCARTÍN A., MARTÍNEZ-HORTA S., FERNÁNDEZ-BOBADILLA R., QUEROL L., ROMERO S., MAÑANAS M. À., RIBA J. Neurophysiological Evidence of Compensatory Brain Mechanisms in Early-Stage Multiple Sclerosis. *PLoS One* 2015; 10(8):e0136786.
- LÓPEZ-GÓNGORA M., QUEROL L., ESCARTÍN A. A one-year follow-up study of the Symbol Digit Modalities Test (SDMT) and the Paced Auditory Serial Addition Test (PASAT) in relapsing-remitting multiple sclerosis: an appraisal of comparative longitudinal sensitivity. *BMC Neurol.* 2015 Mar 22;15:40. doi: 10.1186/s12883-015-0296-2.
- LOPRESTI P. Glatiramer acetate guards against rapid memory decline during relapsing-remitting experimental autoimmune encephalomyelitis. *Neurochem Res.* 2015 Mar; 40(3):473-9.
- LOSSEFF N. A., WEBB S. L., O'RIORDAN J. I., PAGE R., WANG L., BARKER G. J., TOFTS P. S., McDONALD W. I., MILLER D. H., THOMPSON A. J. Spinal cord atrophy and disability in multiple sclerosis: a new reproducible and sensitive MRI method with potential to monitor disease progression. *Brain* 1996; 119:701-08.

- LOUAPRE C., PERLBARG V., GARCÍA-LORENZO D., URBANSKI M., BENALI H., ASSOUD R., *et al.*, Brain networks disconnection in early multiple sclerosis cognitive deficits: an anatomofunctional study. *Hum Brain Mapp* 2014; 35(9):4706-17.
- LÖVBLAD K. O., ANZALONE N., DÖRFLER A., ESSIG M., HURWITZ B., KAPPOS L., LEE S. K., FILIPPI M. MR imaging in multiple sclerosis: review and recommendations for current practice. *AJNR Am J Neuroradiol.* 2010; 31(6):983-9.
- LOVERA J. F., FROHMAN E., BROWN TR *et al.* Memantine for cognitive impairment in multiple sclerosis: a randomized placebo-controlled trial. *Mult Scler* 2010; 16(6):715-723.
- LOWE M. J., HORENSTEIN C., HIRSCH J. G., MARRIE R. A., STONE L., BHATTACHARYYA P. K., GASS A., PHILLIPS M. D. Functional pathway-defined MRI diffusion measures reveal increased transverse diffusivity of water in multiple sclerosis. *Neuroimage* 2006; 32(3):1127-33.
- LUBLIN F. D., REINGOLD S. C., COHEN J. A., CUTTER G. R., SØRENSEN P. S., THOMPSON A. J., *et al.*, Defining the clinical course of multiple sclerosis: The 2013 revisions. *Neurology* 2014; 83(3):278-86.
- LUTZ T., BELLENBERG B., SCHNEIDER R., WEILER F., KÖSTER O., LUKAS C. J Neuroimaging. Central atrophy early in multiple sclerosis: third ventricle volumetry versus planimetry. 2017 May; 27(3):348-354. doi: 10.1111/jon.12410.
- LYNCH S. G., PARMENTER B. A., DENNEY D. R. The association between cognitive impairment and physical disability in multiple sclerosis. *Mult Scler.* 2005; 11(4):469-476.
- MAINERO C., PANTANO P., CARAMIA F., POZZILLI C. fMRI evidence of brain reorganization during attention and memory tasks in multiple sclerosis. *Neuroimage* 2004; 21:858-67.
- MANTEROLA C., URRUTIA S *et al.*, Calidad de vida relacionada con salud. una variable resultado a considerar en investigación clínica. *Int. J. Morphol.* 2013; 31(4):1517-1523.
- MANTYNEN A., ROSTI-OTAJARVI E., KOIVISTO K., LILJA A., HUHTALA H., HAMALAINEN P. Neuropsychological rehabilitation does not improve cognitive performance but reduces perceived cognitive deficits in patients with multiple sclerosis: a randomised, controlled, multi-centre trial. *Mult Scler.* 2014;20(1):99-107.
- MARASESCU R., CEREZO M. Impairment of visuospatial/visuoconstructional skills in multiple sclerosis patients: the correlation with regional lesion load and subcortical atrophy. *Neurologia.* 2016; 31(3):169-175
- MARKOVIC-PLESE S., MCFARLAND H. F. Immunopathogenesis of the multiple sclerosis lesion. *Curr Neurol Neurosci Rep.* 2001; 1(3):257-62.
- MARRIE R. A., MILLER D. M., CHELUNE G. J., COHEN J. A. Validity and reliability of the MSQLI in cognitively impaired patients with multiple sclerosis. *Multiple Sclerosis.* 2003; 9(6):621-626.
- MASDEU J. C., QUINTO C., OLIVERA C., TENNER M., LESLIE D., VISINTAINER P. Open-ring imaging sign. Highly specific for atypical brain demyelination. *Neurology* 2000; 54(7):1427-33.
- MATTIOLI F., STAMPATORI C., ZANOTTI D., PARRINELLO G., CAPRA R. Efficacy and specificity of intensive cognitive rehabilitation of attention and executive functions in multiplesclerosis. *J Neurol Sci.* 2010; 288(1-2):101-5.
- MATTIOLI F., STAMPATORI C., BELLOMI F., SCARPAZZA C., CAPRA R. Natalizumab Significantly Improves Cognitive Impairment over Three Years in MS: Pattern of Disability Progression and Preliminary MRI Findings. *PLoS One.* 2015 Jul 6; 10(7):e0131803.

- MATTIOLI F., STAMPATORI C., CAPRA R. The effect of natalizumab on cognitive function in patients with relapsing-remitting multiple sclerosis: preliminary results of a 1-year follow-up study. *Neurol Sci.* 2011 Feb; 32(1):83-8.
- MCCABE M. P., MCKERN S., MCDONALD E., VOCALES L. M. *J Sex Marital Ther.* 2003; 29 : 305-21
- MCCADDON A. Vitamin B12 in neurology and ageing; clinical and genetic aspects. *Biochimie.* 2013 May; 95(5):1066-76.
- MCDONALD W. I., COMPSTON A., EDAN G., GOODKIN D., HARTUNG H. P., LUBLIN F. D., MCFARLAND H. F., PATY D. W., POLMAN C. H., REINGOLD S. C., SANDBERG-WOLLHEIM M., SIBLEY W., THOMPSON A., VAN DEN NOORT S., WEINSHENKER B. Y., WOLINSKY J. S. Recommended diagnostic criteria for multiple sclerosis: guidelines from the international panel on the diagnosis of multiple sclerosis. *Ann Neurol.* 2001; 50(1):121-27.
- MEIJER K. A., MUHLERT N., CERCIGNANI M., SETHI V., RON M. A., THOMPSON A. J., MILLER D. H., CHARD D., GEURTS J. J., CICCARELLI O. White matter tract abnormalities are associated with cognitive dysfunction in secondary progressive multiple sclerosis. *Mult Scler.* 2016; 22(11): 1429-37.
- MEZZAPESA D. M., ROCCA M. A., RODEGHER M., COMI G., FILIPPI M. Functional cortical changes of the sensorimotor network are associated with clinical recovery in multiple sclerosis. *Hum Brain Mapp.* 2008; 29(5): 562-73.
- MIGLIORE S., GHAZARYAN A., SIMONELLI I., PASQUALETTI P., *et al.* *Behav Neurol.* 2017; 2017: 7404289. doi: 10.1155 / 2017/7404289. Epub 2017 15 de agosto.
- MILLER D. H., BARKHOF F., FRANK J. A., *et al.* Measurement of atrophy in multiple sclerosis: pathological basis, methodological aspects and clinical relevance. *Brain* 2002; 125(Part 8):1676-95.
- MILLER D. H., BARKHOF F., NAUTA J. J. P. Gadolinium enhancement increases the sensitivity of MRI in detecting disease activity in multiple sclerosis. *Brain* 1993; 116:1077-94.
- MILLER D. H., GROSSMAN R. I., REINGOLD S. C., MCFARLAND H. F. The role of magnetic resonance techniques in understanding and managing multiple sclerosis. *Brain* 1998; 121(Pt 1):3-24.
- MILLER D. H. Imaging multiple sclerosis outside the brain. *Funct Neurol.* 2013; 28(2):72.
- MINDEN S. L., SCHIFFER R. B. Affective disorders in multiple sclerosis. Review and recommendations for clinical research. *Archives of Neurology* 1990; 47(1):98-104.
- MINNEBOO A., UITDEHAAG B. M., ADER H. J., BARKHOF F., POLMAN C. H., CASTELIJNS J. A. Patterns of enhancing lesion evolution in multiple sclerosis are uniform within patients. *Neurology* 2005 12; 65(1):56-61.
- MODICA C. M., BERGLAND N., DWYER M. G., RAMASAMY D. P., CARL E., ZIVADINOV R., BENEDICT R. H. Cognitive reserve moderates the impact of subcortical gray matter atrophy on neuropsychological status in multiple sclerosis. *Mult Scler.* 2016 Jan; 22(1):36-42.
- MODREGO P. J., PINA M. A., SIMÓN A., AZUARA M. C. The interrelations between disability and quality of life in patients with multiple sclerosis in the area of Bajo Aragón, Spain: a geographically based survey. *Neurorehabil Neural Repair* 2001; 15:27-35.
- MÖLLER F., POETTGEN J., BROEMEL F., NEUHAUS A., DAUMER M., HEESSEN C. HAGIL (Hamburg Vigil Study): a randomized placebo-controlled double-blind study with modafinil for treatment of fatigue in patients with multiple sclerosis. *Mult Scler.* 2011 Aug; 17(8):1002-9.

- MORGEN K., SAMMER G., COURTNEY S. M., *et al.* Evidence for a direct association between cortical atrophy and cognitive impairment in relapsing-remitting MS. *NeuroImage* 2006; 30(3):891-8.
- MORLET D., FISCHER C. MMN and novelty P3 in coma and other altered states of consciousness: a review. *Brain Topogr.* 2014; 27(4):467-79.
- MOROSO A., RUET A., LAMARGUE-HAMEL D., MUNSCH F., DELOIRE M., OUALLET J. C., CUBIZOLLE S., CHARRÉ-MORIN J., SAUBUSSE A., TOURDIAS T., DOUSSET V., BROCHET B. Preliminary evidence of the cerebellar role on cognitive performances in clinically isolated syndrome. *J Neurol Sci.* 2018 Febrero 2013; 385:1-6. doi: 10.1016
- MORROW S. A., O'CONNOR P. W., POLMAN C. H., GOODMAN A. D., KAPPOS L., LUBLIN F. D., RUDICK R. A., JURGENSEN S., PAES D., FORRESTAL F., BENEDICT R. H. Evaluation of the symbol digit modalities test (SDMT) and MS neuropsychological screening questionnaire (MSNQ) in natalizumab-treated MS patients over 48 weeks. *Mult Scler.* 2010 Nov; 16(11):1385-92.
- MORROW S. A., ROSEHART H. Effects of single dose mixed amphetamine salts--extended release on processing speed in multiple sclerosis: a double blind placebo controlled study. *Psychopharmacology (Berl)* 2015 Dec; 232(23):4253-9.
- MORROW S. A., SMERBECK A., PATRICK K., COOKFAIR D., WEINSTOCK-GUTTMAN B., BENEDICT R. H. Lisdexamfetamine dimesylate improves processing speed and memory in cognitively impaired MS patients: a phase II study. *J Neurol.* 2013 Feb; 260(2):489-97.
- MOSTERT J. P., KOCH M.W., STEEN C., HEERSEMA D. J., DE GROOT J. C., DE KEYSER J. T2 lesions and rate of progression of disability in multiple sclerosis. *Eur J Neurol.* 2010; 17(12):1471-5.
- NÄÄTÄNEN R., KUJALA T., ESCERA C., BALDEWEG T., KREEGIPUU K., CARLSON S., PONTON C. The mismatch negativity (MMN)--a unique window to disturbed central auditory processing in ageing and different clinical conditions. *Clin Neurophysiol.* 2012; 123(3):424-58.
- NAMJOOYAN F., GHANAVATI R., MAJDINASAB N., JOKARI S., JANBOZORGI M. Uses of complementary and alternative medicine in multiple sclerosis. *J Tradit Complement Med.* 2014 Jul; 4(3):145-52.
- NARAYANA P. A., GOVINDARAJAN K. A., GOEL P., DATTA S., LINCOLN J. A., COFIELD S. S., CUTTER G. R., LUBLIN F. D., WOLINSKY J. S. Regional cortical thickness in relapsing remitting multiple sclerosis: A multi-center study. *Neuroimage Clin.* 2012; 2:120-31.
- NARAYANA P. A. Magnetic Resonance Spectroscopy in the Monitoring of Multiple Sclerosis. *J Neuroimaging* 2005; 15(4 Suppl):S46-S57.
- NEBEL K., WIESE H., SEYFARTH J., *et al.* Activity of attention related structures in multiple sclerosis. *Brain Res.* 2007; 1151:150-160.
- NEBEL K., WIESE H., SEYFARTH J., GIZEWSKI E. R., STUDE P., DIENER H. C., LIMMROTH V. Activity of attention related structures in multiple sclerosis patients. *Brain Res.* 2007; 1151:150-60.
- NEJATI S., RAJEZI ESFAHANI S., RAHMANI S., AFROOKHTEH G., HOVEIDA S. The Effect of Group Mindfulnessbased Stress Reduction and Consciousness Yoga Program on Quality of Life and Fatigue Severity in Patients with MS. *J Caring Sci.* 2016 Dec 1; 5(4):325-335.
- NEWTON M. R., BARRETT G., CALLANAN M. M., TOWELL A. D. *Brain* 1989; 112(Pt 6):1637-60.
- NICCOLAI C., PORTACCIO E., GORETTI B., HAKIKI B., GIANNINI M., PASTÒ L., RIGHINI I., FALAUTANO M *et al.*, A comparison of the brief international cognitive assessment for multiple sclerosis and the brief repeatable battery in multiple sclerosis patients. *BMC Neurol.* 2015 Oct 15; 15:204. doi: 10.1186/s12883-015-0460-8.

- NOCENTINI U., BOZZALI M., SPANÒ B., CERCIGNANI M., SERRA L., BASILE B., MANNU R., CALTAGIRONE C., DE LUCA J. Exploration of the relationships between regional grey matter atrophy and cognition in multiple sclerosis. *Brain Imaging Behav.* 2014; 8(3):378-86.
- NOCENTINI U., ROSSINI P. M., CARLESIMO G. A., GRACEFFA A., GRASSO M. G., LUPOI D., OLIVERI M., ORLACCHIO A., POZZILLI C., RIZZATO B., CALTAGIRONE C. Patterns of cognitive impairment in secondary progressive stable phase of multiple sclerosis: correlations with MRI findings. *Eur Neurol.* 2001; 45:11-8.
- NORTVEDT M. W., RIISE T., FRUGARD J., MOHN J., BAKKE A., SKAR A. B., *et al.*, Prevalence of bladder, bowel and sexual problems among multiple sclerosis patients two to five years after diagnosis. *Mult Scler.* 2007; 13:106-12.
- NORTVEDT M. W., RIISE T., MYHR K. M., LANDTBLOM A. M., BAKKE A., NYLAND H. I. Reduced quality of life among multiple sclerosis patients with sexual disturbance and bladder dysfunction. *Mult Scler.* 2001; 7:231-5.
- NOSEWORTHY J. H. Clinical scoring methods for multiple sclerosis. *Ann Neurol* 1994; 36 (Suppl):S80-5.
- NSAMENANG S. A., HIRSCH J. K., TOPCIU R., GOODMAN A. D., DUBERSTEIN P. R. The interrelations between spiritual well-being, pain interference and depressive symptoms in patients with multiple sclerosis. *J Behav Med.* 2016 Apr; 39(2):355-63.
- O'BRIEN A., GAUDINO-GOERIN E *et al.* Relationship of the Multiple Sclerosis Neuropsychological Questionnaire (MSNQ) to functional, emotional, and neuropsychological outcomes. *Archives of Clinical Neuropsychology* 2007 November; 22(8):933-948.
- O'CARROLL C. B., WOODRUFF B. K., LOCKE D. E., HOFFMAN-SNYDER C. R., WELLIK K. E., THAERA G. M., DEMAERSCHALK B. M., WINGERCHUK D. M. Is donepezil effective for multiple sclerosis-related cognitive dysfunction? A critically appraised topic. *Neurologist.* 2012 Jan; 18(1):51-4.
- OGAWA S., LEE T. M., KAY A. R., TANK D. W. Brain magnetic resonance imaging with contrast dependent on blood oxygenation. *Proc Natl Acad Sci. USA* 1990; 87(24):9868-72.
- OKEN B. S., KISHIYAMA S., ZAJDEL D., BOURDETTE D., CARLSEN J., HAAS M., HUGOS C., KRAEMER D. F., LAWRENCE J., MASS M. Randomized controlled trial of yoga and exercise in multiple sclerosis. *Neurology* 2004 Jun 8; 62(11):2058-64.
- OKUDA D. T., MOWRY E. M., CREE B. A., CRABTREE E. C., GOODIN D. S., WAUBANT E., PELLETIER D. Asymptomatic spinal cord lesions predict disease progression in radiologically isolated syndrome. *Neurology* 2011; 76:686-692.
- OREJA-GUEVARA C., LUBRINI G. Deterioro cognitivo en esclerosis múltiple. *Revista Española de Esclerosis Múltiple* 2009 Septiembre; 12.
- OZAKBAS S., CAGIRAN I., ORMECI B., IDIMAN E. Correlations between multiple sclerosis functional composite, expanded disability status scale and health-related quality of life during and after treatment of relapses in patients with multiple sclerosis. *J Neurol Sci.* 2004 Mar 15; 218(1-2):3-7.
- PAGANI E., BAMMER R., HORSFIELD M. A., ROVARIS M., GASS A., CICCARELLI O., FILIPPI M. Diffusion MR imaging in multiple sclerosis: technical aspects and challenges. *AJNR Am J Neuroradiol.* 2007; 28(3):411-20.

- PAGANI E., ROCCA M. A., GALLO A., ROVARIS M., MARTINELLI V., COMI G., FILIPPI M. Regional brain atrophy evolves differently in patients with multiple sclerosis according to clinical phenotype. *AJNR Am J Neuroradiol.* 2005; 26(2):341-6.
- PAN J. W., KRUPP L. B., ELKINS L. E., COYLE P. K. Cognitive dysfunction lateralizes with NAA in multiple sclerosis. *Appl Neuropsychol.* 2001; 8(3):155-60.
- PANTANO P., IANNETTI G. D., CARAMIA F., MAINERO C., DI LEGGE S., BOZZAO L., POZZILLI C., LENZI G. L. Cortical motor reorganization after a single clinical attack of multiple sclerosis. *Brain* 2002; 125(Pt 7):1607-15.
- PANTANO P., MAINERO C., LENZI D., CARAMIA F., IANNETTI G. D., PIATTELLA M. C., PESTALOZZA I., DI LEGGE S., BOZZAO L., POZZILLI C. A longitudinal fMRI study on motor activity in patients with multiple sclerosis. *Brain* 2005; 128(9):2146-53.
- PAPATHANASIOU, A., MESSINIS L., GEORGIU V. Cognitive Impairment in relapsing remitting and secondary progressive multiple sclerosis patients: efficacy of a computerized cognitive screening battery. *ISRN Neurology* 2014; Article ID 151379., 7 pages.
- PARISI L., ROCCA M. A., MATTIOLI F., COPETTI M., CAPRA R., VALSASINA P., *et al.* Changes of brain resting state functional connectivity predict the persistence of cognitive rehabilitation effects in patients with multiple sclerosis. *Mult Scler.* 2013; 20(6):686-94.
- PASTÒ L., PORTACCIO E., GORETTI B., GHEZZI A., LORI S., HAKIKI B., GIANNINI M., RIGHINI I., RAZZOLINI L., NICCOLAI C., MOIOLA L., FALAUTANO M., SIMONE M., VITERBO R. G., PATTI F., CILIA S., POZZILLI C., BIANCHI V., ROSCIO M., MARTINELLI V., COMI G., TROJANO M., AMATO MP; MS Study Group of the Italian Neurological Society. The cognitive reserve theory in the setting of pediatric-onset multiple sclerosis. *Mult Scler.* 2016 Feb 11.
- PATTI F., AMATO M., TROJANO M., *et al.* Cognitive impairment and its relation with disease measures in mildly disabled patients with relapsing-remitting multiple sclerosis: baseline results from the Cognitive Impairment in Multiple Sclerosis (COGIMUS) study. *Mult Scler.* 2009; 15:779-788. 10.1177/1352458509105544.
- PATTI F., AMATO M. P., TROJANO M., BASTIANELLO S., TOLA M. R., GORETTI B., CANIATTI L., DI MONTE E., FERRAZZA P., BRESCIA MORRA V., LO FERMO S., PICCONI O., LUCCICHENTI G; COGIMUS STUDY GROUP. Cognitive impairment and its relation with disease measures in mildly disabled patients with relapsing-remitting multiple sclerosis: baseline results from the Cognitive Impairment in Multiple Sclerosis (COGIMUS) study. *Mult Scler.* 2009;15(7):779-88.
- PATTI F., AMATO M. P., BASTIANELLO S., CANIATTI L., DI MONTE E., FERRAZZA P., GORETTI B., GALLO P., MORRA V. B., LO FERMO S., PICCONI O., TOLA M. R., TROJANO M; COGIMUS STUDY GROUP. Effects of immunomodulatory treatment with subcutaneous interferon beta-1a on cognitive decline in mildly disabled patients with relapsing-remitting multiple sclerosis. *Mult Scler.* 2010 Jan; 16(1):68-77.
- PATTI F., AMATO M. P., TROJANO M., BASTIANELLO S., TOLA M. R., PICCONI O., CILIA S., COTTON S., CENTONZE D., GASPERINI C; COGIMUS STUDY GROUP. Quality of life, depression and fatigue in mildly disabled patients with relapsing-remitting multiple sclerosis receiving subcutaneous interferon beta-1a: 3-year results from the COGIMUS (COGNITIVE IMPAIRMENT IN MULTIPLE SCLEROSIS) study. *Multiple Sclerosis* 2011 Aug; 17(8):991-1001.

- PATTI F, AMATO M. P., TROJANO M., BASTIANELLO S., TOLA M. R., PICCONI O., CILIA S., COTTONE S., GRIMALDI LM; COGIMUS STUDY GROUP. Longitudinal changes in social functioning in mildly disabled patients with relapsing-remitting multiple sclerosisreceiving subcutaneous interferon β -1a: results from the COGIMUS (COGNitive Impairment in MULTiple Sclerosis) study (II). *Qual Life Res.* 2012 Sep; 21(7):1111-21.
- PATTI F, DE STEFANO M., LAVORGNA L., MESSINA S., CHISARI C. G., IPPOLITO D., *et al.*, Lesion Load May Predict Long-Term Cognitive Dysfunction in Multiple Sclerosis Patients. *PLoS One* 2015; 10(3): e0120754.
- PATTI F, MORRA V. B., AMATO M. P., TROJANO M., BASTIANELLO S., TOLA M. R., COTTONE S., PLANT A., PICCONI O; COGIMUS Study Group. Subcutaneous interferon β -1a may protect against cognitive impairment in patients with relapsing-remitting multiple sclerosis: 5-year follow-up of the COGIMUS study. *PLoS One.* 2013 Aug 30;8(8):e74111.
- PEDULLÀ L., BRICHETTO G., TACCHINO A., VASSALLO C., ZARATIN P., BATTAGLIA M. A., BONZANO L., BOVE M. Adaptive vs. non-adaptive cognitive training by means of a personalized App: a randomized trial in people with multiple sclerosis. *J Neuroeng Rehabil.* 2016 Oct 4; 13(1):88.
- PELOSI L., GEESKEN J. M., HOLLY M., HAYWARD M., BLUMHARDT L. D. Working memory impairment in early multiple sclerosis. Evidence from an event-related potential study of patients with clinically isolated myelopathy. *Brain*1997; 120(Pt 11):2039-58.
- PENG Y., LI S., ZHUANG Y., LIU X., WU L., GONG H., LIU D., ZHOU F. Density abnormalities in normal-appearing gray matter in the middle-aged brain with white matter hyperintense lesions: a DARTEL-enhanced voxel-based morphometry study. *Clin Interv Aging* 2016; 11: 615-22.
- PENNER I. K., OPWIS K., KAPPOS L. Relation between functional brain imaging, cognitive impairment and cognitive rehabilitation in patients with multiple sclerosis. *J Neurol.* 2007; 254 (Suppl2):II53-7.
- PENNER I. K., RAUSCH M., KAPPOS L., OPWIS K., RADU E. W. Analysis of impairment related functional architecture in MS patients during performance of different attention tasks. *J Neurol.* 2003; 250: 461-72.
- PÉREZ DE COLOSÍA V., MARTÍN MUÑOZ M., SANZ DE LA TORRE J. C., PLASENCIA BLANCO A., DUQUE SAN JUAN P. Deterioro cognitivo en pacientes con esclerosis múltiple: Estudio con la Batería Neuropsicológica Breve (BNB). *Neurología* 2008; 23 (10):617-72.
- PERKAL H., PORCEL J., NOS C., FREITAS A., GUILLAMÓN I., AYMERICH M., MONTALBÁN X. Validación de la versión española del RAYS [póster]. LVII Reunión Científica de la Sociedad Española de Neurología; Barcelona, (novembre 2005). *Neurología* 2005; 20(9):533.
- PERRINS D. J. D., JAMES P. B. Long-term study on hyperbaric oxygenation for multiple sclerosis. Bedford., United Kingdom: Federation of Multiple Sclerosis Therapy Centers; 1996: website www.ms-selfhelp.org.
- PFFENNINGS L. E., VAN DER PLOEG H. M., COHEN L., BRAMSEN I., POLMAN C. H., LANKHORST G. J., VLEUGELS L. A health-related quality of life questionnaire for multiple sclerosis patients. *Acta Neurol Scand.* 1999 Sep; 100(3):148-55.
- PHILLIPS., C., BAKTIR., M. A., SRIVASTAN, M. Y SALEHI, A. Neuroprotective effects of physical activity on the brain: A closer look at trophic factor signaling. *Frontiers in Cellular Neuroscience* 2014; 8:170.

- PICTON T. W. The P300 wave of the human event-related potential. *J Clin Neurophysiol.* 1992; 9:456-79.
- PIRAS M. R., MAGNANO I., CANU E. D., PAULUS K. S., SATTÀ W. M., SODDU A., CONTI M., ACHENE A., SOLINAS G., AIELLO I. Longitudinal study of cognitive dysfunction in multiple sclerosis: neuropsychological, neuroradiological, and neurophysiological findings. *J Neurol Neurosurg Psychiatry* 2003; 74(7):878-85.
- PIRKO I., M. D; LUCCHINETTI C. F., SRIRAM S., BAKSHI R. Gray matter involvement in multiple sclerosis. *Neurology* 2007; 68:634-42.
- PITTERI *et al.* Cognitive impairment predicts disability progression and cortical thinning in MS: An 8 year study” *M. Mult Scler.* 2017; 23(6):848-854.
- PITTOCK S. J., MAYR W. T., MCCLELLAND R. L., JORGENSEN N. W., WEIGAND S. D., NOSEWORTHY J. H., *et al.*, Quality of life is favorable for most patients with multiple sclerosis: a population-based cohort study. *Arch Neurol.* 2004; 61:679-86.
- PLANCHE V., GIBELIN M., CREGUT D., PEREIRA B., CLAVELOU P. Cognitive impairment in a population-based study of patients with multiple sclerosis: differences between late relapsing-remitting, secondary progressive and primary progressive multiple sclerosis. *Eur J Neurol* (2016) 23:282-9. doi:10.1111/ene. 12715
- PLASSMAN B. L., WILLIAMS JW JR., BURKE J. R., HOLSINGER T., BENJAMIN S. Systematic review: factors associated with risk for and possible prevention of cognitive decline in later life. *Ann Intern Med.* 2010 Aug 3;153(3):182-93.
- PLISKIN N. H., HAMER D. P., GOLDSTEIN D. S., TOWLE V. L., REDER A. T., NORONHA A. Improved delayed visual reproduction test performance in multiple sclerosis patients receiving interferon beta-1b. *Neurology* 1996 47:1463-1468.
- POKRYSZKO-DRAGAN A., DZIADKOWIAK E., ZAGRAJEK M., SLOTWINSKI K., GRUSZKA E., BILINSKA M., PODEMSKI R Cognitive performance, fatigue and event-related potentials in patients with clinically isolated syndrome. *Clin Neurol Neurosurg.* 2016; 149:68-74.
- POKRYSZKO-DRAGAN A., ZAGRAJEK M., SLOTWINSKI K., GRUSZKA E., BILINSKA M., PODEMSKI R. Neuropsychological testing and event-related potentials in the assessment of cognitive performance in the patients with multiple sclerosis: a pilot study. *Clin Neurol Neurosurg.* 2009; 111(6):503-6.
- POLICH J., COMERCHERO M. D. P3a from visual stimuli: typicality, task, and topography. *Brain Topogr.* 2003; 15(3):141-52.
- POLICH J., HERBST K. L. P300 as a clinical assay: rationale, evaluation, and findings. *Int J Psychophysiol.* 2000; 38(1):3-19.
- POLICH J., ROMINE J. S., SIPE J. C., AUNG M., DALESSIO D. J. P300 in multiple sclerosis: a preliminary report. *Int J Psychophysiol.* 1992; 12(2):155-63.
- POLICH J. Updating P300: an integrative theory of P3a and P3b. *Clin Neurophysiol.* 2007; 118(10):2128-48.
- POLMAN C. H., REINGOLD S. C., BANWELL B., CLANET M., COHEN J. A., FILIPPI M., FUJIHARA K., HAVRDOVA E., HUTCHINSON M., KAPPOS L., LUBLIN F. D., MONTALBAN X., O'CONNOR P., SANDBERG-WOLLHEIM M., THOMPSON A. J., WAUBANT E., WEINSHENKER B., WOLINSKY J. S.

- Diagnostic criteria for multiple sclerosis: 2010 revisions to the McDonald criteria. *Ann Neurol*. 2011; 69(2):292-302.
- POONAWALLA A. H., HASAN K. M., GUPTA R. K., AHN C. W., NELSON F., WOLINSKY J. S., NARAYANA P. A. Diffusion-tensor MR imaging of cortical lesions in multiple sclerosis. *Radiology* 2008; 246(3):880-6.
- POPESCU V., SCHOONHEIM M. M., VERSTEEG A., CHATURVEDI N., JONKER M., XAVIER DE MENEZES R., GALLINDO GARRE F., UITDEHAAG B. M., BARKHOF F., VRENKEN H. Grey Matter Atrophy in Multiple Sclerosis: Clinical Interpretation Depends on Choice of Analysis Method. *PLoS One* 2016; 11(1):e0143942.
- PORTACCIO E., GORETTI B., ZIPOLI V., HAKIKI B., GIANNINI M., PASTÒ L., RAZZOLINI L., AMATO M. P. Cognitive rehabilitation in children and adolescents with multiple sclerosis. *Neurol Sci*. 2010 Nov; 31(Suppl 2):S275-8.
- PORTACCIO E., STROMILLO M. L., GORETTI B., HAKIKI B., GIORGIO A., ROSSI F., DE LEUCIO A., DE STEFANO N., AMATO M. P. Natalizumab may reduce cognitive changes and brain atrophy rate in relapsing-remitting multiple sclerosis--a prospective, non-randomized pilot study. *Eur J Neurol*. 2013 Jun; 20(6):986-90.
- POSNER M. I., DEHAENE S. Attentional networks. *Trends Neurosci*. 1994; 17(2):75-9.
- POSNER M. I., PETERSEN SE: The attention system of the human brain. *Annu Rev Neurosci*. 1990; 13:25-42.
- POTAGAS C., GIOGKARARKI E., *et al.*, Cognitive impairment in different MS subtypes and clinically isolated syndromes. *J Neurol Sci*. 2008; 267:100-106.
- PRAKASH R. S., SNOOK E. M., ERICKSON K. I., COLCOMBE S. J., VOSS M. W., MOTL R. W., KRAMER A. F. Cardiorespiratory fitness: A predictor of cortical plasticity in multiple sclerosis. *NeuroImage* 2007; 34:1238-1244.
- PRIETO L., ALONSO J., ANTÓ J. M. La versión española del SF-36 Health Survey: un instrumento para la medida de los resultados clínicos. *Med Clin (Barc)* 1995; 104:771-6.
- PRIETO L., ALONSO J., FERRER M., ANTÓ J. M. Health Status of COPD patients: are results of the SF-36 Health Survey and the Nottingham Health Profile comparable? *J Clin Epidemiol*. 1997; 50:463-73.
- PROTAGAS *et al.* Cognitive impairment in different MS subtypes and Clínicamente aislados síndromes". *J Neurol Sci*. 2008; 267(1-2):100-6
- PUSSWALD G., MILDNER C., ZEBENHOLZER K., AUFF E., LEHRNER J. A neuropsychological rehabilitation program for patients with Multiple Sclerosis based on the model of the ICF. *NeuroRehabilitation*. 2014; 35(3):519-27.
- RAHN K. A., WATKINS C. C., ALT J., RAIS R., STATHIS M., GRISHKAN I., CRAINICEAU C. M., POMPER M. G., ROJAS C., PLETNIKOV M. V., CALABRESI P. A., BRANDT J., BARKER P. B., SLUSHER B. S., KAPLIN A. I. Inhibition of glutamate carboxypeptidase II (GCPII) activity as a treatment for cognitive impairment in multiple sclerosis. *Proc Natl Acad Sci. USA* 2012 Dec 4; 109(49):20101-6.
- RANJEVA J. P., AUDOIN B., AU DUONG M. V., IBARROLA D., CONFORT-GOUNY S., MALIKOVA I., SOULIER E., VIOUT P., ALI-CHÉRIF A., PELLETIER J., COZZONE P. Local tissue damage assessed with statistical mapping analysis of brain magnetization transfer ratio: relationship with function-

- al status of patients in the earliest stage of multiple sclerosis. *AJNR Am J Neuroradiol.* 2005; 26(1):119-27.
- RAO A. B., RICHERT N., HOWARD T., LEWIS B. K., BASH C. N., MCFARLAND H. F., FRANK J. A. Methylprednisolone effect on brain volume and enhancing lesions in MS before and during IFNbeta-1b. *Neurology* 2002; 59:688-94.
- RAO S. M. A manual for the brief, repeatable battery of neuropsychological tests in multiple sclerosis. New York: National Multiple Sclerosis Society., 1991.
- RAO S. M., GRAFMAN J., DIGUILIO D. Memory dysfunction in multiple sclerosis: its relation to working memory, semantic encoding and implicit learning. *Neuropsychology* 1993; 7:364-374.
- RAO S. M., LEO G. J., BERNARDIN L., UNVERZAGT F. Cognitive dysfunction in multiple sclerosis. I. Frequency, patterns, and prediction. *Neurology* 1991 May; 41(5):685-91.
- RAO S. M., LEO G. J., ELLINGTON L., NAUERTZ T., BERNARDIN L., UNVERZAGT F. Cognitive dysfunction in multiple sclerosis: employment and social functioning *neurology* 1991 May; 41(5):692-6.
- RAO S. M., LEO G. J., HAUGHTON V. M., ST. AUBIN-FAUBERT P., BERNARDIN L. Correlation of magnetic resonance imaging with neuropsychological testing in multiple sclerosis. *Neurology* 1989; 39:161-66.
- RAO S. M., MARTIN A. L., HUELIN R., WISSINGER E., KHANKHEL Z., KIM E., FAHRBACH K. Correlations between MRI and Information Processing Speed in MS: A Meta-Analysis. *Mult Scler Int.* 2014; 2014:975803.
- REDDY H., NARAYANAN S., MATTHEWS P. M., HOGE R. D., PIKE G. B., DUQUETTE P., ANTEL J., ARNOLD D. L. Relating axonal injury to functional recovery in MS. *Neurology* 2000; 54:236-39.
- RENELL P. G., JENSEN F., HENRY J. D. Prospective memory in multiple sclerosis. *J Int Neuropsychol Sci.* 2007; 13:410-6.
- REULING I. E. W., POLMAN C. H. Differences in cognitive impairment of relapsing remitting, secondary and primary progressive MS. *Neurology* 2004; 63:335-339.
- REUTER F., ZAARAOUI W., CRESPIY L., FAIVRE A., RICO A., MALIKOVA I., SOULIER E., VIOUT P., RANJEVA J. P., PELETTIER J., AUDOIN B. Frequency of cognitive impairment dramatically increases during the first 5 years of multiple sclerosis. *J Neurol Neurosurg Psychiatry.* 2011 Oct; 82(10):1157-9. doi: 10.1136/jnnp.2010.213744. Epub 2010 Oct 22.
- RIAZI A., HOBART J. C., LAMPING D. L., FITZPATRICK R., THOMPSON A. J. Evidence-based measurement in multiple sclerosis: the psychometric properties of the physical and psychological dimensions of three quality of life rating scales. *RIAZI, A. et al. Mult Scler.* 2003; 9:411-419.
- RIAZI A., HOBART J. C., LAMPING D. L., *et al.* Multiple Sclerosis Impact Scale (MSIS-29): reliability and validity in hospital based samples. *Journal of Neurology, Neurosurgery & Psychiatry* 2002; 73:701-704.
- RICCITELLI G., ROCCA M. A., PAGANI E., RODEGHER M. E., ROSSI P., FALINI A., COMI G., FILIPPI M. Cognitive impairment in multiple sclerosis is associated to different patterns of gray matter atrophy according to clinical phenotype. *Hum Brain Mapp.* 2011; 32(10):1535-43.
- RIVERA-NAVARRO J., BENITO-LEÓN J., MORALES-GONZÁLEZ J. M. Hacia la búsqueda de dimensiones más específicas en la medición de la calidad de vida en la esclerosis múltiple. *Rev Neurol* 2001; 32:705-13.

- ROCCA M. A., CECCARELLI A., RODEGHER M., MISCI P., RICCITELLI G., FALINI A., COMI G., FILIPPI M. Preserved brain adaptive properties in patients with benign multiple sclerosis. *Neurology* 2010a; 74(2):142-49.
- ROCCA M. A., CERCIGNANI M., IANNUCCI G., *et al.* Weekly diffusion-weighted imaging of normal-appearing white matter in MS. *Neurology* 2000; 55:882-84.
- ROCCA M. A., COLOMBO B., FALINI A., GHEZZI A., MARTINELLI V., SCOTTI G., COMI G., FILIPPI M. Cortical adaptation in patients with MS: a cross-sectional functional MRI study of disease phenotypes. *Lancet Neurol.* 2005; 4:618-26.
- ROCCA M. A., DE MEO E., FILIPPI M. Functional MRI in investigating cognitive impairment in multiple sclerosis. *Acta Neurol Scand* 2016; 134(Suppl 200):39-46.
- ROCCA M. A., FALINI A., COLOMBO B., SCOTTI G., COMI G., FILIPPI M. Adaptive functional changes in the cerebral cortex of patients with nondisabling multiple sclerosis correlate with the extent of brain structural damage. *Ann Neurol.* 2002b; 51(3):330-39.
- ROCCA M. A., GALLO A., COLOMBO B., FALINI A., SCOTTI G., COMI G., FILIPPI M. Pyramidal tract lesions and movement-associated cortical recruitment in patients with MS. *Neuroimage* 2004; 23(1):141-47.
- ROCCA M. A., GAVAZZI C., MEZZAPESA D. M., FALINI A., COLOMBO B., MASCALCHI M., SCOTTI G., COMI G., FILIPPI M. A functional magnetic resonance imaging study of patients with secondary progressive multiple sclerosis. *Neuroimage* 2003; 19(4):1770-7.
- ROCCA M. A., HICKMAN S. J., BÖ L., AGOSTA F., MILLER D. H., COMI G., FILIPPI M. Imaging the optic nerve in multiple sclerosis. *Mult Scler.* 2005; 11(5):537-541.
- ROCCA M. A., MATTHEWS P. M., CAPUTO D., GHEZZI A., FALINI A., SCOTTI G., COMI G., FILIPPI M. Evidence for widespread movement-associated functional MRI changes in patients with primary-progressive multiple sclerosis. *Neurology* 2002a; 58(6):866-72.
- ROCCA M. A., VALSASINA P., ABSINTA M., *et al.*, Default-mode network dysfunction and cognitive impairment in progressive MS. *Neurology* 2010b;74:1252-9.
- ROCCA M. A., VALSASINA P., CECCARELLI A., ABSINTA M., GHEZZI A., RICCITELLI G., PAGANI E., FALINI A., COMI G., SCOTTI G., FILIPPI M. Structural and functional MRI correlates of Stroop control in benign MS. *Hum Brain Mapp* 2009; 30:276-90.
- ROCCA M. A., VALSASINA P., HULST H. E., ABDEL-AZIZ K., ENZINGER C., GALLO A., PARETO D., RICCITELLI G., MUHLERT N., CICCARELLI O., BARKHOF F., FAZEKAS F., TEDESCHI G., ARÉVALO M. J., FILIPPI M; MAGNIMS fMRI Study Group. Functional correlates of cognitive dysfunction in multiple sclerosis: A multicenter fMRI Study. *Hum Brain Mapp.* 2014; 35(12):5799-814.
- ROMBERG A., VIRTANEN A., RUUTIAINEN J. Long-term exercise improves functional impairment but not quality of life in multiple sclerosis. *J Neurol.* 2005 Jul; 252(7):839-45.
- ROMERO K., PAVISIAN B., STAINES W. R., FEINSTEIN A. Multiple sclerosis, cannabis, and cognition: A structural MRI study. *Neuroimage Clin.* 2015 Apr 9; 8:140-7.
- ROMERO K., SHAMMI P., FEINSTEIN A. Neurologists' accuracy in predicting cognitive impairment in multiple sclerosis. *Mult Scler Relat Disord.* 2015 Jul; 4(4):291-5. Epub 2015 May 27.
- ROOSEDAAL S. D., HULST H. E., VRENKEN H., FEENSTRA H. E., CASTELIJNS J. A., POWWELS P. J., BARKHOF F., GEURTS J. J. Structural and functional hippocampal changes in multiple sclerosis patients with intact memory function. *Radiology* 2010b; 255(2):595-604.

- ROSENDAAL S. D., SCHOONHEIM M. M., HULST H. E., SANZ-ARIGITA E. J., SMITH S. M., GEURTS J. J., BARKHOF F. Resting state networks change in clinically isolated syndrome. *Brain* 2010a; 133(Pt 6):1612-21.
- ROSTI-OTAJARVI E., MANTYNEN A., KOIVISTO K., HUHTALA H., HAMALAINEN P. Neuropsychological rehabilitation has beneficial effects on perceived cognitive deficits in multiple sclerosis during nine-month follow-up. *J Neurol Sci.* 2013a; 334(1-2):154-60.
- ROSTI-OTAJARVI E., MANTYNEN A., KOIVISTO K., HUHTALA H., HAMALAINEN P. Patient-related factors may affect the outcome of neuropsychological rehabilitation in multiplesclerosis. *J Neurol Sci.* 2013b; 334(1-2):106-11.
- ROVARIS M., BOZZALI M., SANTUCCIO G., GHEZZI A., CAPUTO D., MONTANARI E., BERTOLOTTA A., BERGAMASCHI R., CAPRA R., MANCARDI G., MARTINELLI V., COMI G., FILIPPI M. In vivo assessment of the brain and cervical cord pathology of patients with primary progressive multiple sclerosis *Brain* 2001; 124(Pt 12):2540-9.
- ROVARIS M., COMI G., FILIPPI M. MRI markers of destructive pathology in multiple sclerosis-related cognitive dysfunction. *J Neurol Sci.* 2006 Jun 15; 245(1-2):111-6.
- ROVARIS M., FILIPPI M. Diffusion Tensor MRI in Multiple Sclerosis. *J Neuroimaging* 2007; 17: S27-30.
- ROVARIS M., FILIPPI M. MRI correlates of cognitive dysfunction in multiple sclerosis patients. *J Neurol Virol* 2000; 6(Suppl 2):172-75.
- ROVARIS M., FILIPPI M., FALAUTANO M., MINICUCCI L., ROCCA M. A., MARTINELLI V., *et al.*, Relation between MR abnormalities and patterns of cognitive impairment in multiple sclerosis. *Neurology* 1998; 50:1601-8.
- ROVARIS M., GASS A., BAMMER R., HICKMAN S. J., CICCARELLI O., MILLER D. H., FILIPPI M. Diffusion MRI in multiple sclerosis. *Neurology* 2005; 65(10):1526-32.
- ROVARIS M., HORSFIELD M. A., FILIPPI M. Correlations between magnetization transfer metrics and other magnetic resonance abnormalities in multiple sclerosis. *Neurology* 1999; 53(5 Suppl 3):S40-5.
- ROVARIS M., RICCITELLI G., JUDICA E., POSSA F., CAPUTO D., GHEZZI A., BERTOLOTTA A., CAPRA R., FALAUTANO M., MATTIOLI F., MARTINELLI V., COMI G., FILIPPI M. Cognitive impairment and structural brain damage in benign multiple sclerosis. *Neurology* 2008; 71(19):1521-6.
- ROVIRA A. Resonancia magnética en el diagnóstico de la esclerosis múltiple. *Radiología* 2001; 43(10): 465-76.
- ROVIRA A., ALONSO J., CUCURELLA G., NOS C., TINTORE M., PEDRAZA S., RIO J., MONTALBAN X. Evolution of multiple sclerosis lesions on serial contrast-enhanced T1-weighted and magnetization-transfer MR images. *AJNR Am J Neuroradiol.* 1999; 20(10):1939-45.
- ROVIRA A., PERICOT I., ALONSO J., RIO J., GRIVÉ E., MONTALBAN X. Serial diffusion-weighted MR imaging and proton MR spectroscopy of acute large demyelinating brain lesions: case report. *AJNR Am J Neuroradiol.* 2002; 23:989-94.
- ROVIRA A., WATTJES M. P., TINTORÉ M., TUR C., YOUSRY T. A., SORMANI M. P., DE STEFANO N., FILIPPI M., AUGER C., ROCCA M. A., BARKHOF F., FAZEKAS F., KAPPOS L., POLMAN C., MILLER D., MONTALBAN X., MAGNIMS study group. Evidence-based guidelines: MAGNIMS consen-

- sus guidelines on the use of MRI in multiple sclerosis-clinical implementation in the diagnostic process. *Nat Rev Neurol*. 2015; 11(8):471-82.
- ROY S., FRNDAK S., DRAKE A. S., IRWIN L., ZIVADINOV R., WEINSTOCK-GUTTMAN B., BENEDICT R. H. Differential effects of aging on motor and cognitive functioning in multiple sclerosis. *Mult Scler*. 2017 Sep; 23(10):1385-1393. doi: 10.1177/1352458516679036. Epub 2016 Nov 25.
- RUANO L., PORTACCIO E., GORETTI B., NICCOLAI C., SEVERO M., PATTI F., CILIA S., GALLO P., GROSSI P., GHEZZI A., ROSCIO M., MATTIOLI F., STAMPATORI C., TROJANO M. Age and disability drive cognitive impairment in multiple sclerosis across disease subtypes. *Mult Scler*. 2017 Aug; 23(9):1258-1267.
- RUCHKIN D. S., GRAFMAN J., KRAUSS G. L., JOHNSON R JR., CANOUNE H., RITTER W. Event-related brain potential evidence for a verbal working memory deficit in multiple sclerosis. *Brain* 1994; 117(Pt 2):289-305.
- RUEGER M. A., KEUTERS M. H., WALBERER M., BRAUN R., KLEIN R. SPARING R., *et al*. Multi-session transcranial direct current stimulation (tDCS) elicits inflammatory and regenerative processes in the rat brain. *Plos One* 2012; 7(8):e43776.
- RUET A., DELOIRE M., CHARRÉ-MORIN J., HAMEL D., BROCHET B. Cognitive impairment differs between primary progressive and relapsing-remitting MS. *Neurology*. 2013 Apr 16; 80(16):1501-8. doi: 10.1212/WNL.0b013e31828cf82f. Epub 2013 Mar 20.
- SACCÀ F., COSTABILE T., CAROTENUTO A., LANZILLO R., MOCCIA M., PANE C., RUSO C. V., BARBARULO A. M., CASERTANO S., ROSSI F., SIGNORIELLO E., LUS G., BRESCIA MORRA V. The EDSS integration with the Brief International Cognitive Assessment for Multiple Sclerosis and orientation tests. *Mult Scler*. 2017; 23(9):1289-1296.
- SACCO R., BONAVITA S., ESPOSITO F., TEDESCHI G., GALLO A. The Contribution of Resting State Networks to the Study of Cortical Reorganization in MS. *Mult Scler Int*. 2013; 2013:857807.
- SAMS M., PAAVILAINEN P., ALHO K., NÄÄTÄNEN R. Auditory frequency discrimination and event-related potentials. *Electroencephalogr Clin Neurophysiol*. 1985; 62(6):437-48.
- SANDROFF B. M., DLUGONSKI D., PILUTTI L. A., PULA J. H., BENEDICT R. H., MOTL R. W. Physical activity is associated with cognitive processing speed in persons with multiple sclerosis. *Mult Scler Relat Disord*. 2014a Jan; 3(1):123-8.
- SANDROFF B. M., HILLMAN C. H., BENEDICT R. H., MOTL R. W. Acute effects of walking, cycling, and yoga exercise on cognition in persons with relapsing-remitting multiple sclerosis without impaired cognitive processing speed. *J Clin Exp Neuropsychol*. 2015b; 37(2):209-19.
- SANDROFF B. M., KLAREN R. E., PILUTTI L. A., DLUGONSKI D., BENEDICT R. H., MOTL R. W. Randomized controlled trial of physical activity, cognition, and walking in multiple sclerosis. *J Neurol*. 2014b Feb; 261(2):363-72.
- SANDROFF B. M., MOTL R. W., DE LUCA J. The influence of cognitive impairment on the fitness-cognition relationship in MS. *Med Sci Sports Exerc*. 2017 Jan 23.
- SANDROFF B. M., PILUTTI L. A., BENEDICT R. H., MOTL R. W. Association between physical fitness and cognitive function in multiple sclerosis: does disability status matter? *Neurorehabil Neural Repair*. 2015a Mar-Apr; 29(3):214-23.

- SANDRY J, PAXTON J *et al.* General Mathematical Ability Predicts PASAT Performance in MS Patients: Implications for Clinical Interpretation and Cognitive Reserve. *Journal of the International Neuropsychological Society* 2016 March; 22:375-378
- SANFILIPPO M. P., BENEDICT R. H., WEINSTOCK-GUTTMAN B., BAKSHI R. Gray and white matter brain atrophy and neuropsychological impairment in multiple sclerosis. *Neurology* 2006; 66:685-92.
- SARTORI E., EDAN G. Assessment of cognitive dysfunction in multiple sclerosis. *J Neurol Sci.* 2006; 15., 245(1-2):169-75.
- SASTRE-GARRIGA J., ALONSO J., RENOM M., AREVALO M. J., GONZALEZ I., GALAN I., *et al.* A functional magnetic resonance proof of concept pilot trial of cognitive rehabilitation in multiple sclerosis. *Mult Scler.* 2011; 17(4):457-67.
- SASTRE-GARRIGA J., INGLE G. T., CHARD D. T., RAMIO-TORRENTA L., MCLEAN M. A., MILLER D. H., THOMPSON A. J. Metabolite changes in normal-appearing gray and white matter are linked with disability in early primary progressive multiple sclerosis. *Archives of Neurology* 2005; 62(4):569-73.
- SBARDELLA E., TONA F., PETSAS N., PANTANO P. DTI Measurements in Multiple Sclerosis: Evaluation of Brain Damage and Clinical Implications. *Mult Scler Int.* 2013; 2013:671730.
- SBARDELLA E., TONA F., PETSAS N., UPADHYAY N., PIATTELLA M. C., FILIPPINI N., PROSPERINI L., POZZILLI C., PANTANO P. Functional connectivity changes and their relationship with clinical disability and white matter integrity in patients with relapsing-remitting multiple sclerosis. *Mult Scler.* 2015; 21(13):1681-92.
- SCHEINBERG L., HOLLAND N., LA ROCCA N., LAITIN P., BENNETT A., HALL H. Multiple sclerosis: earning a living. *NY State J Med.* 1980; 80:1395-1400.
- SCHMIERER K., SCARAVILLI F., ALTMANN D. R., BARKER G. J., MILLER D. H. Magnetization transfer ratio and myelin in postmortem multiple sclerosis brain. *Ann Neurol.* 2004; 56(3):407-15.
- SCHOONHEIM M. M., GEURTS J. J., BARKHOF F. The limits of functional reorganization in multiple sclerosis. *Neurology* 2010; 74(16):1246-710.
- SCHOONHEIM M. M., MEIJER K. A., GEURTS J. J. Network collapse and cognitive impairment in multiple sclerosis. *Front Neurol.* 2015; 6:82.
- SCHOONHEIM M. M., POPESCU V., RUEDA LOPES F. C., WIEBENGA O. T., VRENKEN H., DOUW L., POLMAN C. H., GEURTS J. J., BARKHOF F. Subcortical atrophy and cognition: sex effects in multiple sclerosis. *Neurology* 2012; 79(17):1754-61.
- SCHULZ D., KOPP B., KUNKEL A., FAISS J. H. Cognition in the early stage of multiple sclerosis. *Journal of Neurology* 2006; 253:1002-1010.
- SCHURCH B., SCHMID D. M., STOHRER M: Treatment of neurogenic incontinence with botulinum toxin A. *N Engl J Med* 2000; 342:665.
- SCHWARTZMANN L. Calidad de vida relacionada con la salud: aspectos conceptuales. *Ciencia y enfermería* 2003; 9:9-21.
- SCHWID S. R., GOODMAN A. D., WEINSTEIN A., MCDERMOTT M. P., JOHNSON K. P. Cognitive function in relapsing multiple sclerosis: minimal changes in a 10-year clinical trial. *J Neurol Sci.* 2007; 255:57-63.

- SEDIGHI B., PARDAKHTY A., KAMALI H., SHAFIEE K., HASANI B. N. Effect of *Boswellia papyrifera* on cognitive impairment in multiple sclerosis. *Iran J Neurol*. 2014 Jul 4; 13(3):149-53.
- SEPULCRE J., VANOTTI S., HERNÁNDEZ R., SANDOVAL G., CÁCERES F., *et al*. Cognitive impairment in patients with multiple sclerosis using the Brief Repeatable Battery Neuropsychology test. *Mult Scler* 2006; 12(2):187-195.
- SFAGOS C., PAPAGEORGIOU C. C., KOSMA K. K., KODOPADELIS E., UZUNOGLU N. K., VASSILOPOULOS D., RABAVILAS A. D. Working memory deficits in multiple sclerosis: a controlled study with auditory P600 correlates. *J Neurol Neurosurg Psychiatry* 2003; 74(9):1231-5.
- SHATIL E., METZER A., HORVITZ O., MILLER A. Home-based personalized cognitive training in MS patients: a study of adherence and cognitive performance. *NeuroRehabilitation* 2010; 26(2):143-53.
- SIMEONI M. C., AUQUIER P., FERNANDEZ O., *et al*. Validation of the multiple sclerosis international quality of life questionnaire. *Multiple Sclerosis* 2008; 14(2):219-230.
- SIMMONS R. D. Life issues in multiple sclerosis. *Nature Reviews Neurology* 2010; 6(11):603-610. doi:10.1038/nrneuro.2010.143
- SIMON J. H., SCHERZINGER A., RAFF U., *et al*. Computerized method of lesion volume quantitation in multiple sclerosis: error of serial studies. *AJNR Am J Neuroradiol*. 1997; 18:580-82.
- SIMON J. H., ZHANG S., LAIDLAW D. H., MILLER D. E., BROWN M., CORBOY J., BENNETT J. Identification of fibers at risk for degeneration by diffusion tractography in patients at high risk for MS after a clinically isolated syndrome. *J Magn Reson Imaging*. 2006; 24(5):983-8.
- SIMON J. H. Brain atrophy in multiple sclerosis: what we know and would like to know. *Brain* 2002; 125:1676-95.
- SMERBECK A., BENEDICT R. H., ESHAGHI A., VANOTTI S., SPEDO C., BLAHOVA DUSANKOVA J., SAHRAIÁN M. A., MARQUES V. D., LANGDON D. Influence of nationality on the Brief International Cognitive Assessment for Multiple Sclerosis (BICAMS). *Clin Neuropsychol*. 2017 Jul 19: 1-9.
- SOLARI A., MOTTA A., MENDOZZI L., PUCCI E., FORNI M., MANCARDI G., POZZILLI C; CRIMS Trial. Computer-aided retraining of memory and attention in people with multiple sclerosis: a randomized, double-blind controlled trial. *J Neurol Sci*. 2004 Jul 15; 222(1-2):99-104.
- SOLLARI A., MOTTA A. A shortened version of PASAT-3 is feasible. *Mult Scler*. 2007 Sep; 13(8):1020-5. Epub 2007 Jul 10.
- SONG S. K., YOSHINO J., LE T. Q., LIN S. J., SUN S. W., CROSS A. H., ARMSTRONG R. C. Demyelination increases radial diffusivity in corpus callosum of mouse brain. *Neuroimage* 2005; 26(1):132-40.
- SPERLING R. A., DICKERSON B. C., PIHLAJAMAKI M., *et al*., Functional alterations in memory networks in early Alzheimer's disease. *Neuromolecular Med* 2010; 12:27-43.
- SPERLING R. A., GUTTMANN C. R., HOHOL M. J., WARFIELD S. K., JAKAB M., PARENTE M *et al*. Regional magnetic resonance imaging lesion burden and cognitive function in multiple sclerosis: a longitudinal study. *Arch Neurol* 2001; 58:115-21.
- SRINIVASAN R., SAILASUTA N., HURD R., NELSON S., PELLETIER D. Evidence of elevated glutamate in multiple sclerosis using magnetic resonance spectroscopy at 3 T. *Brain* 2005; 128(5):1016-25.
- STAFFEN W., MAIR A., ZAUNER H., UNTERRAINER J., NIEDERHOFER H., KUTZELNIGG A., RITTER S., GOLASZEWSKI S., IGLSEDER B., LADURNER G. Cognitive function and fMRI in patients with

- multiple sclerosis: evidence for compensatory cortical activation during an attention task. *Brain* 2002; 125 (Pt 6):1275-82.
- STANKOFF B., WAUBANT E., CONFAVREUX C., EDAN G., DEBOUVERIE M., RUMBACH L., MOREAU T., PELLETIER J., LUBETZKI C., CLANET M; French Modafinil Study Group. Modafinil for fatigue in MS: a randomized placebo-controlled double-blind study. *Neurology*. 2005 Apr 12; 64(7):1139-43.
- STEFAN K., KUNESCH E., COHEN L. G., BENECKE R., CLASSEN J. Induction of plasticity in the human motor cortex by paired associative stimulation. *Brain* 2000 Mar; 123 Pt 3:572-84.
- STEFAN K., KUNESCH E., BENECKE R., COHEN L. G., CLASSEN J. Mechanisms of enhancement of human motor cortex excitability induced by interventional paired associative stimulation. *J Physiol*. 2002 Sep 1; 543(Pt 2):699-708.
- STENAGER E., STENAGER E. N., JENSEN K. Función sexual en la esclerosis múltiple: un estudio de seguimiento de 5 años. *Ital J Neurol Sci*. 1996; 17:67-9.
- STEVENSON V. L., GAWNE-CAIN M. L., BARKER G. J., THOMPSON A. J., MILLER D. H. Imaging of the spinal cord and brain in multiple sclerosis: a comparative study between fast FLAIR and fast spin echo. *J Neurol* 1997; 244(2):119-24.
- STROBER L., ENGLERT J., MUNSCHAUER F., WEINSTOCK-GUTTMAN B., RAO S., BENEDICT R. H. Sensitivity of conventional memory tests in multiple sclerosis: comparing the Rao Brief Repeatable Neuropsychological Battery and the Minimal Assessment of Cognitive Function in MS. *Mult Scler*. 2009 Sep; 15(9):1077-84. doi: 10.1177/1352458509106615. Epub 2009 Jun 25.
- STROBER L. B., RAO S. M., LEE J. C., FISCHER E., RUDICK R. Cognitive impairment in multiple sclerosis: An 18 year follow-up study. *Mult Scler Relat Disord*. 2014 Jul; 3(4):473-81.
- STUIFBERGEN A. K., BECKER H., PEREZ F., MORISON J., KULLBERG V., TODD A. A randomized controlled trial of a cognitive rehabilitation intervention for persons with multiple sclerosis. *Clin Rehabil*. 2012;26(10):882-93.
- SUMMERS M., FISNIKU L., ANDERSON V., MILLER D., CIPOLOTTI L., RON M. Cognitive impairment in relapsing-remitting multiple sclerosis can be predicted by imaging performed several years earlier. *Mult Scler*. 2008; 14(2):197-204.
- SUMOWSKI J., BENEDICT R., ENZINGER C., FILIPPI M *et al*. Cognition in multiple sclerosis. State of the field and priorities for the future. *Neurology*® 2018; 90:1-11.
- SUMOWSKI J. F., WYLIE G. R., DELUCA J., CHIARAVALLOTI N. Intellectual enrichment is linked to cerebral efficiency in multiple sclerosis: functional magnetic resonance imaging evidence for cognitive reserve. *Brain* 2010; 133(Pt 2):362-74.
- SUMOWSKI J. F., CHIARAVALLOTI N., LEAVITT V. M., DE LUCA J. Cognitive reserve in secondary progressive multiple sclerosis. *Mult Scler*. 2012 Oct; 18(10):1454-8.
- SUMOWSKI J. F., ROCCA M. A., LEAVITT V. M., RICCITELLI G., COMI G., DE LUCA J., FILIPPI M. Brain reserve and cognitive reserve in multiple sclerosis: what you've got and how you use it. *Neurology*. 2013 Jun 11; 80(24):2186-93.
- SUMOWSKI J. F., WYLIE G. R., CHIARAVALLOTI N., DE LUCA J. Intellectual enrichment lessens the effect of brain atrophy on learning and memory in multiple sclerosis. *Neurology* 2010a Jun 15; 74(24):1942-5.

- SUMOWSKI J. F., WYLIE G. R., GONNELLA A., CHIARAVALLOTI N., DE LUCA J. Premorbid cognitive leisure independently contributes to cognitive reserve in multiple sclerosis. *Neurology* 2010b Oct 19; 75(16):1428-31.
- SUMOWSKI J. F. Cognitive reserve as a useful concept for early intervention research in multiple sclerosis. *Front Neurol.* 2015; 6:176.
- SUNDGREN M., PIEHL F., WAHLIN Å., BRISMAR T. Cognitive function did not improve after initiation of natalizumab treatment in relapsing-remitting multiple sclerosis. A prospective one-year dual control group study. *Mult Scler Relat Disord.* 2016 Nov; 10:36-43.
- SWEET L. H., RAO S. M., PRIMEAU M., MAYER A. R., COHEN R. A. Functional magnetic resonance imaging of working memory among multiple sclerosis patients. *J Neuroimaging* 2004; 14:150-57.
- SWIRSKY-SACCHETTI T., MITCHELL D. R., SEWARD J., GONZALES C., LUBLIN F., KNOBLER R *et al.* Neuropsychological and structural brain lesions in multiple sclerosis: a regional analysis. *Neurology* 1992; 42:1291-95.
- TAGLIAZUCCHI E., LAUFS H. Multimodal Imaging of Dynamic Functional Connectivity. *Front Neurol* 2015; 16, 6:10.
- TARIK CATIC, JOSIP CULIG, ENRA SULJIC, ADMIR MASIC Y REFET GOJAK .Validación del cuestionario específico para la enfermedad MSQoL-54 en Bosnia y Herzegovina. *Med Arch.* 2017 Apr; 71(2):103-106. doi: 10.5455 / medaR. H.2017.71.103-106.
- TECCHIO F., ZITO G., ZAPPASODI F, *et al.* Intra-cortical connectivity in multiple sclerosis: a neurophysiological approach. *Brain* 2008;131:1783-1792.
- TECCHIO F., CANCELLI A., COTTONE C., ZITO G., PASQUALETTI P., GHAZARYAN A., ROSSINI P. M., FILIPPI M. M. Multiple sclerosis fatigue relief by bilateral somatosensory cortex neuromodulation. *J Neurol.* 2014 Aug; 261(8):1552-8.
- TEDESCHI G., DINACCI D., COMERCI M., LAVORGNA L., SAVETTIERI G., QUATTRONE A., LIVREA P., PATTI F., MORRA V. B., SERVILLO G., OREFICE G., PACIELLO M., PRINSTER A., CONIGLIO G., BONAVITA S., DI COSTANZO A., BELLACOSA A., VALENTINO P., QUARANTELLI M., BRUNETTI A., SALEMI G., D'AMELIO M., SIMONE I., SALVATORE M., BONAVITA V., ALFANO B. Brain atrophy evolution and lesion load accrual in multiple sclerosis: a 2-year follow-up study. *Mult Scler.* 2009; 15(2):204-11.
- TEKOK-KILIC A., BENEDICT R. H., WEINSTOCK-GUTTMAN B., DWYER M. G., CARONE D., SRINIVASARAGHAVAN B., YELLA V., ABDELRAHMAN N., MUNSCHAUER F., BAKSHI R., ZIVADINOV R. Independent contributions of cortical gray matter atrophy and ventricle enlargement for predicting neuropsychological impairment in multiple sclerosis. *Neuroimage* 2007; 36(4):1294-300.
- TERVANIEMI M., LEHTOKOSKI A., SINKKONEN J., VIRTANEN J., ILMONIEMI R. J., NÄÄTÄNEN R. Test-retest reliability of mismatch negativity for duration., frequency and intensity changes. *Clin Neurophysiol.* 1999; 110(8):1388-93.
- TESAR N., BANDION K., BAUMHACKL U. Efficacy of a neuropsychological training programme for patients with multiple sclerosis - a randomised controlled trial. *Wien Klin Wochenschr.* 2005; 117:747-54.

- THE CANADIAN BURDEN OF ILLNESS STUDY GROUP. Burden of illness of multiple sclerosis: Part II: Quality of life. *Can J Neurol Sci.* 1998 Feb; 25(1):31-8.
- THELEN J. M., LYNCH S. G., BRUCE A. S., HANCOCK L. M., BRUCE J. M. Polypharmacy in multiple sclerosis: relationship with fatigue, perceived cognition, and objective cognitive performance. *J Psychosom Res.* 2014 May; 76(5):400-4.
- THOMPSON D., YU C. Quality of life in patients with coronary heart disease. *Health Qual Life Outcomes* 2003; 1(1). doi:10.1186/1477-7525-1-42
- THORNTON A., RAZ N., TUCKE K. Memory in multiple sclerosis: contextual encoding deficits. *J Internat Neuropsychol Soc.* 2002; 8:395-409.
- THORPE J. W., KIDD D., MOSELEY I. F., *et al.* Serial gadolinium enhanced MRI of the brain and spinal cord in early relapsing-remitting multiple sclerosis. *Neurology* 1996; 46:373-78.
- TIEMANN L., PENNER I. K., HAUPTS M., SCHLEGEL U., CALABRESE P. Cognitive decline in multiple sclerosis: impact of topographic lesion distribution on differential cognitive deficit patterns. *Mult Scler* 2009; 15:1164-1174.
- TILLEMA J. M., PIRKO I. Neuroradiological evaluation of demyelinating disease. *Ther Adv Neurol Disord.* 2013; 6(4):249-68.
- TOMASEVIC L., ZITO G., PASQUALETTI P., FILIPPI M., LANDI D., GHAZARYAN A., LUPOI D., PORCARO C., BAGNATO F., ROSSINI P., TECCHIO F. Cortico-muscular coherence as an index of fatigue in multiple sclerosis. *Mult Scler.* 2013; 19:334-343.
- TOMASSINI V., MATTHEWS P. M., THOMPSON A. J., FUGLØ D., GEURTS J. J., JOHANSEN-BERG H., JONES D. K., ROCCA M. A., WISE R. G., BARKHOF F., PALACE J. Neuroplasticity and functional recovery in multiple sclerosis. *Nat Rev Neurol.* 2012; 8(11):635-46.
- TOMBAUGH T. N. A comprehensive review of the Paced Auditory Serial Addition Test (PASAT). *Arch Clin Neuropsychol Off J Natl Acad Neuropsychol.* 2006 Enero; 21(1):53-76.
- TOOSY A. T., HICKMAN S. J., MISZKIEL K. A., JONES S. J., PLANT G. T., ALTMANN D. R., BARKER G. J., MILLER D. H., THOMPSON A. J. Adaptive cortical plasticity in higher visual areas after acute optic neuritis. *Ann Neurol.* 2005; 57(5):622-33.
- TORTORELLA P., ROCCA M. A., MEZZAPESA D. M., GHEZZI A., LAMANTIA L., COMI G., FILIPPI M. MRI quantification of gray and white matter damage in patients with early-onset multiple sclerosis. *J Neurol.* 2006; 253(7):903-7.
- TOVAR-MOLL F., EVANGELOU I. E., CHIU A. W., RICHERT N. D., OSTUNI J. L., OHAYON J. M., AUH S., EHRMANTRAUT M., TALAGALA S. L., MCFARLAND H. F., BAGNATO F. Thalamic involvement and its impact on clinical disability in patients with multiple sclerosis: A diffusion tensor imaging study at 3T. *AJNR Am J Neuroradiol.* 2009; 30(7):1380-6.
- TRACY V. L., BASSO M. R., MARSON D. C., COMBS D. R., WHITESIDE D. M. Capacity for financial decision making in multiple sclerosis. *J Clin Exp Neuropsychol.* 2017 Feb; 39(1):46-57. Epub 2016 Jul 19.
- TRIANAFYLLOU N. I., VOUMVOURAKIS K., ZALONIS I., SFAGOS K., MANTOUVALOS V., MALLIARA S., PAPAGEORGIOU C. Cognition in relapsing-remitting multiple sclerosis: a multichannel event-related potential (P300) study. *Acta Neurol Scand.* 1992; 85(1):10-3.
- TRUYEN L., VAN WAESBERGHE J. H., VAN WALDERVEEN M. A., VAN OOSTEN B. W., POLMAN C. H., HOMMES O. R., ADÈR H. J., BARKHOF F. Accumulation of hypointense lesions (“black holes”)

- on T1 spin-echo MRI correlates with disease progression in multiple sclerosis. *Neurology* 1996; 47(6):1469-76.
- TWORK S., WIESMETH S., SPINDLER M., WIRTZ M., SCHIPPER S., PÖHLAU D., KLEWER J., KUGLER J. Disability status and quality of life in multiple sclerosis: non-linearity of the Expanded Disability Status Scale (EDSS). *Health Qual Life Outcomes*. 2010 Jun 7;8:55. doi: 10.1186/1477-7525-8-55.
- UYSAI U., IDIMAN F., IDIMAN E., OZAKBAS S., KARAKAS S., BRUCE J. Contingent negative variation is associated with cognitive dysfunction and secondary progressive disease course in multiple sclerosis. *J Clin Neurol*. 2014c; 10(4):296-303.
- VAN DER HIELE K., VAN GORP D. A., HEERINGS M. A., VAN LIESHOUT I., JONGEN P. J., RENEMAN M. F., VAN DER KLINK J. J.; MS@Work Study Group. The MS @Work Study: a 3-year prospective observational study on factors involved with work participation in patients with relapsing-remitting Multiple Sclerosis. *BMC Neurol*. 2015 Aug 12; 15:134.
- VAN DER HIELE K., VAN GORP D., RUIMSCHOTEL R., KAMMINGA N., VISSER L., MIDDELKOOP H. Work Participation and Executive Abilities in Patients with Relapsing-Remitting Multiple Sclerosis. *PLoS One*. 2015 Jun 17; 10(6):e0129228. doi: 10.1371/journal.pone.0129228. eCollection 2015.
- VAN WALDERVEEN M. A., BARKHOF F, TAS M. W., POLMAN C., FREQUIN S. T., HOMMES O. R., THOMPSON A. J., VALK J. Patterns of brain magnetic resonance abnormalities on T2-weighted spin echo images in clinical subgroups of multiple sclerosis: a large cross-sectional study. *Eur Neurol* 1998a; 40(2):91-8.
- VAN WALDERVEEN M. A., KAMPHORST W., SCHELTENS P., VAN WAESBERGHE J. H., RAVID R., VALK J., POLMAN C. H., BARKHOF F. Histopathologic correlate of hypointense lesions on T1-weighted spin-echo MRI in multiple sclerosis. *Neurology* 1998b; 50(5):1282-88.
- VANOTTI S., BENEDICT R. H., ACION L., CÁCERES F; VANEM Workgroup. Validation of the Multiple Sclerosis Neuropsychological Screening Questionnaire in Argentina. *Mult Scler*. 2009 Feb; 15(2):244-50. doi:10.1177/1352458508097924. Epub 2008
- VÁZQUEZ-MARRUFO M., GALVAO-CARMONA A., GONZÁLEZ-ROSA J. J., HIDALGO-MUÑOZ A. R., BORGES M., RUIZ-PEÑA J. L., IZQUIERDO G. Neural correlates of alerting and orienting impairment in multiple sclerosis patients. *PLoS One* 2014; 9(5):e97226.
- VÁZQUEZ-MARRUFO M., GONZÁLEZ-ROSA J., VAQUERO-CASARES E., DUQUE P., BORGES M., IZQUIERDO G. Cognitive evoked potentials in remitting-relapsing and benign forms of multiple sclerosis. *Rev Neurol*. 2009; 48(9):453-8.
- VÁZQUEZ-MARRUFO M., GONZÁLEZ-ROSA J. J., GALVAO-CARMONA A., HIDALGO-MUÑOZ A., BORGES M., PEÑA J. L., IZQUIERDO G. Retest reliability of individual p3 topography assessed by high density EEG. *PLoS One* 2013; 8(5):e62523.
- VÁZQUEZ-MARRUFO M., GONZÁLEZ-ROSA J. J., VAQUERO E., DUQUE P., ESCERA C., BORGES M., IZQUIERDO G., GÓMEZ C. M. Abnormal ERPs and high frequency bands power in multiple sclerosis. *Int J Neurosci*. 2008; 118(1):27-38.
- VELIKONJA O., CURIĆ K., OZURA A., JAZBEC S. S. Influence of sports climbing and yoga on spasticity, cognitive function, mood and fatigue in patients with multiple sclerosis. *Clin Neurol Neurosurg*. 2010 Sep; 112(7):597-601.

- VERLEGER R. Event-related EEG potential research in neurological patients. Chapter 13., *The Cognitive Electrophysiology of Mind and Brain.*, 2003; 309-41.
- VICENS A., IRIARTE J., DE CASTRO P., CASTEJÓN C., MARTÍNEZ-LAGE J. M. Funciones cognitivas en la esclerosis múltiple. *Neurología* 1992; 7:171-175.
- VIN REY F., HEESAKKERS J. Solifenacin in multiple sclerosis patients with overactive bladder: a prospective study. *Adv Urol.* 2011; 2011:834753. doi: 10.1155/2011/834753. Epub 2011 May 5.
- VIRTA J. R., LAATU S., PARKKOLA R., OIKONEN V., RINNE J. O., RUUTIAINEN J. Cerebral acetylcholinesterase activity is not decreased in MS patients with cognitive impairment. *Mult Scler.* 2011 Aug; 17(8):931-8.
- VITKOVITCH M., BISHOP S., DANCEY., RICHARDS A. Stroop interference and negative priming in patients with multiple sclerosis. *Neuropsychologia* 2002; 40(9):1570-1576.
- WACHOWIUS U., TALLEY M., SILVER N., HEINZE H. J., SAILER M. Cognitive impairment in primary and secondary progressive sclerosis. *J Clin Exp Neuropsychol* 2005; 27:65-77.
- WALKER L. A., BERARD J. A., ATKINS H. L., BOWMAN M., LEE H., FREEDMAN M. S. Cognitive change and neuroimaging following immunoablative therapy and hematopoietic stem cell transplantation in multiple sclerosis: A pilot study. *Mult Scler Relat Disord.* 2014 Jan; 3(1):129-35.
- WASSMER E., CHITNIS T., POHL D., AMATO M. P., BANWELL B., GHEZZI A., HINTZEN R. Q., KRUPP L. B., MAKHANI N., ROSTÁSY K., TARDIEU M., TENEMBAUM S., WALDMAN A., WAUBANT E., KORNBORG A. J. Neuropsychological features in childhood and juvenile multiple sclerosis: five-year follow-up. International Pediatric MS Study Group Global Members Symposium report. *Neurology* 2016 Aug 30; 87(9 Suppl 2):S110-6.
- WEINSTEIN A., SCWID S. I., SCHIFFER R. B., McDERMOTT M. P., GIANG. D. W., GOODMAN AD (1999) Neuropsychologic status in multiple sclerosis after treatment with glatiramer. *Arch Neurol.* 56:319-324.
- WERRING D. J., CLARK C. A., BARKER G. J., THOMPSON A. J., MILLER D. H. Diffusion tensor imaging of lesions and normal-appearing white matter in multiple sclerosis. *Neurology* 1999; 52(8):1626-32.
- WHELAN R., LONERGAN R., KIISKI H., NOLAN H., KINSELLA K., BRAMHAM J., O'BRIEN M., REILLY R. B., HUTCHINSON M., TUBRIDY N. A high-density ERP study reveals latency, amplitude, and topographical differences in multiple sclerosis patients versus controls. *Clin Neurophysiol* 2010; 121(9):1420-6.
- WICKS C. R., WARD K., STROUD A., TENNANT A., Ford HL Multiple sclerosis and employment: Associations of psychological factors and work instability. *J Rehabil Med.* 2016 Oct 12; 48(9):799-805. doi: 10.2340/16501977-2142.
- WILKEN J., KANE R. L., SULLIVAN C. L., GUEDEBLATT M., LUCAS S., FALLIS R., YOU X., FOULDS P. Changes in Fatigue and Cognition in Patients with Relapsing Forms of Multiple Sclerosis Treated with Natalizumab: The ENER-G Study. *Int J MS Care.* 2013 Fall; 15(3):120-8.
- WISHART H. A., SAYKIN A. J., McDONALD B. C., MAMOURIAN A. C., FLASHMAN L. A., SCHUSCHU K. R., RYAN K. A., FADUL C. E., KASPER L. H. Brain activation patterns associated with working memory in relapsing remitting MS. *Neurology* 2004; 62(2):234-38.
- WOLFF S. D., BALABAN R. S. Magnetization transfer imaging: practical aspects and clinical applications. *Radiology* 1994; 192:593-599.

- WOODMAN G. F. A Brief Introduction to the Use of Event-Related Potentials (ERPs) in Studies of Perception and Attention. *Atten Percept Psychophys.* 2010; 72(8):2031-46.
- WU N., MINDEN S. L., HOAGLIN D. C., HADDEN L., FRANKEL D. Quality of life in people with multiple sclerosis: Data from the Sonya Slifka Longitudinal Multiple Sclerosis Study. *J Health Hum Serv Adm.* 2007; 30:233-67.
- YORKE A., COHEM E. Functional Assessment of Multiple Sclerosis. *Journal of Physiotherapy* 2015; 61(4):226.
- YU H. J., CHRISTODOULOU C., BHISE V., GREENBLATT D., PATEL Y., SERAFIN D., MALETIC-SAVATIC M., KRUPP L. B., WAGSHUL M. E. Multiple white matter tract abnormalities underlie cognitive impairment in RRMS. *Neuroimage* 2012; 59(4):3713-22.
- ZAKZANIS K. K. Distinct neuro-cognitive profiles in multiple sclerosis subtypes. *Arch Clin Neuropsychol.* 2000; 15:115-36.
- ZANI A. Evoked and Event-Related Potentials. En: RUNEHOV ALC., OVIEDO L., eds. *Encyclopedia of Sciences and Religions.* Dordrecht: Springer, 2013.
- ZHONGHUA YI XUE ZA ZHI (2016). Information processing speed and influential factors in multiple sclerosis; 96 (15): 1173-7. DOI: 10.3760 / CMA.J.ISSN.0376-2491.2016.15.006. CHINESE
- ZHOU F., ZHUANG Y., GONG H., WANG B., WANG X., CHEN Q., WU L., WAN H. Altered inter-sub-region connectivity of the default mode network in relapsing remitting multiple sclerosis: a functional and structural connectivity study. *PLoS One* 2014; 9(7):e101198.
- ZIVADINOV R., BAKSHI R. Role of MRI in multiple sclerosis II: brain and spinal cord atrophy. *Front Biosci* 2004; 9:647-64.
- ZIVADINOV R., COX J. L. Neuroimaging in multiple sclerosis. *Int Rev Neurobiol.* 2007; 79:449-74.
- ZIVADINOV R., SEPCIC J., NASUELLI D., DE MASI R., BRAGADIN L. M., TOMMASI M. A., ZAMBITO-MARSALA S., MORETTI R., BRATINA A., UKMAR M., POZZI-MUCELLI R. S., GROF A., CAZZATO G., ZORZON M. A longitudinal study of brain atrophy and cognitive disturbances in the early phase of relapsing-remitting multiple sclerosis. *J Neurol Neurosurg Psychiatry.* 2001; 70:773-80.
- ZIVADINOV R., STOSIC M., COX J. L., RAMASAMY D. P., DWYER M. G. The place of conventional MRI and newly emerging MRI techniques in monitoring different aspects of treatment outcome. *J Neurol.* 2008; 255 (Suppl 1):S61-74.
- ZORZON M., ZIVADINOV R., BOSCO A., BRAGADIN LM., MORETTI R., BONFIGLI L *et al.*, Sexual dysfunction in multiple sclerosis: a case-control study. I. Frequency and comparison of groups. *Mult Scler.* 1999; 5:418-27.
- ZOU L., WANG H., XIAO Z., FANG Q., ZHANG M., LI T., DU G., LIU Y. Tai chi for health benefits in patients with multiple sclerosis: A systematic review. *PLoS One.* 2017 Feb 9; 12(2):e0170212.