

# Bibliografía web

- Aarsland, D., Marsh, L., y Schrag, A. (2009). Neuropsychiatric symptoms in Parkinson's disease. *Movement Disorders*, 24(15), 2175-2186.
- Adachi, T., Koeda, T., Hirabayashi, S., Maeoka, Y., Shiota, M., Wright, E.C., et al. (2004). The metaphor and sarcasm scenario test: a new instrument to help differentiate high functioning pervasive developmental disorder from attention deficit / hyperactivity disorder. *Brain and Development*, 26, 301-306.
- Adolphs, R. (2003). Cognitive neuroscience of human social behaviour. *Nature Reviews Neuroscience*, 4, 165-178.
- (2009). The social brain: neural basis of social knowledge. *Annual Review of Psychology*, 60, 693.
- (2010). Conceptual challenges and directions for social neuroscience. *Neuron*, 65(6), 752-67.
- Aguado, C., Martínez, J., y Onís, M. (2000). Adaptación y validación al castellano de la versión abreviada de la "Geriatric Depression Scale" (GDS) de Yesavage. *Atención Primaria*, 26(1), 328.
- Agüera-Ortiz, L., Gil-Ruiz, N., Cruz-Orduña, I., Ramos-García, I., Osorio, R.S., Valentí-Soler, M., Olazarán-Rodríguez, J., Dobato-Ayuso, J.L., Lancôt, K., y Martínez-Martín, P. (2015). A novel rating scale for the measurement of apathy in institutionalized persons with dementia: the APADEM-NH. *American Journal of Geriatric Psychiatry*, 23(2), 149-159.
- Agüera-Ortiz, L.F., Gil-Ruiz, N., Cruz-Orduña, I., Ramos-García, M.I., Osorio-Suárez, R.M., Valentí-Soler, M., Martínez-Martín, P., y grupo UIPA-CAFRS (2010). Creación de una escala de medición de la apatía en pacientes con

- demenia tipo alzhéimer institucionalizados: la escala APADEM-NH-66. *Psicogeriatría*, 2(4), 207-219.
- Aguilar, E., Hidalgo, M., Cano, R., López, J., Campillo, M., y Hernández, J. (1995). Estudio prospectivo de la desesperanza en pacientes psicóticos de inicio: características psicométricas de la escala de desesperanza de Beck en este grupo. *Anales de Psiquiatría*, 11(4), 121-125.
- Aguilar, M.J., Urquijo, S., Zabala, M.L., y López, M. (2014). Aportes empíricos a la validación y adaptación al español de las Historias extrañas de Happé. *Revista Argentina de Ciencias del Comportamiento*, 6(2), 1-10.
- Al-Adawi, S., Dorvlo, A.S., Burke, D.T., Huynh, C.C., Jacob, L., Knight, R., et al. (2004). Apathy and depression in cross-cultural survivors of traumatic brain injury. *The Journal of Neuropsychiatry & Clinical Neurosciences*, 16(4), 435-442.
- Albert, S.M., Del Castillo-Castaneda, C., Sano, M., Jacobs, D.M., Marder, K., Bell, K., et al. (1996). Quality of life in patients with Alzheimer's disease as reported by patient proxies. *Journal of the American Geriatrics Society*, 44, 1342-1347.
- Alderman, N., Dawson, K., Rutterford, N.A., y Reynolds, P.J. (2001). A comparison of the validity of self-report measures amongst people with acquired brain injury: a preliminary study of the usefulness of EuroQol-5D. *Neuropsychological Rehabilitation*, 11, 529-37.
- Alexopoulos, G.S., Abrams, R.C., Young, R.C., Shamoian, C.A. (1988). Cornell Scale for Depression in Dementia. *Biological Psychiatry*, 23(3), 271-84.
- Allen, L., Conder, R.L., Green, P., y Cox, D.R. (1997). *CARB '97 manual for the computerized assessment of response bias*. Durham, NC: CogShell.
- Alonso-Fernández, F. (2011). *CET-DE: Cuestionario estructural tetradimensional para la depresión* (6.ª ed.). Madrid: TEA Ediciones.
- Alonso, J., Prieto, L., y Antó, J.M. (1995). La versión española del SF-36 Health Survey (Cuestionario de Salud SF-36): un instrumento para la medida de los resultados clínicos. *Medicina Clínica*, 104, 771-776.
- Altman, E.G., Hedeker, D., Peterson, J.L., y Davis, J.M. (1997). The Altman self-rating mania Scale. *Biological Psychiatry*, 42, 948-955.
- Altman, E.G., Hedeker, D.R., Janicak, P.G., Peterson, J.L., y Davis, J.M. (1994). The Clinician Administered Rating Scale for Mania (CARS-M): development, reliability, and validity. *Biological Psychiatry*, 36, 124-134.
- Álvarez, P., Benavent, P., García Valls, J., Livianos, L., y Rojo, L. (2005). Traducción y adaptación de la Altman Self-Rating Mania Scale al ámbito español. *Actas Españolas de Psiquiatría*, 33, 180-187.
- Anderson, S., Bechara, A., Damasio, H., Tranel, D., y Damasio, A. (1999). Impairment of social and moral behaviour related to early damage in human prefrontal cortex. *Nature Neuroscience*, 2, 1032-1037.

- Anderson, S., Bechara, A., Damasio, H., Tranel, D., y Damasio, A. (1999). Impairment of social and moral behaviour related to early damage in human prefrontal cortex. *Nature Neuroscience*, 2, 1032-1037.
- Andreasen, N.C. (1982). Negative symptoms in schizophrenia. Definition and reliability. *Archives of General Psychiatry*, 39, 784-788.
- Angelelli, P., Paolucci, S., Bivona, U., Piccardi, L., Ciurli, P., Cantagallo, A., et al. (2004). Development of neuropsychiatric symptoms in poststroke patients: a cross-sectional study. *Acta Psychiatrica Scandinava*, 110, 55-63.
- Aparicio-López, C., Rodríguez-Rajo, P., Sánchez-Carrión, R., Enseñat, A., y García-Molina, A. (2019). Rehabilitación de la anosognosia en pacientes con heminegligencia visoespacial. *Revista de Neurología*, 69(5), 190-198.
- Argyle, M., y Cook, M. (1976). *Gaze and Mutual Gaze*. Cambridge: University Press.
- Arnould, A., Rochat, L., Azouvi, P., y Van Der Linden, M. (2013). A multidimensional approach to apathy after traumatic brain injury. *Neuropsychology Review*, 23(3), 210-233.
- Arnsten, A.F.T. (2009). Stress signalling pathways that impair prefrontal cortex structure and function. *Nature Reviews Neuroscience*, 10, 410-422.
- (2009). Stress signalling pathways that impair prefrontal cortex structure and function. *Nature Reviews Neuroscience*, 10, 410-422.
- Ashley, M.J., y Krych, D.K. (1998). *Working with behavior disorders: strategies for traumatic brain injury rehabilitation*. San Antonio, TX: Communication Skill Builders-The Psychological Corporation.
- Asociación Americana de Psicología (2014). *Manual diagnóstico y estadístico de los trastornos mentales (DSM 5)*. Barcelona: Masson.
- Babikian, T., y Boone, K. B. (2007). Intelligence Tests as Measures of Effort (pp. 103-127). En K.B. Boone (ed.), *Assessment of Feigned Cognitive Impairment: A Neuropsychological Perspective*. Nueva York: The Guilford Press.
- Babinski, M. (1914). Contribution a l'étude des troubles mentaux dans l'hémiplégie organique cérébrale (anosognosie). *Revue Neurologique*, 12, 845-848.
- Backhaus, S.L., Fichtenberg, N.L., y Hanks, R.A. (2004). Detection of sub-optimal performance using a floor effect strategy in patients with traumatic brain injury. *The Clinical Neuropsychologist*, 18, 591-603.
- Badía, X., Roset, M., Montserrat, S., Herdman, M., y Segura, A. (1999). The Spanish version of EuroQol: a description and its applications. European Quality of Life scale. *Medicina Clínica (Barc)*, 112, 79-85.
- Badía, X., y Alonso, J. (1994). Adaptación de una medida de la Disfunción Relacionada con la Enfermedad: La versión española del Sickness Impact Profile. *Medicina Clínica*, 102, 90-95.
- Bados, A. (2000). *Agorafobia y ataques de pánico*. Madrid: Pirámide.
- Barco, P.P., Crosson, B., Bolesta, M.M., Werts, D., y Stout, R. (1991). Training awareness and compensation in postacute head injury rehabilitation. En J.S.

- Kreutzer y P.H. Wehman (eds.), *Cognitive rehabilitation for persons with traumatic brain injury: A functional approach* (pp. 129-146). Baltimore, MD: Paul H. Brookes.
- Baron-Cohen, S., Tager-Flusberg, H., y Cohen, D. (2000). *Understanding other minds: perspectives from developmental cognitive neuroscience*. Oxford: University Press.
- Baron-Cohen, S. (1997). Hey! It was just a joke! Understanding propositions and propositional attitudes by normally developing children and children with autism. *Israel Journal Psychiatry and Related Sciences*, 34, 174-178.
- Baron-Cohen, S., Leslie, A., y Frith, U. (1985). Does the autistic child have a “theory of mind”? *Cognition*, 21, 37-46.
- Baron-Cohen, S., Wheelwright, S., Hill, J., Raste, Y., y Plumb, I. (2001). The “Reading the Mind in the Eyes” Test revised version: a study with normal adults, and adults with Asperger’s syndrome or high-functioning autism. *Journal of Child Psychology and Psychiatry*, 42, 241-251.
- Barrash, J., Anderson, S.W., Hathaway-Nepple, J., Jones, R.D., y Tranel, D. (1997). *The Iowa scales of personality change*. Iowa City: University of Iowa College of Medicine, Dept. of Neurology.
- Bauer, M.S., Crits-Christoph, P., Ball, W.A., Dewees, E., McAllister, T., Alahi, P., Cacciola, J., y Whybrow, P.C. (1991). Independent assessment of manic and depressive symptoms by self-rating: scales characteristics and implications for the study of mania. *Archives of general Psychiatry*, 48, 807-812.
- Baumgarten, M., Becker, R., y Gauthier, S. (1990). Validity and Reliability of the Dementia Behavior Disturbance Scale. *Journal of the American Geriatrics Society*, 38, 221-226.
- Bech, P., Rafaelsen, O.J., Kramp, P., y Bolwig, T.G. (1978). The Mania Rating Scale construction and interobserver agreement. *Neuropharmacology*, 17(6), 430-431.
- (1981). Rating scales for affective disorders: Their validity and consistency. *Acta Psychiatrica scandinavica*, 295(64), 1-101.
- Bechara, A., Damasio, A.R., Damasio, H., y Anderson, S. W. (1994). Insensitivity to future consequences following damage to human prefrontal cortex. *Cognition*, 5, 7-15.
- Bechara, A., y Damasio, H. (2002). Decision-making and addiction (part I): impaired activation of somatic states in substance dependent individuals when pondering decisions with negative future consequences. *Neuropsychologia*, 40(10), 1675-1689.
- Beck, A.T., Steer, R.A., y Brown, G.K. (1996). *Beck Depression Inventory-II*. San Antonio, TX: PsychCorp/Pearson.
- (2011a). *Manual. BDI-II. Inventario de Depresión de Beck-II* (adaptación española: Sanz, J., y Vázquez, C.). Madrid: Pearson Educación.

- (2011b). *Manual. BDI-FastScreen. Inventario de Depresión de Beck Fast-Screen para pacientes médicos* (adaptación española: Sanz, J., Izquierdo, A., y García-Vera, M. P.). Madrid: Pearson Educación.
- Beck, A.T., y Steer, R.A. (1993). *Beck Anxiety Inventory manual*. San Antonio, TX: PsychCorp/Pearson.
- Beck, A.T., Weissman, A., Lester, D., y Trexler, L. (1974). The measurement of pessimism: the hopelessness scale. *Journal of Consulting and Clinical Psychology*, 42, 861-865.
- Beck, C., Heithoff, K., Baldwin, B., Cuffel, B., O'Sullivan, P., y Chumbler, N.R. (1997). Assessing disruptive behavioral in older adults: the Disruptive Behavior Scale. *Aging & Mental Health*, 1(1), 71-79.
- Becoña-Iglesias, E., y Cortés-Tomás, M. (2010). *Manual de adicciones para psicólogos especialistas en psicología clínica en formación*. Madrid: So-cidrogalcohol.
- Belanger, H.G., Brown, L.M., Crowell, T.A., Vanderploeg, R.D., y Curtiss, G. (2002). The Key Behaviors Change Inventory and executive functioning in an elderly clinic sample. *Clinical Neuropsychology*, 16, 251-257.
- Benito-León, J., Martínez-Martín, P., Frades, B., et al. (2007). Impact of fatigue in multiple sclerosis: the Fatigue Impact Scale for Daily Use (D-FIS). *Multiple Sclerosis*, 13, 645-651.
- Benton, A., y Hamsher, K.S. (1989). *Multilingual aphasia examination*. Iowa City: University of Iowa.
- Benton, A.L., y Hamsher, K. (1976). *Multilingual Aphasia Examination*. Iowa City: University of Iowa Press.
- Bergner, M., Bobbit, R.A., Carter, W.B., y Gilson, B.S. (1981). The Sickness Impact Profile: Development and Final Revision of a Health Status Measure. *Medical Care*, 19, 787-805.
- Bernal Vargas, L., Riveros Munévar, F., Vinaccia Alpi, S., y Quiceno Sierra, J.M. (2017). Factorial structure and internal consistency of the Fatigue Severity Scale in Colombian population with chronic diseases. *Enfermería Global*, 46, 37-43.
- Berridge, K.C., y Robinson, T.E. (2016). Liking, wanting, and the incentive-sensitization theory of addiction. *American Psychologist*, 71, 670-679.
- Bibby, H., y McDonald, S. (2005). Theory of mind after traumatic brain injury. *Neuropsychologia*, 43(1), 99-114.
- Binder, L.M. (1993). Assessment of malingering after mild head trauma with the Portland Digit Recognition Test. *Journal of Clinical Experimental Neuropsychology*, 15, 170-182.
- Birmingham, E., Bischof, W.F., y Kingstone, A. (2008a). Gaze selection in complex social scenes. *Visual Cognition*, 16(2/3), 341-356.
- (2008b). Social attention and real world scenes: The roles of action, competition, and social content. *Quarterly Journal of Experimental Psychology*, 61(7), 986-999.

- Birmingham, E., Bischof, W.F., y Kingstone, A. (2009a). Get real! Resolving the debate about equivalent social stimuli. *Visual Cognition*, 17(6-7), 904-924.
- Bisiach, E., Vallar, G., Perani, D., Papagno, C., y Berti, A. (1986). Unawareness of disease following lesions of the right hemisphere: anosognosia for hemiplegia and anosognosia for hemianopia. *Neuropsychologia*, 24, 471-482.
- Blake, D.D., Weathers, F.W., Nagy, L.M., Kaloupek, D.G., Klauminzer, G., Charney, D.S., y Keane, T.M. (1990). A clinician rating scale for assessing current and lifetime PTSD: the CAPS-1. *Behavior Therapist*, 18, 187-188.
- Blijd-Hoogewys, E.M.A., Van Geert, P.L.C., Serra, M., y Minderaa, R. (2008). Measuring Theory of Mind in Children. Psychometric Properties of the ToM Storybooks. *Journal of Autism and developmental disorders*, 38, 1907-1930.
- (2010). Development of Theory-of-Mind and the Theory-of-Mind Storybooks-Research in Typically Developing Children and Children with Autism Spectrum Disorders. *European Psychiatric Review*, 3(2), 34-38.
- Boada-Grau, J., Merino-Tejedor, E., Gil-Ripoll, C., Segarra-Pérez, G., y Vigil-Colet, A. (2014). Adaptación al español del Inventario Multidimensional de Fatiga (MFI) al entorno laboral. *Universitas Psychologica*, 13(4), 15-23.
- Boada, M., Cejudo, J.C., Tàrraga, L., López, O., y Kaufer, D. (2002). Neuropsychiatric Inventory Questionnaire (NPI-Q): validación española de una forma abreviada del Neuropsychiatric Inventory (NPI). *Neurología*, 17(6), 317-323.
- Bobes, J., Badía, X., Luque, A., García, M., González, M.P., y Dal-Ré, R. (1999). Validación de las versiones en español de los cuestionarios Liebowitz Social Anxiety Scale, Social Anxiety and Distress Scale y Sheehan Disability Inventory para la evaluación de la fobia social. *Medicina Clínica*, 112, 520-538.
- Bobes, J., Calcedo-Barba, A., García, M., Francois, M., Rico-Villademoros, F., González, M.P., Bascarán, M.T., y Bousoño, M. (2000). Evaluación de las pruebas psicométricas de la versión española de cinco cuestionarios para la evaluación del trastorno de estrés postraumático. *Actas Españolas de Psiquiatría*, 28, 207-218.
- Bobes, J., García Portilla, M.P., Bascarán, M.T., Saiz, P.A., y Bousoño, M. (2002). *Banco de instrumentos básicos para la práctica de la psiquiatría clínica* (2.ª ed.). Barcelona: Ars Medica.
- Bolter, J.F., y Nicolls, R. (1991). *Multi-Digit Memory Test*. San Luis Obispo, CA: Wang Neuropsychological Laboratories.
- Boone, K.B. (2007). A Reconsideration of the Slick *et al.* (1999). Criteria for Malingered Neurocognitive Dysfunction (pp. 29-49). En K.B. Boone (ed.), *Assessment of Feigned Cognitive Impairment: A Neuropsychological Perspective*. Nueva York: The Guilford Press.
- Boone, K.B., Lu, P., y Herzberg, D. (2002). *The b Test*. Los Ángeles: Western Psychological Services.

- Boone, K.B., Lu, P., y Wen, J. (2005). Comparison of various RAVLT scores in the detection of noncredible memory performance. *Archives of Clinical Neuropsychology*, 20, 310-319.
- Borgaro, S.R., y Prigatano, G.P. (2003). Modification of the Patient Competency Rating Scale for use on an acute neurorehabilitation unit: the PCRS-NR. *Brain Injury*, 17(10), 847-853.
- Borghini, G., Astolfi, L., Vecchiato, G., Mattia, D., y Babiloni, F. (2014). Measuring neurophysiological signals in aircraft pilots and car drivers for the assessment of mental workload, fatigue and drowsiness. *Neuroscience Bio-behavioral Review*, 44, 58-75.
- Boyle, P.A., Malloy, P.F., Salloway, S., Cahn-Weiner, D.A., Cohen, R., y Cummings, J.L. (2003). Executive dysfunction and apathy predict functional impairment in Alzheimer disease. *American Journal of Geriatric Psychiatry*, 11, 214-21.
- Brandt, J. (1988). Malingered amnesia. En R. Rogers (ed.), *Clinical assessment of malingering and deception* (pp. 65-83). Nueva York: Guilford Press.
- Brandt, J., Rubinsky, E. y Lassen, G. (1985). Uncovering malingered amnesia. *Annals of the New York Academy of Sciences*, 444, 502-503.
- Brod, M., Stewart, A.L., Sands, L., y Walton, P. (1999). Conceptualization and measurement of quality of life in dementia: the Dementia Quality of Life instrument (DQoL). *Gerontologist*, 39, 25-35.
- Brown, L.J., Potter, J.F., y Foster, B.G. (1990). Caregiver burden should be evaluated during geriatric assessment. *Journal of the American Geriatric Society*, 38, 455-560.
- Brunier, G., y Graydon, J. (1996). A comparison of two methods of measuring fatigue in patients on chronic haemodialysis: visual analogue vs. Likert scale. *International Journal of Nursing Studies*, 33, 338-348.
- Buela-Casal, G., Guillén-Riquelme, A., y Seisdedos Cubero, N. (2011). *Cuestionario de Ansiedad Estado-Rasgo (STAI)*. Madrid: TEA Ediciones.
- Bulbena, A. (2000). *Medición clínica en psiquiatría y psicología*. Barcelona: Masson.
- Bulbena, A., Bobes, J., Luque, A., Dal-Ré, R., Ballesteros, J., y Ibarra, N. (2003). Validación de las versiones en español de la Clinical Anxiety Scale y del Physician Questionnaire para la evaluación de los trastornos de ansiedad. *Medicina Clínica (Barc)*, 121, 367-374.
- Burgener, S., y Twigg, P. (2002). Relationships among caregiver factors and quality of life in care recipients with irreversible dementia. *Alzheimer Disease and Associated Disorders*, 16, 88-102.
- Burns, A., Folstein, S., Brandt, J., y Folstein, M. (1990). Clinical assessment of irritability, aggression, and apathy in Huntington and Alzheimer disease. *Journal of Nerves Mental Disease*, 178, 20-26.
- Buschke, H. (1984). Cued recall in amnesia. *Journal of clinical neuropsychology*, 6, 433-440.

- Bush, S.S., Ruff, R.M., Troster, A.I., Barth, J.T., Koffler, S.P., Pliskin, N.H., *et al.* (2005). Symptom validity assessment: Practice issues and medical necessity (NAN Policy and Planning Committee). *Archives of Clinical Neuropsychology*, 20, 419-426.
- Buswell, G.T. (1935). *How people look at pictures*. Chicago: University of Chicago Press.
- Cacho, J., García-García, R., Arcaya, J., Vicente, J.L., y Lantada, N. (1999). Una propuesta de aplicación y puntuación del test del reloj en la enfermedad de Alzheimer. *Revista de neurología*, 28, 648-655.
- Cacioppo, J.T., Klein, D.J., Berntson, G.G., y Hatfield, E. (1993). The psychophysiology of emotion. En M. Lewis y J.M. Haviland (eds.), *Handbook of Emotions*. Nueva York: Guilford.
- Caeiro, L., Ferro, J.M., y Costa, J. (2013). Apathy secondary to stroke: a systematic review and meta-analysis. *Cerebrovascular Disease*, 35, 23-39.
- Cano, F.J., Rodríguez-Franco, L., y García Martínez, J. (2007). Adaptación española del Inventario de Estrategias de Afrontamiento. *Actas Españolas de Psiquiatría*, 35, 29-39.
- Carmona-Perera, M., Caracuel, A., Verdejo-García, A., y Pérez-García, M. (2013). Adaptation to Spanish population an instrument of moral judgments: Moore's battery of moral dilemmas. *Anales de Psicología*, 29(3), 827-835.
- Carrington, S.J., y Bailey, A.J. (2009). Are there theory of mind regions in the brain? A review of the neuroimaging literature. *Human Brain Mapping*, 30(8), 2313-2335.
- Carrobbles, J.A.I. (1981). Registros fisiológicos. En R. Fernández Ballesteros y J.A.I. Carrobbles (eds.), *Evaluación Conductual*, 364-406. Madrid: Ediciones Pirámide.
- Carver, C.S. (1997). You want to measure coping but your protocol's too long: Consider the Brief COPE. *International Journal of Behavioral Medicine*, 4(1), 92-100.
- Carver, C.S., Scheier, M.F., y Weintraub, J.K. (1989). Assessing coping strategies: A theoretically based approach. *Journal of Personality and Social Psychology*, 56, 267-283.
- Castaño-León, A.M., Navarro-Main, B., Gómez, P.A., Gil, A., Soler, M.D., Lagares, A., Bernabeu, M., Von Steinbüchel, N., y Real, R.G.L. (2017). Quality of Life After Brain Injury: Psychometric Properties of the Spanish Translation of the QOLIBRI. *Evaluation and Health Professions*, 1-18.
- Castaño-Monsalve, B., Bernabeu-Guitart, M., López, R., Bulbena-Vilasar, A., y Quemada, I. (2012). Psychopathological evaluation of traumatic brain injury patients with the Neuropsychiatric Inventory. *Revista Psiquiatría y Salud Mental*, 5(3), 160-166.
- Castellanos-Pinedo, F., Hernández-Pérez, J.M., Zurdo, M., Rodríguez-Fúnez, B., Hernández-Bayo, J.M., García-Fernández, C., *et al.* (2011). Influence of



- premorbid psychopathology and lesion location on affective and behavioral disorders after ischemic stroke. *Journal of Neuropsychiatry Clinical Neuroscience*, 23, 340-347.
- Cederlund, M., Hagberg, B., y Gillberg, C. (2010). Asperger syndrome in adolescent and young adult males. Interview, self- and parent assessment of social, emotional, and cognitive problems. *Research in Developmental Disabilities*, 31, 287-298.
- Chalder, T., Berelowitz, G., Pawlikowska, T., Watts, L., Wessely, S., Wright, D., et al. (1993). Development of a fatigue scale. *Journal of Psychosomatic Research*, 37, 147-153.
- Chambless, D.L., Caputo, G.C., Bright, P., y Gallagher, R. (1984). Assessment of fear of fear in agoraphobics: The Body Sensations Questionnaire and the Agoraphobic Cognitions Questionnaire. *Journal of Consulting and Clinical Psychology*, 52, 1090-1097.
- Chemerinski, E., y Robinson, R.G. (2000). The neuropsychiatry of stroke. *Psychosomatics*, 41, 5-14.
- Chiaravalloti, N.D., y DeLuca, J. (2003). Assessing the behavioral consequences of multiple sclerosis: an application of the Frontal Systems Behavior Scale (FrSBe). *Cognitive and Behavioral Neurology*, 16, 54-67.
- Chouinard, M.J., y Rouleau, I. (1997). The 48-Pictures Test: A two-alternative forced-choice recognition test for the detection of malingering. *Journal of the International Neuropsychological Society*, 3, 545-552.
- Chow, T.W., y Cummings, J. L. (1999). Frontal-subcortical circuits. En Miller, B. L., Cummings, J.L. (eds.), *The human frontal lobes. Functions and disorders*. Nueva York: The Guilford Press.
- Ciurli, P., Formisano, R., Bivona, U., Cantagallo, A., y Angelelli, P. (2011). Neuropsychiatric disorders in persons with severe traumatic brain injury: prevalence, phenomenology, and relationship with demographic, clinical, and functional features. *Journal of Head Trauma Rehabilitation*, 26, 116-126.
- Clarke, D.E., Kob, J.Y., Kuhla, E.A., Van Reekum, R., Salvadora, R., y Marind, R.S. (2011). Are the available apathy measures reliable and valid? A review of the psychometric evidence. *Journal of Psychosomatic Research*, 70, 73-97.
- Clarke, D.E., Van Reekum, R., Patel, J., Simard, M., Gomez, E., y Streiner, D.L. (2007). An appraisal of the psychometric properties of the clinician version of the Apathy Evaluation Scale (AES-C). *International Journal of Methods in Psychiatry Research*, 16, 97-110.
- Cohen-Mansfield, J., Marx, M.S., y Rosenthal, A.S. (1989). A Description of Agitation in a Nursing Home. *Journal of Gerontology*, 44(3), M77-84.
- Cohen, S., Kamarck, T., y Mermelstein, R. (1983). A global measure of perceived stress. *Journal of Health and Social Behavior*, 24(4), 385-396.
- Colom, F., Vieta, E., Martínez-Arán, A., García, M., Reinares, M., Torrent, C., Goikolea, J.M., Banús, S., y Salamero, M. (2002). Versión española de una

- escala de evaluación de la manía: validez y fiabilidad de la Escala de Manía de Young. *Medicina Clínica*, 119, 366-71.
- Comalli, P.E., Wapner, S., y Werner, H. (1962). Interference effects of Stroop Color-Word Test in childhood, adulthood, and aging. *Journal of Genetic Psychology*, 100, 47-153.
- Comeche, M.I., Díaz, M.I., y Vallejo, M.A. (1995). *Cuestionarios, inventarios y escalas: ansiedad, depresión, habilidades sociales*. Madrid: Fundación Universidad-Empresa.
- Conde-Sala, J.L., Reñé-Ramírez, R., Turró-Garriga, O., Gascón-Bayarri, J., Campdelacreu-Fumadó, J., Juncadella-Puig, M., Rico-Pons, I., y Garre-Olmo, J. (2014). Severity of dementia, anosognosia, and depression in relation to the quality of life of patients with Alzheimer disease: discrepancies between patients and caregivers. *American Journal of Geriatric Psychiatry*, 22(2), 138-147.
- Conde, V., Escriba, J.A., e Izquierdo, J.A. (1970). Evaluación estadística y adaptación castellana de la escala Autoplicada para la depresión de Zung. *Archivos de Neurobiología*, 33(2), 185-206.
- Conde, V., Esteban, T., y Useros, E. (1976). Revisión crítica de la adaptación castellana del Cuestionario de Beck. *Revista de Psicología General y Aplicada*, 31, 469-497.
- Conde, V., y Franch, J.I. (1984). *Escalas de evaluación comportamental para la cuantificación de la sintomatología psicopatológica en los trastornos angustiosos y depresivos*. Madrid: Upjohn Farmaquímica.
- Conder, R., Allen, L., y Cox, D. (1992). *Manual for the computerized assessment of response bias*. Durham, NC: CogShell.
- Connor, K.M., y Davidson, J.R.T. (1999). Further psychometric assessment of the TOP-8: a brief interview-based measure of PTSD. *Depress Anxiety*, 9(3), 135-137.
- Contador Castillo, I., Fernández Calvo, B., Cacho Gutiérrez, L.J., Ramos Campos, F., y Hernández Martín., L. (2009). Depresión en la demencia tipo alzhéimer: ¿existe algún efecto sobre la memoria? *Revista de Neurología*, 49(10), 505-510.
- Corcoran, R., Mercer, G., y Frith, C.D. (1995). Schizophrenia, symptomatology and social inference: investigating theory of mind in people with schizophrenia. *Schizophrenia Research*, 17, 5-13.
- Crespo-López, M., y López Martínez, J. (2005). Estrés y estado emocional en cuidadores de familiares mayores con y sin demencia. *Revista Española de Geriatria y Gerontología*, 40(3), 55-61.
- Crespo, M., y Cruzado, J.A. (1997). La evaluación del afrontamiento: adaptación española del cuestionario COPE con muestra de estudiantes universitarios. *Análisis y Modificación de Conducta*, 23, 797-830.
- Crespo, M., y Rivas, M.T. (2015). La evaluación de la carga del cuidador: una revisión más allá de la escala de Zarit. *Clínica y Salud*, 26(1), 9-15.

- Cripe, L.I. (1996). The ecological validity of executive function testing. En R.J. Sbordone y C.J. Long, *Ecological validity of neuropsychological testing*. Florida: GR Press. Lucie Press.
- Cullen, B., Coen, R.F., Lynch, C.A., Cunningham, C.J., Coakley, D., Robertson, I.H., *et al.* (2005). Repetitive behaviour in Alzheimer's disease: description, correlates and functions. *International Journal of Geriatric Psychiatry*, 20, 686-693.
- Cummings, J.L. (1997). The Neuropsychiatric Inventory: assessing psychopathology in dementia patients. *Neurology*, 48(5), 10-16.
- Cummings, J.L., Mega, M., Gray, K., Rosenberg-Thompson, S., Carusi, D.A., y Gornbein, J. (1994). The Neuropsychiatric Inventory: comprehensive assessment of psychopathology in dementia. *Neurology*, 44, 2308-2314.
- Curt, G.A., Breitbart, W., Cella, D., *et al.* (2000). Impact of cancer-related fatigue on the lives of patients: new findings from the Fatigue Coalition. *Oncologist*, 5, 353-360.
- Cutting, J. (1978). Study of anosognosia. *Journal of Neurology, Neurosurgery, and Psychiatry*, 41(6), 548-555.
- Damasio, A. (1994). *Descartes' error. Emotion, reason and the human brain*. Nueva York: Putnam's Sons.
- (1997). Towards a neuropathology of emotion and mood. *Nature*, 386, 769-770.
- (2006). *El error de Descartes* (5.<sup>a</sup> ed.). Barcelona: Editorial Crítica.
- Damasio, A.R., y Van Hoesen, G.W. (1984). Emotional disturbances associated with focal lesions of the limbic frontal lobe. En K.M. Heilman y P. Satz (eds.), *Neuropsychology of human emotion*. Nueva York: Guilford Press.
- Davidson, J.R., Potts, N.L., Richichi, E.A., Ford, S.M., Krishman, K.R., Smith, R.D., y Wilson, W. (1991). The brief social phobia scale. *Journal of Clinical Psychiatry*, 52, 48-51.
- De Arroyabe, E.L., Calvete, E., Hayas, C., y Zubizarreta, A. (2013). Distress of the Caregiver in Acquired Brain Injury: Positive aspects of care to moderate the effects of psychological problems. *The Australian Journal of Rehabilitation Counselling*, 19(2), 84-99.
- Decety, J., y Michalska, K. (2010). Neurodevelopmental changes in the circuits underlying empathy and sympathy from childhood to adulthood. *Developmental Science*, 13(1), 886-899.
- Deffenbacher, J.L., Demm, P.M., y Brandon, A.D. (1986). High general anger. *Behaviour Research and Therapy*, 24, 481-489.
- Delis, D.C., Kramer, H.H., Kaplan, E., y Ober, B.A. (1987). *California Verbal Learning Test*. San Antonio, TX: The Psychological Corporation.
- DeLongis, A., Folkman, S., y Lazarus, R.S. (1988). The impact of daily stress on health and mood: Psychological and social resources as mediators. *Journal of Personality and Social Psychology*, 54, 486-495.

- Denney, R.L. (1999). A brief symptom validity testing procedure for Logical Memory of the Weschler Memory Scale-Revised which can demonstrate verbal memory in the face of claimed disability. *Journal of Forensic Neuropsychology*, 1, 5-26.
- Derogatis, L.R. (1983). *Administration, Scoring and Procedures Manual II for the Revised Version of the SCL-90-R*. Baltimore: John Hopkins University Press.
- Derogatis, L.R., y Melisaratos, N. (1983). The Brief Symptom Inventory: An introductory report. *Psychological Medicine*, 13(3), 595-605.
- Derogatis, L. R. (2001). *Brief Symptom Inventory (BSI)-18. Administration, scoring and procedures manual*. Minneapolis: NCS Pearson.
- Devanand, D.P., Miller, L., Richards, M., Marder, K., Bell, K., Mayeux, R., y Stern, Y. (1992). The Columbia University Scale for Psychopathology in Alzheimer's Disease. *Archives of Neurology*, 49, 371-376.
- Devier, D.J., Pelton, G.H., Tabert, M.H., Liu, X., Cuasay, K., Eisenstadt, R., Marder, K., Stern, Y., y Devanand, D.P. (2009). The impact of anxiety on conversion from mild cognitive impairment to Alzheimer's disease. *International Journal of Geriatric Psychiatry*, 24, 1335-1342.
- Díaz-Olavarieta, C., Cummings, J.L., Velázquez, J., y García de la Cadena, C. (1999). Neuropsychiatric manifestations of multiple sclerosis. *Journal of Neuropsychiatry Clinical Neurosciences*, 11, 51-57.
- Díez Quevedo, C., Rangill, T., Sánchez-Planell, L., Kroenke, K., y Spitzer, R.L. (2001). Validation and utility of the patient health questionnaire in diagnosing mental disorders in 1003 general hospital Spanish inpatients. *Psychosomatic Medicine*, 63(4), 679-86.
- Donoso, A. (2002). Anosognosia en enfermedades cerebrales. *Revista Chilena de Neuropsiquiatría*, 40(2), 69-78.
- Drachman, D.A., Swearer, J.M., O'Donnell, B.F., Mitchell, A.L., y Maloon, A. (1992). The Caretaker Obstreperous-Behavior Rating Assessment (COBRA) Scale. *Journal of the American Geriatrics Society*, 40(5), 463-470.
- Duclos, H., Bejanin, A., Eustache, F., Desgranges, B., y Laisney, M. (2018). Role of context in affective theory of mind in Alzheimer's disease. *Neuropsychologia*, 119, 363-372.
- Dujardin, K., Sockeel, P., Delliaux, M., Destee, A., y Defebvre, L. (2008). The Lille Apathy Rating Scale: validation of a caregiver-based version. *Movement Disorders*, 23, 845-849.
- Durá, E., Andreu, Y., Galdón, M.J., Ferrando, M., Murgui, S., Poveda, R., y Jiménez, Y. (2006). Psychological assessment of temporomandibular disorders patients. Confirmatory analysis of the dimensional structure of the Brief Symptoms Inventory 18 (BSI-18). *Journal of Psychosomatic Research*, 60, 365-370.
- Echeburúa, E. (1995). *Evaluación y Tratamiento de la fobia Social*. Barcelona: Martínez Roca.

- Echeburúa, E., Corral, P., y Fernández-Montalvo, J. (2000). Escala de Inadaptación: propiedades psicométricas en contextos clínicos. *Análisis y Modificación de Conducta*, 26, 325-340.
- Ekman, P. (1972). Universals and Cultural Differences in Facial Expressions of Emotions. En J. Cole, *Nebraska Symposium on Motivation* (pp. 207-282). Lincoln, NB: University of Nebraska Press.
- Ekman, P., y Friesen, W. (1976). *Pictures of Facial Affect*. Palo Alto, CA: Consulting Psychologists Press.
- Epel, E.S., Blackburn, E.H., Lin, J., Dhabhar, F.S., Adler, N.E., Morrow, J.D., y Cawthon, R.M. (2004). Accelerated telomere shortening in response to life stress. *Proceedings of the National Academy of Sciences PNAS*, 101, 17312-17315.
- Ernst, M., y Fudge, J.L. (2009). A developmental neurobiological model of motivated behavior: anatomy, connectivity and ontogeny of the triadic nodes. *Neuroscience & Biobehavioral Reviews*, 33(3), 367-382.
- Ettema, T., Dröes, R., De Lange, J., Mellenbergh, G., y Ribbe, M. (2007). QUALIDEM: development and evaluation of a specific quality of life instrument-validation. *International Journal of Geriatric Psychiatry*, 22, 424-30.
- Euroqol Group (1990). Euroqol a new facility for the measurement of health related quality of life. *Health Policy*, 16, 199-208.
- Everitt, B.J., y Robbins, T.W. (2005). Neural systems of reinforcement for drug addiction: from actions to habits to compulsion. *Nature Neuroscience*, 8, 1481-1489.
- (2016). Drug addiction: updating actions to habits to compulsions ten years on. *Annual Review of Psychology*, 67, 23-50.
- Everly, G.S. (1989). *A clinical Guide to the treatment of the human stress response*. Nueva York: Plenum Press.
- Fabry, J.J. (1980). Depression. En R.H. Woody (ed.), *Encyclopedia of clinical assessment* (vol. 2). San Francisco: Jossey-Bass.
- Fahn, S., Elton, R.L., y UPDRS program members (1987). Unified Parkinsons Disease Rating Scale. En S. Fahn, C.D. Marsden, M. Goldstein y D.B. Calne (eds.), *Recent developments in Parkinsons disease* (vol. 2, pp. 153-163). Florham Park, NJ: Macmillan Healthcare Information.
- Faschingbauer, T., DeVaul, R., y Zisook, S. (1978). *Test ITRD. Texas Revised Inventory of Grief Manual*. Houston: Honeycomb Publications.
- Feinberg, T.E., Roane, D.M., y Ali, J. (2000). Illusory limb movements in anosognosia for hemiplegia. *Journal of Neurology, Neurosurgery and Psychiatry*, 68, 511-513.
- Fellows, L.K., y Farah, M.J. (2007). The role of ventromedial prefrontal cortex in decision making: judgment under uncertainty or judgment per se? *Cerebral Cortex*, 17(11), 2669-2674.

- Fernández-Abascal, E.G. (1997). Estilos y estrategias de afrontamiento. En E.G. Fernández-Abascal, F. Palmero, M. Chóliz y F. Martínez (eds.), *Cuaderno de prácticas de motivación y emoción*. Madrid: Pirámide.
- Fernández-Abascal, E.G., y Palmero, F. (1999). *Emociones y salud*. Barcelona: Ariel Psicología.
- Ferrero-Arias, J., Goñi-Imízcoz, M., González-Bernal, J., Lara-Ortega, F., Da Silva-González, Á., y Díez-Lopez, M. (2011). The Efficacy of Nonpharmacological Treatment for Dementia-related Apathy. *Alzheimer Disease & Associated Disorders*, 25(3), 213-219.
- Figved, N., Klevan, G., Myhr, K.M., Glad, S., Nyland, H., Larsen, J.P., et al. (2005). Neuropsychiatric symptoms in patients with multiple sclerosis. *Acta Psychiatrica Scandinavica*, 112, 463-468.
- Fisk, J.D., Pontefract, A., Ritvo, P.G., Archibald, C.J., y Murray, T.J. (1994). The impact of fatigue on patients with multiple sclerosis. *Canadian Journal of Neurological Science*, 21, 9-14.
- Fisk, J.D., Ritvo, P.G., Ross, L., Haase, D.A., Marrie, T.J., y Schleich, W.F. (1994). Measuring the functional impact of fatigue: initial validation of the fatigue impact scale. *Clinical Infectious Diseases*, 18, 79-83.
- Fisk, J.D., y Doble, S.E. (2002). Construction and validation of a fatigue impact scale for daily administration (D-FIS). *Quality of Life Research*, 11(3), 263-272.
- Fleming, J.M., Strong, J., y Ashton, R. (1996). Self-awareness of deficits in adults with traumatic brain injury: how best to measure? *Brain Injury*, 10, 1-15.
- Folkman, S., y Moskowitz, J.T. (2004). Coping: pitfalls and promise. *Annual Reviews of Psychology*, 55, 745-774.
- Frederick, R.I. (1997). *Validity Indicator Profile manual*. Minneapolis, MN: National Computer Services.
- Friborg, O., Hjemdal, O., Rosenvinge, J., y Martinussen, M. (2003). A new rating scale for adult resilience: What are the central protective resources behind healthy adjustment? *International Journal of Methods in Psychiatric Research*, 12, 65-76.
- Frith, C., y Frith, U. (2012). Social neuroscience. *Annual Review of Psychology*, 63, 287-313.
- Frith, C.D., y Frith, U. (2000). The physiological basis of theory of mind. En S. Baron-Cohen, H. Tager-Flusberg y D. Cohen (eds.), *Understanding Other Minds: Perspective from Developmental Social Neuroscience* (pp. 335-356). Oxford: Oxford University Press.
- Gallacher, J., Bayer, A., Fish, M., Pickering, J., Pedro, S., Dunstan, F., Ebrahim, S., y Ben-Sholmo, Y. (2009). Does anxiety affect risk of dementia?: Findings from the Caerphilly Prospective Study. *Psychosomatic Medicine*, 71, 659-666.

- Galli, S., Tatu, L., Bogousslavsky, J., y Aybek, S. (2018). Conversion, Factitious Disorder and Malingering: A Distinct Pattern or a Continuum? *Frontiers of Neurology Neuroscience*, 42, 72-80.
- García-García, J., Landa, V., Trigueros, M., Calvo, P., y Garminde, I. (1995). *Adaptación a nuestro medio del Inventario Texas Revisado de Duelo*. Barcelona: Actas del I Congreso de la Sociedad Española de Cuidados Palitativos.
- García-Ramos García, R. (2010). *Validación al castellano de la escala LARS de apatía en la enfermedad de Parkinson* (tesis doctoral). Madrid: Universidad Complutense de Madrid.
- García-Ramos, R., Villanueva-Iza, C., Catalán, M.J., Reig-Ferrer, A., y Matías-Guío, J. (2014). Validation of a Spanish Version of the Lille Apathy Rating Scale for Parkinson's Disease. *Scientific World Journal*, article ID 849834, 7.
- Garre-Olmo, J., Planas-Pujol, X., López-Pousa, S., Weiner, M.F., Turón-Estrada, A., Juvinyà, D., Ballester, D., y Vilalta Franch, J. (2010). Cross-cultural adaptation and psychometric validation of a Spanish version of the Quality of Life in Late-Stage Dementia Scale. *Quality of Life Research*, 19(3), 445-453.
- Gil-Moreno-De-Mora, G., Palmi Guerrero, J., y Prat-Subirana, J.A. (2017). Valoración de la percepción subjetiva de la fatiga en motoristas de competición Rally-Raid Dakar. *Acción Psicológica*, 14(1), 93-104.
- Gil, D., Fernández-Modamio, M., Bengochea, R., y Arrieta, M. (2012). Adaptación al español de la prueba de teoría de la mente Hinting Task. *Revista de Psiquiatría y Salud Mental*, 5(2), 79-88.
- Glaser, R., y Kiecolt-Glaser, J.K. (2005). Stress-induced immune dysfunction: implications for health. *Nature Reviews Immunology*, 5, 243-251.
- Glenn, M.B., O'Neil-Pirozzi, T., Goldstein, R., Burke, D., y Jacob, L. (2001). Depression amongst outpatients with traumatic brain injury. *Brain Injury*, 15, 811-818.
- Godfrey, H.P., Harnett, M.A., Knight, R.G., Marsh, N.V., Kesel, D.A., Partridge, F.M., et al. (2003). Assessing distress in caregivers of people with a traumatic brain injury (TBI): a psychometric study of the Head Injury Behaviour Scale. *Brain Injury*, 17(5), 427-435.
- Goetz, C.G., Tilley, B.C., Shaftman, S.R., Stebbins, G.T., Fahn, S., Martínez-Martín, P., et al. (2008). Movement Disorder Society-sponsored revision of the Unified Parkinson's Disease Rating Scale (MDS-UPDRS): scale presentation and clinimetric testing results. *Movement Disorders*, 23, 2129-2170.
- Goldberg, D., Bridges, K., Duncan-Jones, P., y Grayson, D. (1988). Detecting anxiety and depression in general medical settings. *British Medical Journal*, 297, 897-899.
- Goldberg, D.P., y Hillier, V.F. (1979). A scaled version of the General Health Questionnaire. *Psychological Medicine*, 9, 139-45.

- Goldstein, R.Z., y Volkow, N.D. (2002). Drug addiction and its underlying neurobiological basis: neuroimaging evidence for the involvement of the frontal cortex. *American Journal of Psychiatry*, 159, 1642-1652.
- Gómez Peña, M., Penelo, E., Fernández-Aranda, F., Álvarez Moya, E., Santamaría, J.J., et al. (2011). Motivation to change and pathological gambling: analysis of the relationship with clinical and psychopathological variables. *British Journal of Clinical Psychology*, 50(2), 196-210.
- González de la Rivera, J.L., De las Cuevas, C., Rodríguez Abuín, M., y Rodríguez Pulido, F. (2002). *SCL-90-R. Cuestionario de 90 síntomas*. Madrid: TEA.
- González Roscigno, E.A., Mujica Díaz, A.L., Terán Mendoza, O.E., Guerrero Alcedo, J.M., y Arroyo Alvarado, D. de J. (2016). Sintomatología frontal y trastornos de personalidad en usuarios de drogas ilícitas. *Drugs and Addictive Behavior*, 1(1), 17-28.
- Goodman, W.K., Price, L.H., Rasmussen, S.A., Mazure, C., Fleischmann, R.L., Hill, C.L., Heninger, G.R., y Charney, D.S. (1989). Yale-Brown Obsessive-Compulsive Scale, I: Development, use and reability. *Archives of General Psychiatry*, 46, 1006-1011.
- Goran, D.A., y Fabiano, R.J. (1993). The scaling of the Katz Adjustment Scale in a traumatic brain injury rehabilitation sample. *Brain Injury*, 7, 219-229.
- Gort, A.M., March, J., Gómez, X., De Miguel, M., Mazarico, S. y Ballesté, J. (2005). Escala de Zarit reducida en cuidados paliativos. *Medicina Clínica*, 124, 651-653.
- Gottfries, C.G., Brane, G., Gullberg, B., y Steen, G. (1982). A New Rating Scale for Dementia Syndromes. *Archives of Gerontology and Geriatrics*, 1, 311-330.
- Grace, J., Stout, J.C., y Malloy, P.F. (1999). Assessing frontal lobe behavioral syndromes with the Frontal Lobe Personality Scale. *Assessment*, 6, 269-284.
- Grace, J., y Malloy, P. (2001). *Frontal Systems Behavior Scale (FrSBe): Professional Manual*. Lutz, FL: Psychological Assessment Resources.
- Gray, J.M., Shepherd, M., y McKinlay, W.W. (1994). Negative symptoms in the traumatically brain-injured during the first year postdischarge, and their effect on rehabilitation status, work status and family burden. *Clinical Rehabilitation*, 8(3), 188-197.
- Green, P., Allen, L., y Astner, K. (1996). *The Word Memory Test: A manual for the oral and computerized forms*. Durham, NC: CogniSyst, Inc.
- Greiffenstein, M.F., y Baker, W.J. (2001). Comparison of premorbid and postinjury MMPI-2 profiles in late postconcussion claimants. *The Clinical Neuropsychologist*, 15, 162-170.
- Griffin, G.A., Glassmire, D.M., Henderson, E.H., y McCann, C. (1997). Rey II: Redesigning the Rey Screening Test of Malingering. *Journal of Clinical Psychology*, 53, 757-766.



- Grossman, A.B., Woolley-Levine, S., Bradley, W.G., y Miller, R.G. (2007). Detecting neurobehavioral changes in amyotrophic lateral sclerosis. *Amyotrophic lateral sclerosis and other motor neuron disorders*, 8, 56-61.
- Gualart-Balet, M., Jiménez-Cortés, M.P., Tuquet-Calvo, H., Pelegrín-Valero, C., Olivera-Pueyo, J., Benabarre-Ciria, S., y Tirapu-Ustároz, J. (2015). Validación española de la Iowa Rating Scale for Personality Change (IRSPC) para la valoración de los cambios de personalidad en pacientes con daño cerebral adquirido. *Revista de Neurología*, 60(1), 17-29.
- Guerra-Silla, M.G., Gutiérrez-Robledo, L.M., Villalpando-Berumen, J.M., Pérez-Zepeda, M.U., Montana-Álvarez, M., Reyes-Guerrero, J., y Rosas-Carrasco, O. (2011). Psychometric evaluation of a Spanish language version of the Screen for Caregiver Burden (SCB) in caregivers of patients with mixed, vascular and Alzheimer's dementia. *Journal of Clinical Nursing*, 20, 23-24.
- Hama, S., Yamashita, H., Shigenobu, M., Watanabe, A., Hiramoto, K., Kurisu, K., et al. (2007). Depression or apathy and functional recovery after stroke. *International Journal of Geriatric Psychiatry*, 22(10), 1046-1051.
- Hama, S., Yamashita, H., Shigenobu, M., Watanabe, A., Kurisu, K., Yamawaki, S., y Kitaoka, T. (2007). Post-stroke affective or apathetic depression and lesion location: Left frontal lobe and bilateral basal ganglia. *European Archives of Psychiatry and Clinical Neuroscience*, 257, 149-152.
- Hamilton, M. (1959). The assessment of anxiety states by rating. *British Journal of Medicine and Psychology*, 32, 50-55.
- (1960). A Rating scale for depression. *Journal of Neurology, Neurosurgery and Psychiatry*, 23, 56-62.
- Hammond, F.M., Davis, C.S., Whiteside, Y.O., Philbrick, P., y Hirsch, M.A. (2011). Marital adjustment and stability following traumatic brain injury: a pilot qualitative analysis of spouse perspectives. *The Journal of Head Trauma Rehabilitation*, 26, 69-78.
- Happé, F. (1995). *Autism: An introduction of Psychological Theory*. Harvard: Harvard University Press.
- Happé, F., y Frith, U. (1994). Theory of mind in autism. En E. Schopler y G. Mesibov (eds.), *Learning and cognition in autism*. Nueva York: Plenum Press.
- Hart, T., Whyte, J., Kim, J., y Vaccaro, M. (2005). Executive function and self-awareness of real-world behavior and attention deficits following traumatic brain injury. *Journal of Head Trauma Rehabilitation*, 20, 333-347.
- Hartman Maier, A., Soroker, N., Oman, S.D., y Katz, N. (2001). Anosognosia for hemiplegia in stroke rehabilitation. *Neurorehabilitation and neural Repair*, 15, 213-222.
- Hartman Maier, A., Soroker, N., Oman, S.D., y Katz, N. (2003). Awareness of disabilities in stroke rehabilitation clinical trial. *Disability and Rehabilitation*, 7, 25(1), 35-44.

- Hartman Maeir, A., Soroker, N., Ring, H., y Katz, N.J. (2002). Awareness of deficits in stroke rehabilitation. *Rehabilitation Medicine*, 34(4), 158-64.
- Havet-Thomassin, V., Allain, P., Etcharry-Bouyx, F., y Le Gall, D. (2006). What about theory of mind after severe brain injury? *Brain Injury*, 20(1), 83-91.
- Haynes, S.N., Levine, S., Scotch, N., Feinleib, M., y Kannel, W.B. (1978). The relationship of psychological factors to coronary heart disease in the Framingham Study: I. Methods and risk factors. *American Journal of Epidemiology*, 107, 362-363.
- Herrero, M.J., Blanch, J., Peri, J.M., De Pablo, J., Pintor, L., y Bulbena, A. (2003). A validation study of the hospital anxiety and depression scale (HADS) in a Spanish population. *General Hospital Psychiatry*, 25(4), 277-283.
- Hiscock, M., y Hiscock, C.K. (1989). Refining the forced choice method for the detection of malingering. *Journal of Clinical and Experimental Neuropsychology*, 11, 967-974.
- Hobfoll, S.E., Dunahoo, C.L., y Monnier, J. (1993). Preliminary test manual: *The strategic approach to coping scale (SACS)*. Manuscrito no publicado. Kent State University.
- Hogrefe, G.J., Wimmer, H., y Perner, J. (1986). Ignorance versus false belief: A developmental lag in attribution of epistemic states. *Child Development*, 57, 567-582.
- Homaifar, B.Y., Brenner, L.A., Gutiérrez, P.M., Harwood, J.F., Thompson, C., Filley, C.M., Kelly, J.P. y Adler, L.E. (2009). Sensitivity and specificity of the Beck Depression Inventory-II in persons with traumatic brain injury. *Archives of Physical Medicine and Rehabilitation*, 90, 652-656.
- Hoshmand, L., y Austin, G. (1987). Validation studies of a multifactor cognitive-behavioral Anger Control Inventory. *Journal of Personality Assessment*, 51, 417-432.
- Huertas-Hoyas, E., Pedrero-Pérez, E.J., Martínez-Campos, M., y Laselle-López, M. (2016). Inventario de síntomas prefrontales (ISP) en daño cerebral adquirido: concordancia entre puntuaciones de paciente, familiar y profesional. *Revista de Neurología*, 63, 385-92.
- Huizink, A.C., Mulder, E.J., Robles de Medina, P.G., Visser, G.H., y Buitelaar, J.K. (2004). Is pregnancy anxiety a distinctive syndrome? *Early Human Development*, 79, 81-91.
- Iqbala, N., Caswell, H.L., Harea, D.J., Pilkington, O., Mercer, S., y Duncan, S. (2009). Neuropsychological profiles of patients with juvenile myoclonic epilepsy and their siblings: a preliminary controlled experimental video-EEG case series. *Epilepsy Behavior*, 14, 516-21.
- Iruthayarajah, J., Alibrahim, F., Mehta, S., Janzen, S., McIntyre, A., y Teasell, R. (2018). Cognitive behavioural therapy for aggression among individuals with moderate to severe acquired brain injury: a systematic review and meta-analysis. *Brain Injury*, 1-7.

- Itier, R.J., Villate, C., y Ryan, J.D. (2007). Eyes always attract attention but gaze orienting is task-dependent: Evidence from eye movement monitoring. *Neuropsychologia*, 45(5), 1019-1028.
- Iverson, G., Franzen, M., y McCracken, L. (1991). Evaluation of an objective assessment technique for the detection of malingered memory deficits. *Law and Human Behavior*, 15, 667-676.
- Jacobsen, B.S., y Lowery, B.J. (1992). Further analysis of the psychometric properties of the Levine Denial of Illness Scale. *Psychosomatic Medicine*, 54(3), 372-381.
- Janik, S.W., Wellens, R., Goldberg, M.L., y Delloso, L.F. (1978). Eyes at the centre of focus in the visual examination of human faces. *Perceptual and Motor Skills*, 47(3), 857-858.
- Johnson, J., y McCown, W. (1997). *Family Therapy of Neurobehavioral Disorders. Integrating Neuropsychology and Family therapy*. Nueva York: The Haworth Press, Inc.
- Jones, B., y Mishkin, M. (1972). Limbic lesions and the problem of stimulus-reinforcement associations. *Experimental Neurology*, 36, 362-377.
- Jorge, R.E. (2015). Mood disorders. *Handbook of Clinical Neurology*, 128, 613-631.
- Kaland, N., Moller-Nielsen, A., Callesen, K., Mortensen, E., Gottlieb, D., y Smith, L. (2002). A new “advanced” test of theory of mind: evidence from children and Adolescents with Asperger syndrome. *Journal of Clinical Psychology and Psychiatry*, 43, 517-528.
- Kaland, N., Moller-Nielsen, A., Smith, L., Mortensen, E., Callesen, K., y Gottlieb, D. (2005). The Strange Stories test-a replication study of children and adolescents with Asperger syndrome. *European Journal of Child and Adolescent Psychiatry*, 14, 73-82.
- Kaland, N., Smith, L., y Mortensen, E.L. (2007). Response times of children and adolescents with Asperger syndrome on an ‘advanced’ test of theory of mind. *Journal of Autism and Developmental Disorders*, 37, 197-209.
- Kanner, A.D., Coyne, J.C., Schaefer, C., y Lazarus, R.S. (1981). Comparison of two modes of stress measurement: daily hassles and uplifts versus mayor life events. *Journal of Behavior Medicine*, 4, 1-39.
- Kaplan, E.F., Goodglass, H., y Weintraub, S. (1983). *Boston Naming Test*. Filadelfia: Lea & Febiger.
- (2001). *The Boston Naming Test* (2.<sup>a</sup> ed.). Filadelfia: Lippincott Williams & Wilkins.
- Kapur, N. (1994). The coin-in-the-hand test: a new “bed-side” test for the detection of malingering in patients with suspected memory disorder. *Journal of Neurology, Neurosurgery and Psychiatry*, 57, 385-386.
- Kaszniak, A.W., y Allender, J. (1985). Psychological assessment of depression in older adults. En G.M. Chaisson-Stewart (ed.), *Depression in the elderly: An interdisciplinary approach*. Nueva York: Wiley.

- Kay, S.R., Fiszbein, A., Vital-Herne, M., y Fuentes, L.S. (1990). The Positive and Negative Syndrome Scale-Spanish adaptation. *Journal of Nervous Mental Disease*, 178, 510-517.
- Kay, S.R., Opler, L.A., y Lindenmayer, J.P. (1989). The Positive and Negative Syndrome Scale (PANSS): rationale and standardisation. *British Journal of Psychiatry, suppl.* (7), 59-67.
- Kelly, G., Brown, S., Todd, J., y Kremer, P. (2008). Challenging behaviour profiles of people with acquired brain injury living in community settings. *Brain Injury*, 22(6), 457-470.
- Kerr, S.L., y Neale, J.M. (1993). Emotion perception in schizophrenia: specific deficit or further evidence of generalized poor performance? *Journal of Abnormal Psychology*, 102, 312-318.
- Kilmer, R., Demakis, G., Hammond, F., Grattan, K., Cook, J., y Kornev, A. (2006). Use of the Neuropsychiatric Inventory in traumatic brain injury: a pilot investigation. *Rehabilitation Psychology*, 51, 232-238.
- Kim, S.H., Facundo, M., Todd, K., Sankar, B., y Robinson, R. (1999). Irritability after traumatic brain injury. *The Journal of Nervous and Mental Disease*, 6, 327-335.
- Kitwood, T., y Bredin, K.A. (1992). New approach to the evaluation of dementia care. *Journal of Advanced Health Nursing Care*, 1, 41-60.
- Kivela, S.L. (1992). Psychological assessment and rating scales: Depression and other age-related affective disorders. En M. Bergener *et al.* (eds.), *Aging and mental disorders*. Nueva York: Springer.
- Knesevich, J.W., Martin, R.L., Berg, L., y Danziger, W. (1983). Preliminary report on affective symptoms in the early stages of senile dementia of the Alzheimer's type. *American Journal of Psychiatry*, 140, 233-235.
- Knight, R., Devereux, R., y Godfrey, H. (1998). Caring for a family member with a traumatic brain injury. *Brain Injury*, 12, 467-481.
- Knight, R.G., Ross, R.A., Collins, J.I., y Parmenter, S.A. (1985). Some norms, reliability and preliminary validity data for an S-ROS: Inventory of Anger: The Subjective Anger Scale (SAS). *Personality and Individual Differences*, 6, 331-339.
- Koenigs, M., Huey, E.D., Calamia, M., Raymond, V., Tranel, D., y Grafman, J. (2008). Distinct regions of prefrontal cortex mediate resistance and vulnerability to depression. *Journal of Neuroscience*, 28, 12341-12348.
- Koenigs, M., Young, L., Adolphs, R., Tranel, D., Hauser, M., y Damasio, A. (2007). Damage to the prefrontal cortex increases utilitarian moral judgments. *Nature*, 447(7138), 808-911.
- Kolb, B., y Whishaw, I.Q. (2006). *Neuropsicología humana*. Madrid: Médica Panamericana.
- Kolitz, B.P., Vanderploeg, R.D., y Curtiss, G. (2003). Development of the Key Behaviors Change Inventory: a traumatic brain injury behavioral outcome

- assessment instrument. *Archives of Physical Medicine and Rehabilitation*, 84, 277-84.
- Koob, G.F., Stinus, L., Le Moal, M., y Bloom, F.E. (1989). Opponent process theory of motivation: neurobiological evidence from studies of opiate dependence. *Neuroscience & Biobehavioral Reviews*, 13, 135-140.
- Korkman, M., Kirk, U., y Temp, S. (2007). *NEPSY II*. San Antonio, TX: Psychological Corporation.
- Korkman, M., Kirk, U., y Temp, S. (2014). *NEPSY II*. Madrid: Pearson Educación.
- Kreutzer, J.S., Marwitz, J.H., Seel, R., y Serio, C.D. (1996). Validation of a neurobehavioral functioning inventory for adults with traumatic brain injury. *Archives of Physical Medicine and Rehabilitation*, 77(2), 116-124.
- Krupp, L.B., LaRocca, N.G., Muir-Nash, J., y Steinberg, A.D. (1989). The fatigue severity scale Application to patients with multiple sclerosis and systemic lupus erythematosus. *Archives of Neurology*, 46, 1121-1123.
- Kulisevsky, J., Litvan, I., Berthier, M.L., Pascual-Sedano, B., Paulsen, J.S., y Cummings, J.L. (2001). Neuropsychiatric assessment of Gilles de la Tourette patients: comparative study with other hyperkinetic and hypokinetic movement disorders. *Movement Disorders*, 16, 1098-1104.
- Kuopio, A.M., Marttila, R.J., Helenius, H., Toivonen, M., y Rinne, U.K. (2000). The quality of life in Parkinson's disease. *Movement Disorders*, 15, 216-223.
- Kwag, K.H., Martin, P., Russell, D., Franke, W., y Kohut, M. (2011). The impact of perceived stress, social support, and home-based physical activity on mental health among older adults. *International Journal of Aging and Human Development*, 72, 137-154.
- Lane-Brown, A.T., y Tate, R.L. (2009). Measuring apathy after traumatic brain injury: psychometric properties of the Apathy Evaluation Scale and the Frontal Systems Behavior Scale. *Brain Injury*, 23, 999-1007.
- Lansing, C.R., y McConkie, G.W. (2003). Word identification and eye fixation locations in visual and visual-plus-auditory presentations of spoken sentences. *Perception & Psychophysics*, 65(4), 536-552.
- Larson, M.J., Perlstein, W.M., Demery, J.A., y Stigge-Kaufman, D.A. (2006). Cognitive control impairments in traumatic brain injury. *Journal of Clinical and Experimental Neuropsychology*, 28, 968-86.
- Lawton, M.P., Moos, M., Kleban, M.H., Glicksman, A., y Rovine, M. (1991). A two-factor model of caregiving appraisal and psychological well-being. *J. Gerontol B Psychol Sci Soc Sci*, 46, 181-9.
- Lazarus, R.S., y Folkman, S. (1984). Coping and adaptation. En W.D. Gentry (ed.), *The handbook of behavioural medicine*. Nueva York: Guilford.
- (1984), *Stress, appraisal, and coping*. Nueva York: Springer.
- (1989), *Estrés y procesos cognitivos*. Barcelona: Editorial Martínez Roca.

- Le Heron, C., Apps, M.A.J., y Husaina, M. (2018). The anatomy of apathy: A neurocognitive framework for amotivated behaviour. *Neuropsychologia*, *118*, 54-67.
- Le Moal, M., Stinus, L., y Simon, H. (1979). Increased sensitivity to (+)amphetamine self-administered by rats following meso-cortico-limbic dopamine neurone destruction. *Nature*, *280*(5718), 156-158.
- Lebert, F., Pasquier, F., y Petit, H. (1996). Évaluation Comportementale Dans la Démence de Type Alzheimer par le Questionnaire de Dyscontrôle Comportemental. *Presse Medicale*, *25*(14), 665-667.
- Lee, K.A., Hicks, G., y Nino-Murcia, G. (1991). Validity and reliability of a scale to assess fatigue. *Psychiatry Research*, *36*, 291-298.
- Leentjens, A.F., Dujardin, K., Marsh, L., Martinez-Martin, P., Richard, I.H., Starkstein, S.E., Weintraub, D., Sampaio, C., Poewe, W., Rascol, O., Stebbins, G.T., y Goetz, C.G. (2008). Apathy and anhedonia rating scales in Parkinson's disease: Critique and recommendations. *Movement Disorders*, *23*, 2004-2014.
- León-Salas, B., Olazarán-Rodríguez, J., Dobato-Ayuso, J.L., Prieto-Jurczynska, C., Agüera-Ortiz, L., Osorio-Suárez, R., et al. (2009). Propiedades psicométricas de la versión española de la escala Quality of Life-Alzheimer's Disease (QOL-AD) en pacientes con enfermedad de Alzheimer institucionalizados. *Neurología*, *24*, 715-716.
- Lerdal, A., Kottorp, A., Gay, C., Aouizerat, B.E., Portillo, C.J., y Lee, K.A. (2011). A 7-item version of the fatigue severity scale has better psychometric properties among HIV-infected adults: an application of a Rasch model. *Quality of Life Research*, *20*, 1447-1456.
- Lerdal, A., y Kottorp, A. (2011). Psychometric properties of the Fatigue Severity Scale—Rasch analyses of individual responses in a Norwegian stroke cohort. *International Journal of Nursing Studies*, *48*, 1258-1265.
- Levenstein, S., Prantera, C., Varvo, V., Scribano, M.L., Berto, E., Luzzi, C., y Andreoli, A. (1993). Development of the Perceived Stress Questionnaire: a new tool for psychosomatic research. *Journal of Psychosomatic Research*, *37*(1), 19-32.
- Levin, H.S., High, W.M., Goethe, K.E., Sisson, R.A., Overall, J.E., Rhoades, H.M., Eisenberg, H.M., Zalisky, Z., y Gary, H.E. (1987). The Neurobehavioral Rating Scale: Assessment of the behavioral sequelae of head injury by the clinician. *Journal of Neurology, Neurosurgery and Psychiatry*, *50*, 183-193.
- Levine, J., Rudy, T., y Kerns, R. (1994). A two factor model of denial of illness: aconfirmatory factor analysis. *Journal of Psychosomatic Research*, *38*(2), 99-110.
- Levine, J., Warrenburg, S., Kerns, R., Schwartz, G., Delanay, R., Fontana, A., Gradman, A., Smith, S., Allen, S., y Cascione, R. (1987). The role of denial

- in recovery from coronary heart disease. *Psychosomatic Medicine*, 49(2), 109-117.
- Levy, M.L., Cummings, J.L., Fairbanks, L.A., Masterman, D., Miller, B.L., Craig, A.H., *et al.* (1998). Apathy is not depression. *Journal of Neuropsychiatry and Clinical Neuroscience*, 10, 314-319.
- Levy, R., y Dubois, B. (2006). Apathy and the functional anatomy of the prefrontal cortex-basal ganglia circuits. *Cerebral Cortex*, 16, 916-928.
- Lewis, D.J., y Duncan, C.P. (1956). Effect of different percentages of money reward on extinction of a lever-pulling response. *Journal of Experimental Psychology*, 52(1), 23.
- (1957). Explication and resistance to extinction of a lever-pulling response as functions of percentage of reinforcement and amount of reward. *Journal of Experimental Psychology*, 54(2), 23.
- Lewis, M. (2015). *The biology of desire. Why addiction is not a disease*. Nueva York: Public Affairs.
- (2017). Addiction and the brain: Development, not disease. *Neuroethics*, 10, 7-18.
- Lezak, M.D. (1995). *Neuropsychological assessment* (3.<sup>a</sup> ed). Nueva York: Oxford University Press.
- Liebowitz, M.R. (1987). Social Phobia. *Modern Problems of Pharmacopsychiatry*, 22, 141-173.
- Ligthart, S.A., Richard, E., Franssen, N.L., Eurelings, L.S.M., Beem L., Eikelenboom P., van Gool W.A., y Moll van Charante E.P. (2012). Association of vascular factors with apathy in community-dwelling elderly individuals. *Archives of General Psychiatry*, 69, 636-642.
- Litvan, I., Cummings, J.L., y Mega, M.S. (1998). Neuropsychiatric features of corticobasal degeneration, *Journal of Neurology Neurosurg Psychiatry*, 65, 717-721.
- Livianos, L., Rojo, L., Guillén, J.L., Villavencio, D., Pino, A., Mora, R., Vila, M.L. y Domínguez, A. (2000). Adaptación de la Escala para la Valoración de la Manía por Clínicos (EVMAC). *Actas Españolas de Psiquiatría*, 28 (3), 169-177.
- Llanero-Luque, M., Ruiz-Sánchez de León, J.M., Pedrero-Pérez, E., Olivar-Arroyo, A., Bouso-Saiz, J.C., Rojo-Mota, G., y Puerta-García, C. (2008). Sintomatología disejecutiva en adictos a sustancias en tratamiento mediante la versión española del Cuestionario Disejecutivo (DEX-Sp). *Revista de Neurología*, 47(9), 1-15.
- Lobo, A., Chamorro, L., Luque, A., Dal-Re, R., Badia, X., y Baro, E. (2002). Validación de las versiones en español de la Montgomery-Asberg Depression Rating Scale y la Hamilton Anxiety Rating Scale para la evaluación de la depresión y de la ansiedad. *Med Clin (Barc)*, 118(13), 493-499.

- Lobo, A., Pérez-Echevarría, M.J., y Artal, J. (1986) Validity of the scaled version of the General Health Questionnaire (GHQ-28) in a Spanish population. *Psychological Medicine*, 16, 135-140.
- Logsdon, R.G., Gibbons, L.E., McCurry, S.M., y Teri, L. (1999). Quality of life in Alzheimer's disease: patient and caregiver reports. *Journal of Mental Health and Aging*, 5, 21-32.
- López de Arroyabe, E., y Calvete, E. (2005). Síntomas de duelo en familiares de personas con daño cerebral adquirido. Grief symptoms in relatives of people with traumatic brain injury. *Mapfre Medicina*, 16, 266-276.
- (2012). Evaluación de las necesidades de los familiares de personas afectadas de Daño Cerebral Adquirido mediante el Cuestionario de Necesidades Familiares. *Anales de Psicología*, 28, 728-735.
- López de Arroyabe, E., Calvete, E., Las Hayas, C., y Zubizarreta, A. (2013). Distress of the caregiver in acquired brain injury: Positive aspects of care to moderate the effects of psychological problems. *Australian Journal of Rehabilitation Counselling*, 19(2), 84-99.
- López, S.R., y Moral, M.S. (2005). Validación del Índice de Esfuerzo del Cuidador en la población española. *Enfermería Comunitaria*, 1, 12-17.
- Losada, A., Montorio, I., Izal, M. y Márquez, M. (2006). *Estudio e intervención sobre el malestar psicológico de los cuidadores de personas con demencia. El papel de los pensamientos disfuncionales*. Madrid: Instituto de Mayores y Servicios Sociales (IMSERSO).
- Lozoya-Delgado, P., Ruiz-Sánchez de León, J.M., y Pedrero-Pérez, E.J. (2012). Validación de un cuestionario de quejas cognitivas para adultos jóvenes: relación entre las quejas subjetivas de memoria, la sintomatología prefrontal y el estrés percibido. *Revista de Neurología*, 54, 137-150.
- Lu, P., Boone, K.B., Jiménez, N., y Razani, J. (2004). Failure to inhibit the reading response on the Stroop Test: A pathognomonic indicator of suspect effort. *Journal of Clinical and Experimental Neuropsychology*, 26, 180-189.
- Lucas-Carrasco, R., Gómez-Benito, J., Rejas, J., y Ott, B.R. (2013). The cornell brown scale for quality of life in dementia: Spanish adaptation and validation. *Alzheimer Disease and Associated Disorders*, 27(1), 44-50.
- Lucas-Carrasco, R., Lamping, D.L., Banerjee, S., Rejas, J., Smith, S.C., y Gómez-Benito, J. (2010). Validation of the spanish version of the demqol system. *International Psychogeriatrics*, 22(4), 589-597.
- Lucas-Carrasco, R., y Gómez-Benito, J. (2009). Psychometric properties of the Spanish version of the DQOL. *XII Congress of the International Federation of Psychiatric Epidemiology (IFPE)*. Viena, Austria.
- Lueken, U., Seidl, U., Volker, L., Schweiger, E., Kruse, A., y Schroder, J. (2007). Development of a short version of the Apathy Evaluation Scale specifically adapted for demented nursing home residents. *American Journal of Geriatric Psychiatry*, 15, 376-85.



- Lupien, S.J., De Leon, M., De Santi, S., Convit, A., Tarshish, C., y Nair, N.P. (1998). Cortisol levels during human aging predict hippocampal atrophy and memory deficits. *Nature Neuroscience*, 1, 69-73.
- Malec, J.F., y Lezak, M.D. (2003). *Manual for the Mayo-Portland Adaptability Inventory (MPAI-4) for adults, children and adolescents*. Portland, OR: Oregon Health and Sciences University.
- Man, W., Polkey, M., Donaldson, N., Gray, B., y Moxham, J. (2004). Community pulmonary rehabilitation after hospitalisation for acute exacerbations of chronic obstructive pulmonary disease. *BMJ*, 200329, 1209-1213.
- Manes, F., Paradiso, S., y Robinson, R.G. (1999). Neuropsychiatric effects of insular stroke. *The Journal of Nervous and Mental Disease*, 187, 707-712.
- Manzano, J.M.A., Llorca, G., Ledesma, A., y López-Ibor, J.J. (1994). Adaptación Española de la Alzheimer's Disease Assessment Scale (ADAS). *Actas Luso-Españolas de Neurología, Psiquiatría y Ciencias Afines*, 22(2), 64-70.
- Marcel, A., Tegne'r, R., y Nimmo-Smith, I. (2004). Anosognosia for plegia: specificity, extension, partiality and disunity of bodily unawareness. *Cortex*, 40, 19-40.
- Marin, R.S. (1991). Apathy: a neuropsychiatric syndrome. *Journal of Neuropsychiatry and Clinical Neuroscience*, 3, 243-54.
- Marin, R.S., Biedrzycki, R.C., y Firinciogullari, S. (1991). Reliability and validity of the Apathy Evaluation Scale. *Psychiatry Research*, 38, 143-62.
- Marin, R.S., Firinciogullari, S., y Biedrzycki, R.C. (1993). The sources of convergence between measures of apathy and depression. *Journal of Affective Disorders*, 28, 117-24.
- Marks I.M., y Mathews A.M. (1979). Brief standard self-ratings for phobic patients. *Behaviour Research and Therapy*, 17, 263-267.
- Marsh, N.V., Kersel, D.A., Havill, J.H., y Sleigh, J.W. (1998). Caregiver burden at 1 year following severe traumatic brain injury. *Brain Injury*, 12(12), 1045-1059.
- Martí-Esquitino, J.(2015). *Patología de la conciencia: la anosognosia*. Málaga: Monsalvez2007.
- Martín, M., Salvadó, I., Nadal, S., Miji, L.C., Rico, J.M., Lanz, P., y Taussing, M.I. (1996). Adaptación para nuestro medio de la Escala de Sobrecarga del Cuidador (Caregiver Burden Interview) de Zarit. *Revista de Gerontología*, 6, 338-346.
- Martínez de la Iglesia, J., Onís, M.C., Dueñas, H.R., Albert, C.C., Aguado, T.C., y Luque, L.R. (2002). Versión española del cuestionario de Yesavage abreviado (GDS) para el despistaje de depresión en mayores de 65 años: adaptación y validación. *Revista de Medicina Familiar y Comunitaria*, 12, 620-630.
- Martínez-Martín, P. (2004). Calidad de vida. *Medicina Clínica (Barc)*, 5, 86-90.

- Martínez-Martín, P., Catalán, M., Benito-León, J., *et al.* (2006). Impact of fatigue in Parkinson's disease: the Fatigue Impact Scale for Daily Use (D-FIS). *Quality of Life Research*, 15, 597-606.
- Martínez-Martín, P., Rodríguez-Blázquez, C., Álvarez-Sánchez, M., Arakaki, T., Bergareche-Yarza, A., Chade, A., *et al.* (2013). Expanded and independent validation of the Movement Disorder Society-Unified Parkinson's Disease Rating Scale (MDS-UPDRS). *Journal of Neurology*, 260(1), 228-236.
- Martínez-Martín, P., Skorvanek, M., Rojo-Abuín, J.M., Gregova, Z., Stebbins, G.T., Goetz, C.G., y miembros del QUALPD Study Group (2018). Validation study of the hoehn and yahr scale included in the MDS-UPDRS. *Movement Disorders*, 33(4), 651-652.
- Mathews, A.M., Gelder, M., y Johnston, D.W. (1986 [1981]). *Agorafobia. Naturaleza y tratamiento*. Barcelona: Fontanella.
- Mathias, J.L. (2003). Neurobehavioral functioning of persons with Parkinson's disease. *Applied Neuropsychology*, 10, 57-68.
- Mazaux, J.M., Masson, F., Levin, H.S., Alaoui, P., Maurette, P., y Barat, M. (1997). Long-term neuropsychological outcome and loss of social autonomy after traumatic brain injury. *Archives of Physical Medicine and Rehabilitation*, 78(12), 1316-1320.
- McAdams, L.A., Harris, M.J., Heaton, S.C., Bailey, A., Fell, R., y Jeste, D.V. (1997). Validity of specific subscales of the positive and negative symptom scales in older schizophrenia outpatients. *Schizophrenia Research*, 27, 219-226.
- McConaughy, E.A., Prochaska, J.O., y Velicer, W.F. (1983). Stages of change in psychotherapy: Measurement and sample profiles. *Psychotherapy: Theory, Research and Practice*, 20, 368-375.
- McDonald, C.R., Norman, M.A., Delis, D.C., Tecoma, E.S., y Iragui, V.J. (2004). Cognitive inhibition, mood, and behavior in frontal lobe epilepsy: relationship to seizure lateralization and structural MRI findings. *Epilepsia*, 45, 346.
- McEwen, B.S. (1999). Stress and hippocampal plasticity. *Annual Review of Neuroscience*, 22, 105-122.
- McGivney, S.A., Mulvihill, M., y Taylor, B. (1994). Validating the GDS depression screen in the nursing home. *Journal of the American Geriatrics Society*, 42, 490-492.
- McIntosh, R.C., Rosselli, M., Uddin, L.Q., y Antoni, M. (2015). Neuropathological sequelae of Human Immunodeficiency Virus and apathy: a review of neuropsychological and neuroimaging studies. *Neuroscience Biobehavioral Reviews*, 55, 147-164.
- McLean, A.J., Temkin, N.R., Dikmen, S., y Wyler, A.R. (1983). The behavioral sequelae of head injury. *Journal of Clinical and Experimental Neuropsychology*, 5, 361-376.

- Mega, M.S., Cummings, J.L., Fiorello, T., y Gornbein, J. (1996). The spectrum of behavioral changes in Alzheimer's disease. *Neurology*, *46*, 130-135.
- Mendoza, Y., Cuello, P., y López, V. (2016). Análisis psicométrico del inventario de sintomatología prefrontal en sujetos adictos y no adictos. *Revista Chilena de Neuropsicología*, *11*, 24-29.
- Merchán Dueñas, T.C. (2008). *Validación de la Apathy Evaluation Scale modificada de Marín (ASP-mM), para la evaluación de apatía, en pacientes con enfermedad de Parkinson que acuden a la consulta externa del Servicio de Neurología del Hospital Carlos Andrade Marín de la ciudad de Quito* [tesis doctoral]. Ecuador: Universidad Católica del Ecuador.
- Michaud, K., Forget, H., y Cohen, H. (2009). Chronic glucocorticoid hypersecretion in Cushing's syndrome exacerbates cognitive aging. *Brain Cognition*, *71*, 1-8
- Migliorelli, R., Tesón, A., Sabe, L., Petracca, G., Petracchi, M., Leiguarda, R., y Starkstein, S.E. (1995). Anosognosia in Alzheimer's disease: a study of associated factors. *Journal Neuropsychiatry and Clinical Neurosciences*, *7*, 338-344.
- Miguel Tobal, J.J., Casado Morales, M.I., Cano Vindel, A., y Spielberger, C. (2001). *Manual del Inventario de Expresión de Ira Estado-Rasgo. STAXI-2*. Madrid: TEA Ediciones.
- Miller, W.R. (1983). Motivational interviewing with problem drinkers. *Behavioural Psychotherapy*, *11*, 147-172.
- Miller, W.R., y Rose, G.S. (2009). Toward a theory of motivacional Interviewing. *American Psychologist*, *64*, 527-537.
- Miller, W.R., y Tonigan, J.S. (1996). Assessing drinkers' motivation for change: The Stages of Change Readiness and Treatment Eagerness Scale (SOCRATES). *Psychology of Addictive Behaviors*, *10*, 81-89.
- Millis, S.R., y Putnam, S.H. (1994). The Recognition Memory Test in the assessment of memory impairment after financially compensable mild head injury: A replication. *Perceptual and Motor Skills*, *79*, 384-386.
- Moldover, J.E., Goldberg, K.B., y Prout, M.F. (2004). Depression after traumatic brain injury: a review of evidence for clinical heterogeneity. *Neuropsychology Review*, *14*(3), 143-154.
- Molinero, C., Bonete, S., Gómez-Pérez, M.M., y Calero, M.D. (2015). Estudio normativo del "test de 60 caras de Ekman" para adolescentes españoles. *Behavioral Psychology/Psicología Conductual*, *23*(2), 361-371.
- Monnier, J., Hobfoll, S.E., Dunahoo, C.L., Hulsizer, M.R., y Jonson, R. (1998). There's more than rugged individualism in coping. Part 2: Construct validity and further model testing. *Anxiety, Stress, and Coping*, *11*, 247-272.
- Montgomery, S.A., y Asberg, M. (1979). A new depression scale designed to be sensitive to change. *British Journal of Psychiatry*, *134*, 382-389.

- Montón, C., *et al.* (1993). Escalas de ansiedad y depresión de Goldberg: una guía de entrevista eficaz para la detección del malestar psíquico. *Atención Primaria*, 12, 345-349.
- Moore, A., Clark, B., y Kane, M. (2008). Who Shalt Not Kill? Individual Differences in Working Memory Capacity, Executive Control and Moral Judgment. *Psychological Science*, 6(19), 549-557.
- Morawetz, C., Bode, S., Baudewig, J., y Heekeren, H.R. (2017). Effective amygdala-prefrontal connectivity predicts individual differences in successful emotion regulation. *Social Cognitive and Affective Neuroscience*, 12(4), 569-585.
- Morgan, J.E., y Sweet, J.J. (2009). *Neuropsychology of malingering casebook*. Nueva York: Psychology Press.
- Motzkin, J.C., Philippi, C.L., Wolf, R.C., Baskaya, M.K., y Koenigs, M. (2015). Ventromedial prefrontal cortex is critical for the regulation of amygdala activity in humans. *Biological Psychiatry*, 77(3), 276-284.
- Movement Disorder Society Task Force on Rating Scales for Parkinson's Disease (2003). The Unified Parkinson's Disease Rating Scale (UPDRS): status and recommendations. *Movement Disorders*, 18(7), 738-750.
- Müllera, T., y Appsa, M.A.J. (2019). Motivational fatigue: A neurocognitive framework for the impact of effortful exertion on subsequent motivation. *Neuropsychologia*, 123, 141-151.
- Muñoz Céspedes, J.M., y Tirapu Ustárroz, J. (2001). *Rehabilitación Neuropsicológica*. Madrid: Síntesis.
- Muñoz Céspedes, J.M., Fernández Guinea, S., y Pelegrín Valero, C. (1999). La intervención neuropsicológica en las personas con daño cerebral traumático. Una perspectiva integradora. En A. Ruano, J.M. Muñoz Céspedes y C. Cid Rojo (coords.), *Psicología de la Rehabilitación*. Madrid: Mapfre.
- Muñoz Céspedes, J.M., Pérez-Nieto, M.A., e Iruarrizaga, I. (2001). La evaluación y tratamiento de la ira en personas con daño cerebral traumático. Una perspectiva neuropsicológica. *Revista de Psicología General y Aplicada*, 54(3), 407-423.
- Muris, P., Steerneman, P., Meesters, C., Merckelbach, H., Horselenberg, R., Van den Hogen, T., y Van Dongen, L. (1999). The TOM Test: A new instrument for assessing theory of mind in normal children and children with pervasive developmental disorders. *Journal of Autism and Developmental Disorders*, 29, 67-80.
- Murman, D.L., Chen, Q., Powell, M.C., Kuo, S.B., Bradley, C.J., y Colenda, C.C. (2002). The incremental direct costs associated with behavioral symptoms in AD. *Neurology*, 59(11), 1721-1729.
- Murphy, M.E. (2010). *Everyday executive function in people with schizophrenia: investigating the Dysexecutive Questionnaire*. Kansas City: University of Missouri.

- Myers, C., y Wilks, D. (1999). Comparison of Euroqol EQ-5D and SF-36 in patients with chronic fatigue syndrome. *Quality of Life Research*, 8, 9-16.
- Naqvi, N.H., Rudrauf, D., Damasio, H., y Bechara, A. (2007). Damage to the insula disrupts addiction to cigarette smoking. *Science*, 315(5811), 531-534.
- Nelson, L., Satz, P., y D'Elia, L.F. (1994). *Neuropsychology Behavior and Affect Profile*. Palo Alto, CA: Mind Garden.
- Nitch, S. R., y Glassmire, D. M. (2007). Non-forced-choice measures to detect noncredible cognitive performance. En K. B. Boone (Ed.), *Assessment of feigned cognitive impairment: A neuropsychological perspective* (p. 78-102). The Guilford Press.
- Novaco, R.W. (1975). *Anger Control: The Development and Evaluation of an Experimental Treatment*. Lexington: D.C. Heath.
- (1977). Stress inoculation: A cognitive therapy for anger and its application to a case of depression. *Journal of Consulting and Clinical Psychology*, 45, 600-608.
- O'Bryant, S., Hilsabeck, R., Fisher, J., y McCaffrey, R. (2003). Utility of the Trail Making Test in the assessment of malingering in a sample of mild traumatic brain injury. *The Clinical Neuropsychologist*, 17, 69-74.
- Omdal, R., Sjøholm, H., Koldingnes, W., et al. (2005). Fatigue in patients with lupus is not associated with disturbances in cerebral blood flow as detected by SPECT. *Journal of Neurology*, 252, 78-83.
- Ongur, D. (2000). The organization of networks within the orbital and medial prefrontal cortex of rats, monkeys and humans. *Cerebral Cortex*, 10(3), 206-219.
- Orfei, M.D., Robinson, R.G., Prigatano, G.P., Starkstein, S., Rüsçh, N., Bria, P., Caltagirone, C., y Spalletta, G. (2007). Anosognosia for hemiplegia after stroke is a multifaceted phenomenon: a systematic review of the literature. *Brain*, 130(12), 3075-3090.
- Ott, B.R., Lafleche, G., Whelihan, W.M., Buongiorno, G.W., Albert, M.S., y Fogel, B.S. (1996). Impaired awareness of deficits in Alzheimer disease. *Alzheimer Disease and Associated Disorders*, 10, 68-76.
- Overall, J.E., y Gorham, D.R. (1962). The Brief Psychiatric Rating Scale. *Psychology Reports*, 10, 790-812.
- Owensworth, T.L., McFarland, K., y Young, R.M. (2000). Development and Standardization of the Self-Regulation Skills Interview (SRSI): A New Clinical Assessment Tool for Acquired Brain Injury. *The Clinical Neuropsychologist*, 14(1), 76-92.
- Palmero, F., Fernández-Abascal, E.G., Martínez, F., y Chóliz, M. (2002). *Psicología de la motivación y la emoción*. Madrid: McGraw Hill.
- Pascual, L.F., Saz, P., Larumbe, R., Martínez-Lage, P., Muruzábal, J., Morales, F., Lobo, A., Martínez-Lage, J.M., y Sastre y Hernández M. (1997). Estandarización en una población española de la escala ADAS (Alzheimer's Disease Assessment Scale). *Neurología*, 12(6), 24-32.

- Patel, V., y Hope, R.A. (1992). A rating scale for aggressive behaviour in the elderly—the RAGE. *Psychological Medicine*, 22(1), 211-221.
- Paul, D., Franzen, M.D., Cohen, S.H., y Fremouw, W. (1992). An investigation into the reliability and validity of two tests used in the detection of dissimulation. *International Journal of Neuropsychology*, 14, 1-9.
- Paulsen, J.S., Ready, R.E., Hamilton, J.M., Mega, M.S., y Cummings, J.L. (2001). Neuropsychiatric aspects of Huntington's disease, *Journal of Neurology Neurosurg Psychiatry*, 71, 310-314.
- Paz Domingo, M.V., Ruiz Sánchez de León, J.M., Paz Solís, J.F., Gandía González, M., Mateos González, A., y Pedrero Pérez, E.J. (2017). Dolor crónico: relación con sintomatología prefrontal y estrés percibido. *Revista de la Sociedad Española de Dolor*, 24(4), 179-187.
- Pearlin, L.I., Mullan, J.T., Semple, S.J., y Skaff, M.M. (1990). Caregiving and the stress process: An overview of concepts and their measures. *Gerontologist*, 30, 583-94.
- Pedersen, K.F., Larsenm, J.P., y Aarsland, D. (2008). Validation of the Unified Parkinson's Disease Rating Scale (UPDRS) section I as a screening and diagnostic instrument for apathy in patients with Parkinson's disease. *Parkinsonism and Related Disorders*, 14, 183-186.
- Pedrero-Pérez, E., Ruiz-Sánchez de León, J.M., Llanero-Luque, M., Rojo-Mota, G., Olivar-Arroyo, A., y Puerta García, C. (2009a). Sintomatología frontal en adictos a sustancias en tratamiento mediante la versión española de la Escala de Comportamiento Frontal. *Revista de Neurología*, 48(12), 624-631.
- (2011). Evaluación de los síntomas prefrontales: propiedades psicométricas y datos normativos del Cuestionario Disejutivo (DEX) en una muestra de población española. *Revista de Neurología*, 52(7), 394-404.
- Pedrero-Pérez, E., Ruiz-Sánchez de León, J.M., Rojo-Mota, G., Llanero-Luque, M., Olivar-Arroyo, A., Bouso-Saiz, J.C., y Puerta García, C. (2009b). Versión española del Cuestionario Disejutivo (DEX-Sp): propiedades psicométricas en adictos y población no clínica. *Adicciones*, 21(2), 155-166.
- Pedrero-Pérez, E.J., y Ruiz-Sánchez de León, J.M. (2013). Quejas subjetivas de memoria, personalidad y sintomatología prefrontal en adultos jóvenes. *Revista de Neurología*, 57(7), 289-96.
- (2019). Síntomas prefrontales en la vida diaria: normalización y estandarización del inventario de síntomas prefrontales abreviado en población española. *Revista de Neurología*, 68(11), 459-467.
- Pedrero-Pérez, E.J., Rojo-Mota, G., Ruiz-Sánchez de León, J.M., Fernández-Méndez, L.M., Morales-Alonso, S., y Prieto-Hidalgo, A. (2014). Reserva cognitiva en adictos a sustancias en tratamiento: relación con el rendimiento cognitivo y las actividades cotidianas. *Revista de Neurología*, 59 (11), 481-489.

- Pedrero-Pérez, E.J., Ruiz Sánchez de León, J.M., Morales-Alonso, S., Pedrero-Aguilar, J., y Fernández-Méndez, L.M. (2015a). Sintomatología prefrontal en la vida diaria: evaluación de cribado mediante el Inventario de Síntomas Prefrontales Abreviado (ISP-20). *Revista de Neurología*, 60(9), 385-93.
- Pedrero-Pérez, E.J., Ruiz-Sánchez de León, J.M. y Winpenny-Tejedor, C. (2015b). Dysexecutive Questionnaire (DEX): Unrestricted structural analysis in large clinical and non-clinical samples. *Neuropsychological Rehabilitation*, 25(6), 879-894.
- Pedrero-Pérez, E.J., Ruiz-Sánchez de León, J.M., Lozoya-Delgado, P., Rojo-Mota, G., Llanero-Luque, M., y Puerta García, C. (2013). Sintomatología prefrontal y trastornos de la personalidad en adictos a sustancias. *Revista de Neurología*, 56(4), 205-213.
- Pedrero-Pérez, E.J., Ruiz-Sánchez de León, J.M., Lozoya-Delgado, P., Rojo-Mota, G., Llanero-Luque, M., y Puerta-García, C. (2015c). La “Escala de Estrés Percibido”: Estudio psicométrico sin restricciones en población no clínica y adictos a sustancias en tratamiento. *Behavioral Psychology*, 23(2), 305-324.
- Pedrero-Pérez, E.J., Ruiz-Sánchez de León, J.M., Rojo-Mota, G., Morales-Alonso, S., Pedrero-Aguilar, J., Lorenzo-Luque, I., y González, A. (2016). Inventario de síntomas prefrontales (ISP): validez ecológica y convergencia con medidas neuropsicológicas. *Revista de Neurología*, 63(6), 241-51.
- Pedrero-Pérez, E.J., Santed Germán, M.Á. y Pérez García, A.M. (2012). Adaptación española de la Escala multiaxial de afrontamiento estratégico (SACS) de Hobfoll. *Psicothema*, 24, 455-460.
- Pelegrín-Valero, C., Martín-Carrasco, M., y Tirapu-Ustarroz, J. (1995). La escala NRS: La versión española de la Neurobehavioural Rating Scale. *Anales de Psiquiatría*, 2(3), 22-32.
- Peña-Casanova, J. (2019). *Programa integrado de exploración neuropsicológica Test Barcelona 2*. Barcelona: Test Barcelona Services, S.L.
- Peña-Casanova, J., Aguilar, M., Santacruz, P., Bertran-Serra, I., Hernández, G., Sol, J.M., Pujol, A., Blesa, R. y Grupo NORMACODEM (1997). Adaptación y Normalización españolas de la Alzheimer’s Disease Assessment Scale (ADAS). *Neurología*, 12(2), 69-77.
- Peña-Casanova, J., Gramunt-Fombuena, N., Quiñones-Ubeda, S., Sánchez-Benavides, G., Aguilar, M., Badenes, D., et al. (2009a). Spanish multicenter normative studies (NEURONORMA project): norms for the Rey-Osterrieth complex figure (copy and memory), and free and cued selective reminding test. *Archives of Clinical Neuropsychology*, 24, 371-393.
- Peña-Casanova, J., Quiñones-Ubeda, S., Gramunt-Fombuena, N., Aguilar, M., Casas, L., Molinuevo, J.L., et al. (2009b). Spanish multicenter normative studies (NEURONORMA project): norms for Boston naming test and token test. *Archives of Clinical Neuropsychology*, 24, 343-354.

- Peña-Casanova, J., Quiñones-Ubeda, S., Quintana-Aparicio, M., Aguilar, M., Badenes, D. Molinuevo, J.L., *et al.* (2009c). Spanish Multicenter Normative Studies (NEURONORMA Project): Norms for Verbal Span, Visuospatial Span, Letter and Number Sequencing, Trail Making Test, and Symbol Digit Modalities Test. *Archives of Clinical Neuropsychology*, 24, 321-41.
- Peterson, R.A., y Reiss, S. (1993). *Anxiety Sensitivity Index Revised Test Manual*. Worthington, OH: International Diagnosis System Publishing Corporation.
- Pia, L., Neppi-Modona, M., Ricci, R., y Berti, A. (2004). The anatomy of anosognosia for hemiplegia: a meta-analysis. *Cortex; a Journal Devoted to the Study of the Nervous System and Behavior*, 40(2), 367-377.
- Plutchik, R., Van Praag, H. M., Conte, H. R., y Picard, S. (1989). Correlates of Suicide and Violence Risk: The Suicide Risk Measure. *Comprehensive Psychiatry*, 30(4), 296-302.
- Ponsford, J., Olver J., Ponsford, M., y Nelms, R. (2003). Long-term adjustment of families following traumatic brain injury where comprehensive rehabilitation has been provided. *Brain Injury*, 17(6), 153-168.
- Porzolt, F., Kojer, M., Schmidl, M., Greimel, E., Sigle, J., Richter, J., *et al.* (2004). A new instrument to describe indicators of well-being in old-old patients with severe dementia-The Vienna List. *Health and Quality of Life Outcomes*, 2, 1-8.
- Premack, D., y Woodruff, G. (1978). Does chimpanzee have a theory of mind??. *Behavioral Brain Sciences*, 4, 9-30.
- Prigatano, G.P., Bruna, O., Mataró, M., Muñoz Céspedes, J.M., Fernández, S., y Junqué, C. (1998). Initial disturbances of consciousness and resultant impaired awareness in spanish patients with traumatic brain injury. *Journal of Head Trauma Rehabilitation*, 13(5), 1-15.
- Prigatano, G.P. (1992). Personality disturbances associated with traumatic brain injury. *Journal of Consulting and Clinical Psychology*, 60(3), 360-368.
- (2005). Disturbances of self-awareness and rehabilitation of patients with traumatic brain injury: a 20-year perspective. *The Journal of Head Trauma Rehabilitation*, 20(1), 19-29.
- Prigatano, G.P., y Klonoff, P.S. (1998). A Clinician's Rating Scale for Evaluating Impaired Self-Awareness and Denial of Disability After Brain Injury. *The Clinical Neuropsychologist*, 12(1), 56-67.
- Prigatano, G.P., y Schacter, D.L. (1991). *Awareness of deficits after brain injury: clinical and theoretical issues*. Nueva York: Oxford University Press.
- Prigatano, G.P., y Fordyce, D.J. (1986). Cognitive dysfunction and psychosocial adjustment after brain injury. En G.P. Prigatano, D.J. Fordyce, H.K. Zeiner, J.R. Roueche, M. Pepping y B.Wood (eds.), *Neuropsychological rehabilitation after brain injury* (pp. 1-17). Baltimore, MD: The Johns Hopkins University Press.



- Prochaska, J.O., y DiClemente, C.C. (1983). Stages and processes of self-change of smoking. Towards an integrative model of change. *Journal of Consulting and Clinical Psychology*, 51, 390-395.
- Pugh, R. (1983). An association between hostility and poor adherence to treatment in patients suffering from depression. *British Journal of Medical Psychology*, 56(2), 205-208.
- Pujol, J., Aspiazu, P., Salamero, M., y Cuevas, R. (2001). Sintomatología depresiva de la demencia. Escala de Cornell: validación de la versión en castellano. *Revista de Neurología*, 33(4), 397-398.
- Rabins, P.V., Kasper, J.D., Kleinman, L., y Black, B.S. (1999). Concepts and methods in the development of the ADRQL: an instrument for assessing health-related quality of life in persons with Alzheimer's disease. *Journal of Mental Health and Aging*, 5, 33-48.
- Radloff, L. (1977). The CES-D Scale: A self-report depression scale for research in the general population. *Applied Psychological Measurement*, 1, 385-401.
- Raich, R.M. (1997). Evaluación y tratamiento de un caso de trastorno obsesivo-compulsivo. En D. Maciá y F. Méndez (eds.), *Aplicaciones clínicas de la evaluación y modificación de conducta*. Madrid: Pirámide.
- Ramos-Brieva, J.A., Montejo Iglesias, M.L., Lafuente López, R., Ponce de León Hernández, C., y Moreno Sarmiento, A. (1991). Validación de la Escala-Criba Geriátrica para la Depresión. *Actas Luso-Españolas de Neurología, Psiquiatría y Ciencias Afines*, 19, 174-177.
- Ramos-Brieva, J.A. (1986). Validación de la versión castellana de la escala de Hamilton para la depresión. *Actas Luso-Esp Neurol Psiquiatr*, 14, 324-334.
- Ramos, F. (1982). *Personalidad, depresión y muerte* [tesis doctoral]. Madrid: Editorial de la Universidad Complutense de Madrid.
- Ratnavalli, E., Brayne, C., Dawson, K., y Hodges, J.R. (2002). The prevalence of frontotemporal dementia. *Neurology*, 58(11), 1615-1621.
- Ravina, B., Camicioli, R., Como, P.G., Marsh, L., Jankovic, J., Weintraub, D., y Elm, J. (2007). The impact of depressive symptoms in early Parkinson disease. *Neurology*, 69(4), 342-347.
- Ready, R., Ott, B., Grace, J., y Fernández, I. (2002) The Cornell-Brown Scale for Quality of Life in Dementia. *Alzheimer Disease and Associated Disorders*, 16, 109-215.
- Ready, R.E., Friedman, J., Grace, J., y Fernandez, H. (2004). Testosterone deficiency and apathy in Parkinson's disease: a pilot study. *Journal of Neurology Neurosurgery and Psychiatry*, 75, 1323-1326.
- Ready, R.E., Ott, B.R., Grace, J., y Cahn-Weiner, D.A. (2003). Apathy and executive dysfunction in mild cognitive impairment and Alzheimer disease. *American Journal of Geriatric Psychiatry*, 11, 222-228.
- Redish, A.D., Jensen, S., y Johnson, A. (2008). Addiction as vulnerabilities in the decision process. *Behavioral and Brain Sciences*, 31, 461-487.

- Reisberg, B., Borenstein, J., Salob, S.P., Ferris, S.H., Franssen, E., y Georgotas, A. (1987). Behavioral symptoms in Alzheimer's disease: phenomenology and treatment. *Journal of Clinical Psychiatry*, 48 supl, 9-15.
- Reitan, R.M. (1985). Validity of the Trail Making Test as an indication of organic brain damage. *Perceptual Motor Skills*, 8, 271-276.
- Reitan, R.M., y Wolfson, D. (1985). *The Halstead-Reitan neuropsychological test battery. Theory and clinical interpretation* [2.<sup>a</sup> ed.]. Tucson, AZ: Neuropsychology Press.
- Remor, E. (2006). Psychometric properties of a European Spanish version of the perceived Stress Scale (PSS). *Spanish Journal of Psychology*, 9, 86-93.
- Remor, E.A., y Carrobles, J.A. (2001). Versión española de la Escala de estrés percibido (PSS-14): estudio psicométrico en una muestra VIH+. *Ansiedad y Estrés*, 7, 195-201.
- Resnick, B., Zimmerman, S.I., Magaziner, J., y Adelman, A. (1998). Use of the Apathy Evaluation Scale as a measure of motivation in elderly people. *Rehabilitation Nursing*, 23, 141-147.
- Rey, A. (1941). L'examen psychologie dans las cas d'encephalopathie traumatique. *Archives de Psychologie*, 28, 286-340.
- (1964). *The clinical exam in psychology*. París: Presses Universitaires de France.
- Ríos, J.R., y Hernández, M. (2005). Prevalencia de depresión en pacientes con epilepsia y su calidad de vida. *Revista de Sanidad Militar*, 59, 311-317.
- Robert, P.H., Claiet, S., Benoit, M., Koutaich, J., Bertogliati, C., Tible, O., et al. (2002). The Apathy Inventory: assessment of apathy and awareness in Alzheimer's disease, Parkinson's disease and mild cognitive impairment. *International Journal of Geriatric Psychiatry*, 17, 1099-1105.
- Robinson, B.C. (1983). Validation od caregiver strain index. *Journal of Gerontology*, 38(3), 344-348.
- Robinson, R.G., Boston, J.D., Starkstein, S.E., y Price, T.R. (1988). Comparison of mania with depression following brain injury: casual factors. *American Journal of Psychiatry*, 145, 172-178.
- Robinson, T.E., y Berridge, K.C. (1993). The neural basis of drug craving: an incentive-sensitization theory of addiction. *Brain Research Reviews*, 18, 247-291.
- Rodríguez-Marín, J., Terol, M.C., López-Roig, S., y Pastor, M.A. (1992). Evaluación del afrontamiento del estrés: propiedades psicométricas del cuestionario de formas de afrontamiento de acontecimientos estresantes. *Revista de Psicología de la Salud*, 4, 59-84.
- Rodríguez-Martos, A., Rubio, G., Aubà, J., Santo-Domingo, J., Torralba, L., y Campillo, M. (2000). Readiness to change questionnaire: Reliability study of its Spanish version. *Alcohol & Alcoholism*, 35, 270-275.

- Rogers, R., Harrell, E. H. y Liff, C. D. (1993). Feigning neuropsychological impairment: A critical review of methodological and clinical considerations. *Clinical Psychology Review*, 13(3), 255-274.
- Rollnick, S., Heather, N., Gold, R., y Hall, W. (1992). Development of a short "readiness to change" questionnaire for use in brief, opportunistic interventions among excessive drinkers. *British Journal of Addiction*, 87, 743-754.
- Román, F., Rojas, G., Román, N., Iturry, M., Blanco, R., Leis, A., Bartoloni, L., Allegri, R., y Argencog, X. (2012). Baremos del Test de la Mirada en español en adultos normales de Buenos Aires. *Revista Neuropsicología Latinoamericana*, 4(3), 1-5.
- Rosen, H.J., Allison, S.C., Schauer, G.F., Gorno-Tempini, M.L., Weiner, M.W., y Miller B.L. (2005). Neuroanatomical correlates of behavioural disorders in dementia. *Brain*, 128, 2612-25.
- Rosen, J., Burgio, L., Kollar, M., Cain, M., Allison, M., Fogleman, M., Michael, M., y Zubenko, G.S. (1994). The Pittsburgh Agitation Scale: A User-Friendly Instrument for Rating Agitation in Dementia Patients. *American Journal of Geriatric Psychiatry*, 2(1), 52-59.
- Rosen, W.G., Mohs, R.C., y Davis, K.L. (1984) A New Rating Scale for Alzheimer's Disease. *American Journal of Psychiatry*, 141(11), 1356-1364.
- Rosenthal, I.A., Hutcherson, C.A., Adolphs, R., y Stanley, D.A. (2019). Deconstructing theory-of-mind impairment in high-functioning adults with autism. *Current biology*, 29(3), 513-519.
- Ross, S.R., Benning, S.D., y Adams, Z. (2007). Symptoms of executive dysfunction are endemic to secondary psychopathy: an examination in criminal offenders and noninstitutionalized young adults. *Journal of Personality Disorders*, 21, 384-99.
- Rubio, G., Montero, I., Jáuregui, J., Villanueva, R., Marín, J.J., y Santodomingo, J. (1998). Validación de la escala de riesgo suicida de Plutchik en población española. *Arch. Neurobiol.*, 61, 143-152.
- Ruiperez, M.A., Ibañez, M.I. Lorente, E., Moro, M., y Ortet, G. (2001). Psychometric Properties of the Spanish Version of BSI. *European Journal of Psychological Assessment*, 17(3), 241-250.
- Ruiz-Sánchez de León, J.M., Pedrero Pérez, E.J., Lozoya Delgado, P., Llanero Luque, M., Rojo Mota, G., y Puerta García, C. (2012). Inventario de Síntomas Prefrontales (ISP) para la evaluación clínica de las adicciones en la vida diaria: proceso de creación y propiedades psicométricas. *Revista de Neurología*, 54(11), 649-663.
- Ruiz-Sánchez de León, J.M., Pedrero-Pérez, E.J. y Lozoya-Delgado, P. (2014). Caracterización neuropsicológica de las quejas de memoria en la población general: relación con la sintomatología prefrontal y el estrés percibido. *Anales de Psicología*, 30(2), 676-683.
- Ruiz-Sánchez de León, J.M., Motos-Muñoz, M., Juárez, M., Sancho Cantus, D., Pedrero-Pérez, E.J., Fuente, C., y De la Rubia Ortí, J.E. (2019). Prefrontal

- Symptomatology in People with Amyotrophic Lateral Sclerosis: Clinical Utility and Psychometric Properties of Prefrontal Symptoms Inventory (PSI). *Prensa Médica Argentina*, 105(3), 1-4.
- Ruiz-Sánchez de León, J.M., Pedrero-Pérez, E.J., Gálvez, S., Fernández-Méndez, L.M., Lozoya-Delgado, P., y Departamento de Neuropsicología del Centro de Tratamiento de la Lesión Cerebral LES CER (2015). Utilidad clínica y propiedades psicométricas del inventario de síntomas prefrontales (ISP) en el daño cerebral adquirido y las demencias degenerativas. *Revista de Neurología*, 61(9), 387-394.
- Ruiz-Sánchez de León, J.M., y Pedrero Pérez, E.J. (2014). *Neuropsicología de la adicción*. Madrid: Editorial Médica Panamericana.
- (2019). *Neuropsicología de las conductas adictivas*. Madrid: Editorial Síntesis.
- Ruiz Sánchez de León, J.M., y González-Marqués, J. (2012). Evaluación neuropsicológica de la simulación. En H. González-Ordi, P. Santamaría Fernández y P. Capilla Ramírez (eds.), *Estrategias de detección de la simulación. Un manual clínico multidisciplinar* (pp. 275-300). Madrid: TEA Ediciones.
- Rush, B.K., McNeil, R.B., Gamble, D.M., Luke, S.H., Richie, A.N., Albers, C.S., et al. (2010). Behavioral symptoms in long-term survivors of ischemic stroke. *Journal of Stroke and Cerebrovascular Disease*, 19, 326-332.
- Ryden, M.B. (1988). Aggressive behavior in persons with dementia who live in the community. *Alzheimer Disease and Associated Disorders*, 2(4), 342-355.
- Sal y Rosas, H.J., Vega Diensmaier, J.M., Mazzotti Suarez, G., Vidal, H., Guimas, B., Adrianzén, C., y Vivar, R. (2002). Validación de una versión en español de la escala Yale-Brown para el trastorno obsesivo-compulsivo. *Actas Españolas de Psiquiatría*, 30(1), 30-35.
- Salazar, G.M., Martínez, M.K., y Ayala, V.H. (2001). *Adaptación y validación de la escala "Stages of change readiness and treatment eagerness scale, SOCRATES"*. México: Universidad Nacional Autónoma de México.
- Salek, S.S., Walker, M.D., y Bayer, A.J. (1999) The Community Dementia Quality of Life Profile (CDQLP): a factor analysis. *Quality of Life Research*, 8, 660.
- Sandin, B., Chorot, P., Santed, M.A., Jiménez, P., y Romero, M. (1994). Ansiedad cognitiva y somática: relación con otras variables de ansiedad y psicósomáticas. *Revista de Psicología General y Aplicada*, 47(3), 313-320.
- Sandín, B., y Chorot, P. (2003). Cuestionario de afrontamiento del estrés (CAE): Desarrollo y validación preliminar. *Revista de Psicopatología y Psicología Clínica*, 8, 39-54.
- Santed, M.A., Chorot, B., y Sandín, B. (1991). *Cuestionario de Estrés Diario (CED)*. Madrid: Universidad Nacional de Educación a Distancia.

- Santos, C.O., Caeiro, L., Ferro, J.M., Albuquerque, R., y Figueira, M.L. (2006). Anger, hostility and aggression in the first days of acute stroke. *European Journal of Neurology*, 13, 351-358.
- Sanz-Blasco, R. (2018). *Regulación emocional*. Madrid: Editorial Grupo 5.
- Sanz-Blasco, R., y Casado-Morales, M.I. (2010). Relación entre estrés y enfermedad de Alzheimer: una revisión teórica de los nuevos hallazgos. *EduPsykhé*, 9(2), 177-188.
- Sanz-Carrillo, C., García-Campayo, J., Rubio, A., Santed, M.A., y Montoro, M. (2002). Validation of the Spanish version of the Perceived Stress Questionnaire. *Journal of Psychosomatic Research*, 52, 167-172.
- Sanz, J., García-Vera, M.P., y Fortún, M. (2012). El “Inventario de ansiedad de Beck” (BAI): propiedades psicométricas de la versión española en pacientes con trastornos psicológicos. *Behavioral Psychology-Psicología Conductual*, 20, 563-583.
- Sanz, J., Perdigón, A.L., y Vázquez, C. (2003). Adaptación española del Inventario para la Depresión de Beck-II (BDI-II): propiedades psicométricas en población general. *Clínica y Salud*, 14(3), 249-280.
- Sanz, J., y García-Vera, M.P. (2007). Análisis psicométrico de las versiones breves del “Inventario para la Depresión de Beck” de 1978 (BDI-IA). *Psicología Conductual*, 15, 191-214
- Schagen, S., Schmand, B., De Sterke, S., y Lindeboom, J. (1997). Amsterdam Short-Term Memory Test: A new procedure for the detection of feigned memory deficits. *Journal of Clinical and Experimental Neuropsychology*, 19, 43-51.
- Schalock, R.L., y Verdugo, M.A. (2003). *Calidad de vida: manual para profesionales de la educación, salud y servicios sociales*. Madrid: Alianza.
- (2013). *Discapacidad e inclusión. Manual para la docencia*. Salamanca: Amarú Ediciones.
- Scholte op Reimer, W.J.M., De Haan, R.J., Pijnenborg, J.M.A., Limburg, M., y Van den Bos, G.A.M. (1998). Assessment of burden in partners of stroke patients with the Sense of competence Questionnaire. *Stroke*, 29, 373-379.
- Schulz, R., O'Brien, A.T., Bookwala, J., y Fleissner, K. (1995) Psychiatric and physical morbidity effects of dementia caregiving: Prevalence, correlates, and causes. *Gerontologist*, 35, 771-791.
- Sedó, M.A. (2007). *Test de los cinco dígitos (FDT)*. Madrid: TEA.
- Selai, C., Trimble, M., Rossor, M., y Harvey, R. (2001). Assessing quality of life in dementia. Preliminary psychometric testing of the Quality of Life Assessment Schedule (QOLAS). *Neuropsychol Rehabilitation*, 1, 219-43.
- Shankar K.K., Walker, M., Frost, D., y Orrell, M.W. (1999) The development of a valid and reliable scale for rating anxiety in dementia (RAID). *Ageing and Mental Health*, 3, 39-49.

- Shear, M.K., Brown, T.A., Barlow, D.H., Money, R., Sholomskas, D., Woods, S., Gorman, J. y Papp, L. (1997). Multicenter collaborative panic disorder severity scale. *American Journal of Psychiatry*, *154*, 1571-1575.
- Sherer, M., Bergloff, P., Boake, C., High, W., y Levin, E. (1998). The Awareness Questionnaire: factor structure and internal consistency. *Brain Injury*, *12*(1), 63-68.
- Sherman, D.S., Boone, K.B., Lu, P., y Razani, J. (2002). Re-examination of a Rey Auditory Verbal Learning Test/Rey Complex Figure discriminant function to detect suboptimal effort. *The Clinical Neuropsychologist*, *16*, 242-250.
- Shulman, L.M., Taback, R.L., Rabinstein, A.A., y Weiner, W.J. (2002). Non-recognition of depression and other non-motor symptoms in Parkinson's disease. *Parkinsonism and Related Disorders*, *8*(3) 193-197.
- Siegel, S.M. (1985). The Multidimensional Anger Inventory. En M.A. Chesney y N.H. Rosenman (ed.), *Anger and hostility in cardiovascular and behavioral disorders*. Washington, DC: Hemisphere.
- Sierra, J.C., Buela-Casal, G., y Miró, E. (2001). Evaluación y tratamiento de la ansiedad generalizada. En G. Buela-Casal y J.C. Sierra (eds.), *Manual de evaluación y tratamientos psicológicos*, 133-185. Madrid: Biblioteca Nueva.
- Silver, J.M., McAllister, T.M., y Yudofsky, S.C. (2005). *Textbook of traumatic brain injury*. Arlington: American Psychiatric Publishing.
- Sivrioglu, E.Y., Sivrioglu, K., Ertan, T., Ertan, F.S., Cankurtaran, E., Aki, O., Uluduz, D., Ince, B., y Kirli, S. (2009). Reliability and validity of the Geriatric Depression Scale in detection of poststroke minor depression. *Journal of Clinical and Experimental Neuropsychology*, *31*, 999-1006.
- Slick, D., Hopp, G., Strauss, E., y Thompson, G.B. (1997). *VSVT: Victoria Symptom Validity Test (Version 1.0)*. Odessa, Florida: Psychological Assessment Resources.
- Slick, D.J., Sherman, E.M., y Iverson, G.L. (1999). Diagnostic criteria for malingering neurocognitive dysfunction: Proposed standards for clinical practice and research. *The Clinical Neuropsychologist*, *13*, 545-561.
- Smets, E.M., Garssen, B., Bonke, B., y De Haes, J.C. (1995). The Multidimensional Fatigue Inventory (MFI) psychometric qualities of an instrument to assess fatigue. *Journal of Psychosomatic Research*, *39*, 315-325.
- Smith, S.C., Lamping, D.L., Banerjee, S., Harwood, R.H., Foley, B., Smith, P., et al. (2007). Development of a new measure of health-related quality of life for people with dementia: DEMQOL. *Psychological Medicine*, *37*, 737-746.
- Snaith, R.P., Baugh, S.J., Clayden, A.D., Husain, A., y Sipple, M.A. (1982). The Clinical Anxiety Scale: an instrument derived from the Hamilton Anxiety Scale. *The British Journal of Psychiatry*, *141*, 518-523.

- Sockeel, P., Dujardin, K., Devos, D., Denève, C., Destée, A., y Defebvre, L. (2006). The Lille Apathy Rating Scale (LARS), a new instrument for detecting and quantifying apathy: validation in Parkinson's disease. *Journal of Neurology, Neurosurgery and Psychiatry*, 77, 579-584.
- Soler, J., Pérez, V., Puigdemont, D., Pérez, J., Figueres, M., y Álvarez, E. (1997). Estudio de validación del Center for Epidemiologic Studies Depression Scale (CES-D) en una población española de pacientes con trastornos afectivos. *Actas Luso Españolas de Neurología y Psiquiatría y Ciencias Afines*, 25(4), 243-249.
- Solomon, R.L. (1980). The opponent-process theory of acquired motivation: the costs of pleasure and the benefits of pain. *American Psychologist*, 35, 691-712.
- Somme, J., Fernández-Martínez, M., Molano, A., y Zarranz, J.J. (2013). Neuropsychiatric symptoms in amnesic mild cognitive impairment: Increased risk and faster progression to dementia. *Current Alzheimer Research*, 10, 86-94.
- Spalletta, G., Girardi, P., Caltagirone, C., y Orfei, M.D. (2012). Anosognosia and neuropsychiatric symptoms and disorders in mild Alzheimer disease and mild cognitive impairment. *Journal of Alzheimer's Disease*, 29, 761-772.
- Spearing, M.K., Post, R.M., Leverich, G.S. Brandt, D., y Nolen, W. (1997). Modification of the Clinical Global Impressions (CGI). Scale for use in bipolar illness (BP): the CGIBP. *Psychiatry Research*, 73, 159-171.
- Spielberger, C.D. (1999). *The State-Trait Anger Expression Inventory-2: Professional Manual*. Odessa, FL: Psychological Assessment Resources.
- Spielberger, C.D., Gorsuch, R.L., y Lushene, R.E. (1970). *Manual for the State/Trait Anxiety Inventory*. Palo Alto: Consulting Psychologists Press.
- Spinella, M. (2003). Relationship between drug use and prefrontal associated traits. *Addiction Biology*, 8, 67-74.
- Spinella, M., y Lyke, J. (2004). Executive personality traits and eating behaviour. *International Journal of Neuroscience*, 114, 83-93.
- Spinella, M., Yang, B., y Lester, D. (2004). Prefrontal system dysfunction and credit card debt. *International Journal of Neuroscience*, 114, 1323-1332.
- Spitzer, R.L., Kroenke, K., y Williams, J.B. (1999). Validation and utility of a self-report version of PRIME-MD: the PHQ primary care study. Primary Care Evaluation of Mental Disorders. Patient Health Questionnaire. *JAMA*, 282(18), 1737-44.
- Stancin, T., Wade, S.L., Walz, N.C., Yeates, K.O., y Taylor, H.G. (2010). Family adaptation 18 months following traumatic brain injury in early childhood. *Journal of Developmental and Behavioral Pediatrics*, 31, 317-325.
- Starkstein, S.E., Fedoroff, J.P., Price, T.R., Leiguarda, R., y Robinson, R.G. (1992a). Anosognosia in patients with cerebrovascular lesions. A study of causative factors. *Stroke*, 23, 1446-1453.

- (1993). Apathy following cerebrovascular lesions. *Stroke*, 24, 1625-1630.
- Starkstein, S.E., Ingram, L., Garau, M.L., y Mizrahi, R. (2005). On the overlap between apathy and depression in dementia. *Journal of Neurology, Neurosurgery and Psychiatry*, 76, 1070-1074.
- Starkstein, S.E., Jorge, R., Mizrahi, R., y Robinson, R.G. (2006a). A prospective longitudinal study of apathy in Alzheimer's disease. *Journal of Neurology, Neurosurgery and Psychiatry*, 77, 8-11.
- (2006b). A diagnostic formulation for anosognosia in Alzheimer's disease. *Journal of Neurology, Neurosurgery and Psychiatry*, 77, 719-725.
- Starkstein, S.E., Mayberg, H.S., Preziosi, T.J., Andrezejewski, P., Leiguarda, R., y Robinson, R.G. (1992b). Reliability, validity, and clinical correlates of apathy in Parkinson's disease. *Journal of Neuropsychiatry and Clinical Neurosciences*, 4, 134-139.
- Starkstein, S.E., Petracca, G., Chemerinski, E., y Kremer, J. (2001). Syndromic validity of apathy in Alzheimer's disease. *American Journal of Psychiatry*, 158, 872-877.
- Starkstein, S.E., y Leentjens, A.F. (2008). The nosological position of apathy in clinical practice. *Journal of Neurology, Neurosurgery and Psychiatry*, 79, 1088-1092.
- Starkstein, S.E., y Merello, M. (2007). The Unified Parkinson's Disease Rating Scale: validation study of the mentation, behavior, and mood section. *Movement Disorders*, 22, 2156-2161.
- Steer, R.A., Rissmiller, D.J., y Beck, A.T. (2000). Use of the Beck Depression Inventory-II with depressed geriatric inpatients. *Behaviour Research and Therapy*, 38, 311-318.
- Steerneman, P. (1994). *Theory-of-mind screening-scale*. Leuven/Apeldoorn: Garant.
- Stella, F., Bucken-Gobbi, L.T., Gobbi, S., y Ana-Simões, C.S. (2007). Síntomas depresivos y trastorno motor en pacientes con enfermedad de Parkinson. *Revista de Neurología*, 45(10), 594-598.
- Stern, R.A. (1996). *Visual Analogue Mood Scales. Professional Manual*. Odesa, FL: Psychological Assessment Resources, Inc.
- Stone, V., Baron-Cohen, S., y Knoght, R. (1998). Frontal lobe contributions to theory of mind. *Journal of Cognitive Neuroscience*, 10, 640-656.
- Strauss, M.E., y Sperry, S.D. (2002). An informant-based assessment of apathy in Alzheimer disease. *Neuropsychiatry, Neuropsychology and Behavioral Neurology*, 15, 176-183.
- Stuss, D.T. (1991). Disturbance of self-awareness after frontal system damage. En G.P. Prigatano y D.L. Schacter (eds.), *Awareness of deficit after brain injury. Clinical and theoretical issues* (pp. 63-83). Nueva York: Oxford University Press.



- Stuss, D.T., Van Reekum, R., y Murphy, K.J. (2000). Differentiation of states and causes of apathy. En J.C. Borod (ed.), *The Neuropsychology of emotion* (pp. 340-363). Oxford: Oxford University Press.
- Sullivan, K., Winner, E., y Hopfield, N. (1995). How children tell a lie from a joke: The role of second-order mental attributions. *British Journal of Developmental Psychology*, *13*, 191-204.
- Sullivan, K., Zaitchik, D., y Tager-Flusberg, H. (1994). Preschoolers can attribute second-order beliefs. *Developmental Psychology*, *30*(3), 395-402.
- Sunderland, T., Alterman, I.S., Yount, D., Hill, J.L., Tariot, P.N., Newhouse, P.A., Mueller, E.A., Mellow, A.M., y Cohen, R.M. (1988). New Scale for the Assessment of Depressed Mood in Demented Patients. *American Journal of Psychiatry*, *145*(8), 955-959.
- Szurek, S. (1998). *Evaluación de la Depresión en la Tercera Edad: Escala de Evaluación Psicogeriátrica* [tesis doctoral]. Madrid: Facultad de Psicología de la Universidad Complutense de Madrid.
- Tager-Flusberg, H. (2003). Exploring the relationships between theory of mind and socialcommunicative functioning in children with autism. En B. Rapacholi y V. Slaughter (eds.), *Individual differences in theory of mind: Implications for typical and atypical development* (pp. 197-212). Londres: Psychology Press.
- Tariot, P.N., Mack, J.L., Patterson, M.B., Edland, S.D., et al. (1995). The Behavioral Rating Scale for Dementia of the Consortium to Establish a Registry for Alzheimer's Disease. *American Journal of Psychiatry*, *152*(9), 1349-1357.
- Tateno, A., Jorge, R.E., y Robinson, R.G. (2003). Clinical correlates of aggressive behavior after traumatic brain injury. *The Journal of Neuropsychiatry and Clinical Neurosciences*, *15*(2), 155-60.
- Tellez, N., Rio, J., Tintore, M., et al. (2005). Does the Modified Fatigue Impact Scale offer a more comprehensive assessment of fatigue in MS? *Multiple Sclerosis*, *11*, 198-202.
- Templer, D.I. (1970). The construction and validation of Death Anxiety Scale. *Journal of General Psychology*, *82*, 165-177.
- Terada, S., Ishizu, H., Fujisawa, Y., Fujita, D., Yokota, O., Nakashima, H., et al. (2002). Development and evaluation of a healthrelated quality of life questionnaire for the elderly with dementia in Japan. *International Journal of Geriatric Psychiatry*, *17*, 851-858.
- Terán-Mendoza, O., Sira-Ramos, D., Guerrero-Alcedo, J., y Arroyo-Alvarado, D. (2016). Sintomatología frontal, estrés autopercibido y quejas subjetivas de memoria en adictos a sustancias. *Revista de Neurología*, *62*, 296-302.
- Teri, L., Truax, P., Logsdon, R., Uomoto, J., Zarit, S., y Vitaliano, P.P. (1992). Assessment of Behavioral Problems in Dementia: The Revised Memory and Behavior Problems Checklist. *Psychology and Aging*, *7*(4), 622-631.

- Thompson, J.C., Harris, J., Sollom, A.C., Stopford, C.L., Howard, E., Snowden, J.S., y Craufurd, D. (2012). Longitudinal evaluation of neuropsychiatric symptoms in Huntington's disease. *Journal of Neuropsychiatry and Clinical Neurosciences*, 24, 53-60.
- Thompson, J.R., Bryant, B., Campbell, E.M., Craig, E.M., Hughes, C., Rotholz, D.A., Schalock, R.L., Silverman, W., Tasse, M.J., y Wehmeyer, M. (2004). *Supports Intensity Scale. Users manual*. Washington, DC: American Association on Mental Retardation.
- Thornton, M., y Travis, S. (2003). Analysis of the reliability of the Modified Caregiver Strain Index. *Journal of Gerontology: Social Sciences*, 58B, 127-132.
- Thorpe, S.J., Rolls, E.T., y Maddison, S. (1983). The orbitofrontal cortex: neuronal activity in the behaving monkey. *Experimental Brain Research*, 49, 93-115.
- Tirapu Ustárroz, J. (2007). La evaluación neuropsicológica. *Intervención Psicosocial*, 16(2), 189-211.
- Tirapu Ustárroz, J., y Ruiz Sánchez de León, J.M. (2011). Evaluación neurocognitiva en las adicciones. En E.J. Pedrero Pérez, J.M. Ruiz Sánchez de León, A. Verdejo García, M. Llanero Luque y E. Ambrosio Flores (eds.), *Neurociencias y adicciones* (pp. 111-134). Madrid: Sociedad Española de Toxicomanías.
- Tirapu Ustárroz, J., Pérez, G., Erekatxo, M., y Pelegrín, C. (2007). ¿Qué es la teoría de la mente? *Revista de Neurología*, 44, 479-489.
- Tirapu-Ustárroz, J., García-Molina, A., Luna-Lario, P., Roig-Rovira, T., y Pelegrín-Valero, C. (2008). Modelos de funciones y control ejecutivo (I). *Revista de Neurología*, 46(11), 684-692.
- Tobin, D. L., Holroyd, K. A., Reynolds, R. V., y Kigal, J. K. (1989). The hierarchical factor structure of the Coping Strategies Inventory. *Cognitive Therapy Research*, 13, 343-361.
- Tombaugh, T.N. (1996). *Test of Memory Malingering*. Toronto, Ontario, Canada: Multi-Health Systems.
- Tomkins, D.M., y Sellers, E.M. (2001). Addiction and the brain: the role of neurotransmitters in the cause and treatment of drug dependence. *Canadian Medical Association Journal*, 164(6), 817-821.
- Trigg, R., Skevington, S.M., y Jones, R.W. (2007). How can we best assess the quality of life of people with dementia? The Bath Assessment of Subjective Quality of Life in Dementia (BASQID). *Gerontologist*, 47, 789-797.
- Tyrer, P., Owen, R.T., y Cicchetti, D.V. (1984). The brief scale for anxiety: A subdivision of the comprehensive psychopathological rating scale. *Journal of Neurology, Neurosurgery and Psychiatry*, 47, 970-975.
- Vanheule, S., Desmet, M., Groenvynck, H., Rosseel, Y., y Fontaine, J. (2008). The factor structure of the Beck Depression Inventory-II: An evaluation. *Assessment*, 15, 177-187.

- Vázquez, C. (1995). Evaluación de trastornos depresivos y bipolares. En A. Roa (comp.), *Evaluación en Psicología Clínica y de la salud* (pp. 163-209). Madrid: CEPE.
- Vázquez, C., y Sanz, J. (1999). Fiabilidad y validez de la versión española del Inventario para la Depresión de Beck de 1978 en pacientes con trastornos psicológicos. *Clínica y Salud, 10*, 59-81.
- Velligan, D.I., Ritch, J.L., Sui, D., DiCocco, M., y Huntzinger, C.D. (2002). Frontal Systems Behavior Scale in schizophrenia: relationships with psychiatric symptomatology, cognition and adaptive function. *Psychiatry Research, 113*, 227-36.
- Verdugo, M.A., Arias, B., y Ibañez, A. (2007). *SIS. Escala de Intensidad de Apoyos. Manual. Adaptación española*. Madrid: TEA.
- Verdugo, M.A., Gómez, L.E., Fernández, M., Aguayo, V., y Arias, B. (2018). *Escala CAVIDACE: Evaluación de la calidad de vida de personas con daño cerebral*. Salamanca: INICO.
- Verhaeghe, S., Delfloor, T., y Grypdonck, M. (2005). Stress and coping among families of patients with traumatic brain injury: a review of the literature. *Journal of Clinical Nursing, 14*, 1004-1012.
- Vickery, C.D., Berry, D.T.R., Dearth, C.S., Vagini, V.L., Baser, R.E., Cragar, D.E., et al. (2004). Head injury and the ability to feign neuropsychological deficits. *Archives of Clinical Neuropsychology, 19*, 37-48.
- Vieta, E., Bobes, J., Ballesteros, J., González-Pinto, A., Luque, A., y Ibarra, N. (2008). Validity and reliability of the spanish versions of the Bech-Rafaelson's mania and melancholia scales for bipolar disorders. *Acta Psychiatrica Scandinavica, 117*(3), 207-15.
- Vieta, E., Torrent, C., Martínez-Arán, A., Colom, F., Reinares, M., Benabarre, A., Comes, M., y Goikolea, J.M. (2002). Escala sencilla de evaluación del curso del trastorno bipolar: CGI-BP-M. *Actas Españolas de Psiquiatría, 30*(5), 301-304.
- Vilalta-Franch, J., Lozano-Gallego, M., Hernández-Ferrándiz, M., Llinàs-Reglà, J., López-Pousa, S., y López, O.L. (1999). Neuropsychiatric inventory. Propiedades psicométricas de su adaptación al español. *Revista de Neurología, 29*(1), 15-19.
- Viner, R., y Christie, D. (2005). Fatigue and somatic symptoms. *BMJ (Clinical Research Ed.), 330*, 1012-1015.
- Visser-Keizer, A.C., Meyboom de Jong, B., Deelman, B.G., Berg, I.J., y Gerritsen, M.J.J. (2002). Subjective changes in emotion, cognition and behaviour after stroke: actors affecting the perception of patients and partners. *Journal of Clinical and Experimental Neuropsychology, 24*(8), 1032-1045.
- Vitaliano, P.P., Russo, J., Young, H.M., Becker, J., y Maiuro, R.D. (1991). The Screen for Caregiver Burden. *Gerontologist, 31*, 76-83.
- Von Steinbüchel, N., Wilson, L., Gibbons, H., Muehlan, H., Schmidt, H., Schmidt, S., y Truelle, J. L. (2012). QOLIBRI overall scale: a brief index of

- health-related quality of life after traumatic brain injury. *Journal of Neurology, Neurosurgery & Psychiatry*, 83(11), 1041-1047.
- Von Stenbüchel, *et al.*, (2010). Quality of Life after Brain Injury (QOLIBRI): scale validity and correlates of quality of life. *Journal of Neurotrauma*, 27(7), 1157-1165.
- Wagner, H. (1989). The physiological differentiation of emotions. En H. Wagner y A. Manstead (eds.), *Handbook of Social Psychophysiology*. Nueva York: Wiley.
- Ware, J.E., Snow, K.K., Kosinski, M., y Gandek, B. (1993). *SF-36 Health Survey. Manual and Interpretation Guide*. Boston, MA: The Health Institute, New England Medical Center.
- Warrington, E. (1984). *Recognition Memory Test*. Windsor, UK: NFER Nelson.
- Warrington, E.K., y James, M. (1991). *Visual Object and Space Perception Battery*. Suffolk: Thames Valley Test Co.
- (1993). *Batería de test para la percepción visual de objetos y del espacio (VOSP)*. Madrid: TEA Ediciones.
- Wechsler, D. (2012). *Escala de inteligencia de Wechsler para adultos (WAIS-IV)*. Madrid: Pearson.
- (2013). *Escala de Memoria Wechsler para Adultos (WMS-IV)*. Madrid: Pearson.
- Weiner, M.F., Martin-Cook, K., Svetlik, D.A., Saine, K., Foster, B., y Fontaine, C.S. (2000). The Quality of Life in Late-Stage Dementia (QUALID) scale. *JAMA*, 1, 114-116.
- Weintraub, D., Oehlberg, K.A., Katz, I.R., y Stern, M.B. (2006). Test characteristics of the 15-item geriatric depression scale and Hamilton Depression Rating Scale in Parkinson disease. *American Journal of Geriatric Psychiatry*, 14, 169-175.
- Wellman, H.M., y Liu, D. (2004). Scaling theory-of-mind tasks. *Child Development*, 75, 523-541.
- Whitehead, L. (2009). The measurement of fatigue in chronic illness: a systematic review of unidimensional and multidimensional fatigue measures. *Journal of Pain and Symptom Management*, 37(1), 107-128.
- Whitehouse, P., Orgogozo, J., Becker, R., Gauthier, S., Pontecorvo, M., Erzigkeit, H., Rogers, S., Mohs, R.C., Bodick, N., Bruno, G., y Dal-Bianco, P. (1997). Quality-of-life assessment in dementia drug development. *Alzheimer Disease and Associated Disorders*, 11, 56-60.
- Wielenga-Boiten, J.E., Heijenbrok-Kal, M.H., y Ribbers, G.M. (2015). The Relationship of Health Locus of Control and Health-Related Quality of Life in the Chronic Phase After Traumatic Brain Injury. *The Journal of Head Trauma Rehabilitation*, 30(6), 424-431.
- Wiggins, E.C., y Brandt, J. (1988). The detection of simulated amnesia. *Law and Human Behavior*, 12, 57-78.

- Willer, B., Flaherty, P.M., y Coallier, S. (2001). Families living with acquired brain injury. En R.L. Wood y T.M. McMillan (eds.), *Neurobehavioural disability and social handicap following traumatic brain injury* (pp. 47-63). Nueva York: Oxford University Press.
- Wilson, B.A., Alderman, N., Burgess, P.W., Emslie, H., y Evans, J.J. (1996). *Behavioural assessment of the dysexecutive syndrome*. Bury St. Edmunds, UK: Thames Valley Test.
- Wimmer, H., y Perner, J. (1983). Beliefs about beliefs: representation and constraining function of wrong beliefs in young children's understanding of deception. *Cognition*, 21, 103-128.
- Winner, E., y Leekam, S. (1991). Distinguishing irony from deception: understanding the speaker's second-order intention. *British Journal of Developmental Psychology*, 9, 257-270.
- Wittchen, H.U., y Boyer, P. (1998). Screening for anxiety disorders. Sensitivity and specificity of the Anxiety Screening Questionnaire (ASQ-15). *The British Journal of Psychiatry*, 173(S34), 10-17.
- Wood, R.L., Williams, C., y Lewis, R. (2010). Role of alexithymia in suicide ideation after traumatic brain injury. *Journal of the International Neuropsychological Society*, 16, 1108-1114.
- Yarbus, A.L. (1967). *Eye movements and vision*. Nueva York: Plenum.
- Yesavage, J.A. y Brink, T.L. (1983). Development and validation of a geriatric depression screening scale: a preliminary report. *Journal of Psychiatric Research*, 17(1), 37-49.
- Young, A.W., Perrett, D., Calder, A., et al. (2002). *Facial Expressions of Emotion: Stimuli and Test*. Bury St Edmunds: Thames Valley Test Co.
- Young, R.C., Biggs, J.T., Ziegler, V.E., y Meyer, D.A. (1978). A rating scale for mania: reliability, validity, and sensitivity. *British Journal of Psychiatry: the Journal of Mental Science*, 133, 429-435.
- Yudofsky, S.C., Silver, J.M., y Jackson, W. (1986). The Overt Aggression Scale for the objective rating of verbal and physical aggression. *American Journal of Psychiatry*, 143, 35-39.
- Zald, D.H., y Andreotti, C. (2010). Neuropsychological assessment of the orbital and ventromedial prefrontal cortex. *Neuropsychologia*, 48(12), 3377-3391.
- Zarit, S.H., Reever, K.E., y Bach-Peterson, J. (1980). Relatives of the impaired elderly. Correlates of feelings of burden. *Gerontologist*, 20, 649-654.
- Zelin, M.I., Adler, G., y Myerson, P.G. (1972). Anger Self-Report: An objective questionnaire for the measurement of aggression. *Journal of Consulting and Clinical Psychology*, 39, 340.
- Zigmong, A.S., y Snaith, R.P. (1983). The hospital anxiety and depression. *Acta Psychiatrica Scandinavica*, 67, 361-370.
- Zung, W.W.K. (1965). A Self-rating Depression Scale. *Archives of General Psychiatry*, 12, 63-70.