

Bibliografía

- Addis, M. E., & Martell, C. R. (2004). *Overcoming Depression One Step at a Time: The New Behavioral Activation Approach to Getting Your Life Back*. New York: New Harbinger.
- Arciero, G., & Bondolfi, G. (2009). *Selfhood, identity and personality styles*. Oxford: Wiley.
- Ambrosini, A., Stanghellini, G., & Langer, A. I. (2011). El Typus melancholicus de Tellenbach en la actualidad: una revisión sobre la personalidad premórbida vulnerable a la melancolía. *Actas Españolas de Psiquiatría*, 39, 302-311.
- Armento, M. E. A., & Hopko, D. R. (2007). The Environmental Reward Observation Scale (EROS): Development, validity, and reliability. *Behavior Therapy*, 38, 107-119.
- (2009). Behavioral Activation of a breast cancer patient with coexistent Major Depression and Generalized Anxiety Disorder. *Clinical Case Studies*, 8(1), 25-37.
- Barraca, J. (2009a). La Activación Conductual (AC) y la Terapia de Activación Conductual para la Depresión (TACD): Dos protocolos de tratamiento desde el modelo de la activación conductual. *EduPsykhé*, 8, 23-47.
- (2009b). Habilidades clínicas en la terapia conductual de tercera generación. *Clínica y Salud*, 20 (2), 109-117.
- (2010). Aplicación de la Activación Conductual en un paciente con sintomatología depresiva. *Clínica y Salud*, 21 (2), 183-197.
- (2011). *El viaje al ahora. Una guía sencilla para llevar la atención plena a nuestro día a día*. Bilbao: Desclée de Brouwer.
- (2014). Aplicación de la Activación Conductual (AC) en un paciente con Distimia. Comunicación presentada en el Simposio Invitado "Casos clínicos abordados con terapias de tercera generación" en el *X Congreso Nacional de Casos Clínicos de Trastornos Psicopatológicos*. Granada, 22-24 de mayo de 2014.

- Barraca, J. y Pérez-Álvarez, M. (2010). Adaptación Española del Environmental Reward Observation Scale (EROS). *Ansiedad y Estrés*, 16 (1), 95-107.
- (2015). Traducción del *Reward Probability Index* – RPI. Manuscrito no publicado.
- Barraca, J., Pérez-Álvarez, M., & Lozano Bleda, J. H. (2011). Avoidance and activation as keys to depression: Adaptation of the Behavioral Activation for Depression Scale (BADs) in a Spanish sample. *Spanish Journal of Psychology*, 14, 998-1009.
- Barth, J., Munder, T., Gerger, H, Neike et al. (2013) Comparative Efficacy of Seven Psychotherapeutic Interventions for Patients with Depression: A Network Meta-Analysis. *PLoS Med* 10(5): e1001454. doi:10.1371/journal.pmed.1001454.
- Becerra-Gálvez, A. L. Lugo-Gonzalez, I. V. & Reynoso-Erazo, L. (2013). Terapia de activación conductual en pacientes oncológicos: Una propuesta de intervención. *Psicooncología*, 10(1), 163-168.
- Beck, A. T., Rush, A. J., Shaw, B. F., & Emery, G. (1979). *Cognitive therapy of depression*. New York: Guilford.
- Beck, A. T., Steer, R. A., & Brown, G. K. (1996). *Manual for the Beck Depression Inventory-II*. San Antonio, TX: Psychological Corporation.
- Busch, A. M., Whited, M. C., Appelhans, B. M., Schneider, K. L., Wa-ring, M. E., DeBiasse, M. A., Oleski, J. L., Crawford, S. L., Pagoto, S. L. (2013). Reliable change in depression during behavioral weight loss treatment among women with major depression. *Obesity*, 21(3), E211-8. doi: 10.1002/oby.20113.
- Carey, T. A. (2011). Exposure and reorganization: The what and how of effective psychotherapy. *Clinical Psychology Review*, 31, 236–248.
- Carvalho, J. P., Gawrysiak, M. J., Hellmuth, J. C., McNulty, J. K., Magidson, J. F., Lejuez, C. W., Hopko, D. R. (2011). The Reward Probability Index: Design and validation of a scale measuring access to environmental reward. *Behavior Therapy*, 42(2), 249-262.
- Chu, B. C., Colognori, D., Weissman, A. S., & Bannon, K. (2009). An initial description and pilot of group behavioral activation therapy for anxious and depressed youth. *Cognitive and Behavioral Practice*, 16(4), 408-419.
- Ciamarella, A., & Poli, P. (2001). Assessment of depression among cancer patients: The role of pain, cancer type, and treatment. *Psycho-Oncology*, 10, 156-165
- Collado, A., Long, K. E., MacPherson, L., & Lejuez, K. W. (2014). The efficacy of a behavioral activation intervention among depressed US Latinos with limited English language proficiency: study protocol for a randomized controlled trial. *Trials*, 15: 231.
- Cogle, J. R. (2012). What Makes a Quality Therapy? A Consideration of Parsimony, Ease, and Efficiency. *Behavior Therapy*, 43, 468-481.

- Cuijpers, P., van Straten, A., & Warmerdam, L. (2007). Behavioral activation treatments of depression: A meta-analysis. *Clinical Psychology Review, 27*, 318-326.
- Davidson, T. M., Yuen, E. K., Felton, J. W., McCauley, J., Gros, K. S., & Ruggiero, K. J. (2014). Feasibility assessment of a brief, web-based behavioral activation intervention for adolescents with depressed mood. *International Journal of Psychiatry in Medicine, 48*(1), 69-82.
- Danziger, K. (1997). *Naming the Mind How Psychology Found Its Language*. Toronto: York University.
- Daughters, S. B., Braun, A. R., Sargeant, M. N., Reynolds, E. K., Hopko, D. R., Blanco, C., & Lejuez, C. (2008). Effectiveness of a brief behavioral treatment for inner-city illicit drug users with elevated depressive symptoms: The Life Enhancement Treatment for Substance Use (LETS Act!). *Journal of Clinical Psychiatry, 69*(1), 122-129
- Dimidjian S., Hollon S. D., Dobson K. S., Schmalting K. B., Kohlenberg, R. J., Addis, M. E., Gallop, R., McGlinchey, J. B., Markley, D. K., Gollan, J. K., Atkins, D. C., & Dunner, D. L. (2006). Randomized Trial of Behavioral Activation, Cognitive Therapy, and Antidepressant Medication in the Acute Treatment of Adults With Major Depression. *Journal of Consulting and Clinical Psychology, 74*, 658-670.
- Dimidjian, S., Martell, C. R., Addis, M. E., & Herman-Dunn, R. (2008). Behavioral Activation for Depression. In D. H. Barlow (Ed.), *Clinical Handbook of Psychological Disorders. A step-by-step treatment manual (4^a ed.)* (pp. 328-364). New York: Guilford.
- Dobson, K. S., Hollon, S. D., Dimidjian, S., Schmalting, K. B., Kohlenberg, R. J., Gallop, R., et al. (2008). Randomized trial of behavioral activation, cognitive therapy, and anti-depressant medication in the prevention of relapse and recurrence of major depression. *Journal of Consulting and Clinical Psychology, 76*, 468-477.
- D'Zurilla, T. J. & Goldfried, M. R. (1971). Problem solving and behavior modification. *Journal of Abnormal Psychology, 78*, 107-126.
- D'Zurilla, T. J., & Nezu, A. M. (1999). *Problem-solving therapy: A social competence approach to clinical intervention (2nd ed.)*. New York: Springer.
- Ekers, D., Webster, L., Van Straten, A., Cuijpers, P., Richards, D., et al. (2014) Behavioural Activation for Depression: An Update of Meta-Analysis of Effectiveness and Sub Group Analysis. *PLoS ONE 9*(6): e100100. doi: 10.1371/journal.pone.0100100
- Fernández-Rodríguez, C., Villoria-Fernández, E., Amigo, I., Padierna, C., Gracia, J. M., Fernández, R., et al. (2011). Terapia de activación conductual en pacientes con cáncer. *Anales De Psicología, 27*(2), 278-291.
- Fernández-Rodríguez, C., Villoria-Fernández, E., Fernández-García, P., & González-Fernández, S. (2014). Efectos de la activación conductual en la cali-

- dad de vida y estado emocional de los pacientes con cáncer de pulmón. *Psicooncología*, 11(2-3), 199-215
- Ferster, C. B. (1973). A functional analysis of depression. *American Psychologist*, 28, 850-870.
- First, M. B., Spitzer, R. L., Gibbon, M., & Williams, J. B. W. (1997). *User's guide for the Structured Clinical Interview for DSM-IV Axis I Disorders*. Washington, DC: American Psychiatric Press.
- First, M. B., Spitzer, R. L., Gibbon, M., & Williams, J. B. W., & Benjamin, L. (1996). *User's guide for the Structured Clinical Interview for DSM-IV Axis II Personality Disorders (SCID-II)*. New York: Biometrics Research Department, New York State Psychiatric Institute.
- Fuchs, T. (2013). Existential vulnerability: toward a psychopathology of limit situations. *Psychopathology*, 46, 301-308.
- García-Toro, M., Ibarra, O., Gili, M., Serrano, M. J., Oliván, B., Vicens, E., & Roca, M. (2012). Four hygienic-dietary recommendations as add-on treatment in depression: a randomized-controlled trial. *Journal of Affective Disorders*, 140(2), 200-203.
- Gortner, E. T., Gollan, J. K., Dobson, K. S., & Jacobson, N. S. (1998). Cognitive-behavioral treatment for depression: Relapse prevention. *Journal of Consulting and Clinical Psychology*, 66, 377-384.
- Gros, D. F., Price, M., Strachan, M., Yuen, E. K., Milanak, M. E., & Acierno, R. (2012). Behavioral activation and therapeutic exposure: An investigation of relative symptom changes in PTSD and depression during the course of integrated behavioral activation, situational exposure, and imaginal exposure techniques. *Behavior Modification*, 36(4), 580-599.
- Hacking, I. (1995). The looping effects of human kinds. In D. Sperber, D. Premack, & A. Premack (Eds.), *Causal Cognition: An Interdisciplinary Approach* (pp. 351-383). Oxford: Oxford University Press.
- Hamilton, M. A. (1960). A rating scale for depression. *Journal of Neurology, Neurosurgery, and Psychiatry*, 23, 56-61.
- Hayes, S. C., Strosahl, K. D., & Wilson, K. G. (1999). *Acceptance and commitment therapy: An experiential approach to behavior change*. New York: Guilford.
- Hopko, D. R., Armento, M. E. A., Robertson, S., Ryba, M. M., Carvalho, J. P., Colman, L. K., et al. (2011). Brief behavioral activation and problem-solving therapy for depressed breast cancer patients: Randomized trial. *Journal of Consulting and Clinical Psychology*, 79, 834-849.
- Hopko, D. R.; Cannity, K., McIndoo, C. C., File, A. A., Ryba, M. M., Clark, C. G., & Bell, J. L. (2015). Behavior therapy for depressed breast cancer patients: Predictors of treatment outcome. *Journal of Consulting and Clinical Psychology*, 83(1), 225-231

- Hopko, D. R., Lejuez, C., & Hopko, S. D. (2004). Behavioral activation as an intervention for coexistent depressive and anxiety symptoms. *Clinical Case Studies, 3*(1), 37-48.
- Hopko, D. R., Magidson, J. F., & Lejuez, C. W. (2011). Treatment failure in behavior therapy: Focus on behavioral activation for depression. *Journal of Clinical Psychology, 67*(11), 1106-1116.
- Hopko, D., Sanchez, L., Hopko, S., Dvir, S., & Lejuez, C. (2003). Behavioral activation and the prevention of suicidal behaviors in patients with borderline personality disorder. *Journal of Personality Disorders, 17*, 460-478.
- Houghton, S., Curran, J., & Saxon, D. (2008). An uncontrolled evaluation of group behavioural activation for depression. *Behavioural and Cognitive Psychotherapy, 36*, 235-239.
- Jacob, M., Keeley, M. L., Ritschel, L., & Craighead, W. E. (2013). Behavioural activation for the treatment of low-income, african american adolescents with major depressive disorder: A case series. *Clinical Psychology & Psychotherapy, 20*(1), 87-96.
- Jacobson, N. S., Dobson, K.S., Truax, P. A., Addis, M. E., Koerner, K., Gollan, J. K., Gortner, E., & Prince, S. E. (1996). A component analysis of cognitive-behavioral treatment for depression. *Journal of Consulting and Clinical Psychology, 64*, 295-304.
- Jahoda, A., Melville, C. A., Pert, C., Cooper, S., Lynn, H., Williams, C., Davidson, C.. (2014). A feasibility study of behavioural activation for depressive symptoms in adults with intellectual disabilities. *Journal of Intellectual Disability Research*, Dec 14. Epub 2014. doi: 10.1111/jir.12175.
- Jakupcak, M., Roberts, L. J., Martell, C., Mulick, P., Michael, S., Reed, R., et al. (2006). A pilot study of behavioral activation for veterans with posttraumatic stress disorder. *Journal of Traumatic Stress, 19*(3), 387-391.
- Jakupcak, M., Wagner, A., Paulson, A., Varra, A., & McFall, M. (2010). Behavioral activation as a primary care-based treatment for PTSD and depression among returning veterans. *Journal of Traumatic Stress, 23*(4), 491-495.
- Kanter, J. W., Busch, A. M. y Rusch, L. C. (2009). *Behavioral Activation: Distinctive Features*. New York: Routledge.
- Kanter, J. W., Mulick, P. S., Busch, A. M., Berlin, K. S. y Martell, C. R. (2007). The Behavioral Activation for Depression Scale (BADs): Psychometric Properties and Factor Structure. *Journal of Psychopathology and Behavior Assessment, 29*, 191-202.
- Kanter, J. W., Rusch, L. C., Busch, A. M., & Sedivy, S. K. (2009). Validation of the Behavioral Activation for Depression Scale (BADs) in a community sample with elevated depressive symptoms. *Journal of Psychopathology and Behavioral Assessment, 31*, 36-42
- Kanter, J. W., Santiago-Rivera, A. L., Santos, M. M., Nagy, G., López, M., Hurtado, G. D., & West, P. (2014). A randomized hybrid efficacy and effective-

- ness trial of behavioral activation for latinos with depression. *Behavior Therapy*, 46 (2), 177-192.
- Kohlenberg, R. J., & Tsai, M. (1991). *Functional analytic psychotherapy: Creating intense and curative therapeutic relationships*. New York: Plenum.
- Lazzari, C., Egan, S. J., & Rees, C. S. (2011). Behavioral activation treatment for depression in older adults delivered via videoconferencing: A pilot study. *Cognitive and Behavioral Practice*, 18(4), 555-565.
- Lejuez, C.W., Hopko, D.R., Acierno, R., Daughters, S.B, & Pagoto, S.L. (2011). Ten year revision of the brief behavioral activation treatment for depression: revised treatment manual. *Behavior Modification*, 35, 111-161.
- Lejuez, C. W., Hopko, D. R. y Hopko, S. D. (2001). A brief Behavioral Activation Treatment for Depression: Treatment Manual. *Behavior Modification*, 25, 225-286.
- Lewinsohn, P. M. (1974). A behavioral approach to depression. In R. M. Friedman, & M. M. Katz (Eds.), *The psychology of depression: Contemporary theory and research* (pp. 157-185). New York: John Wiley.
- Lewinsohn, P. M., Muñoz, R. F., Youngren, M. A., & Zeiss, A. M. (1978). *Control your depression*. Englewoods Cliffs, NJ: Prentice Hall.
- Linehan, M. M. (1993) *Cognitive-behavioral treatment of borderline personality disorder*. New York: Guilford.
- Lucas, R. E., Clark, A. E., Georgellis, Y., & Diener, E. (2004). Unemployment alters the set-point for life satisfaction. *Psychological Science*, 15, 8-13.
- Lundervold, D., Talley, C., & Buermann, M. (2008). Effect behavioral of behavioral activation treatment on chronic fibromyalgia pain: replication and extension. *International Journal of Behavioral Consultation and Therapy*, 4, 146-157.
- MacPherson, L., Tull, M. T., Matusiewicz, A. K., Rodman, S., Strong, D. R., Kahler, C. W., Lejuez, C. W. (2010). Randomized controlled trial of behavioral activation smoking cessation treatment for smokers with elevated depressive symptoms. *Journal of Consulting and Clinical Psychology*, 78(1), 55-61.
- Maj, M. (2011). When does depression become a mental disorder? *British Journal of Psychiatry*, 199, 85-86.
- Martell, C. R., Addis, M. E., & Jacobson, N. S. (2001). *Depression in Context: Strategies for Guided Action*. New York: W. W. Norton.
- Martell, C. R., Dimidjian, S., & Herman-Dunn, R. (2010). *Behavioral Activation for Depression: A Clinician's Guide*. New York: Guilford.
- McCauley, E., Schloretd, K., Gudmundsen, G., Martell, C., & Dimidjian, S. (2011). Expanding behavioral activation to depressed adolescents: Lessons learned in treatment development. *Cognitive and Behavioral Practice*, 18(3), 371-383.

- Merikangas, K. R., Risch, N. J., & Weissman, M. M. (1994). Comorbidity and co-transmission of alcoholism, anxiety and depression. *Psychological Medicine*, 24, 69-80.
- Mineka, S., Watson, D., & Clark, L. A. (1998). Comorbidity of anxiety and unipolar mood disorders. *Annual Review of Psychology*, 49, 377-412
- Moss, K., Scogin, F., Di Napoli, E., & Presnell, A. (2012). A self-help behavioral activation treatment for geriatric depressive symptoms. *Aging & Mental Health*, 16(5), 625-635.
- Mulick, P. S., & Naugle, A. E. (2004). Behavioral activation for comorbid PTSD and major depression: A case study. *Cognitive and Behavioral Practice*, 11(4), 378-387.
- (2010). Behavioral activation in the treatment of comorbid posttraumatic stress disorder and major depressive disorder. *International Journal of Behavioral Consultation and Therapy*, 5(3-4), 330-350.
- Pagoto, S., Bodenlos, J. S., Schneider, K. L., Olendzki, B., Spates, C. R., & Ma, Y. (2008). Initial investigation of behavioral activation therapy for comorbid major depressive disorder and obesity. *Psychotherapy: Theory, Research, Practice, Training*, 45(3), 410-415.
- Pasterfield, M., Bailey, D., Hems, D., McMillan, D., Richards, D., & Gilbody, S. (2014). Adapting manualized behavioural activation treatment for older adults with depression. *The Cognitive Behaviour Therapist*, 7, e5 DOI: <http://dx.doi.org/10.1017/S1754470X14000038>
- Pérez Álvarez, M. (2007). La activación conductual y la desmedicalización de la depresión. *Papeles del Psicólogo*, 28, 97-110.
- (2014). *Las terapias de tercera generación como terapias contextuales*. Madrid: Síntesis.
- Plagge, J. M., Lu, M. W., Lovejoy, T. I., Karl, A. I., & Dobscha, S. K. (2013). Treatment of comorbid pain and PTSD in returning veterans: A collaborative approach utilizing behavioral activation. *Pain Medicine*, 14(8), 1164-1172.
- Polenick, C. A., & Flora, S. R. (2013). Behavioral activation for depression in older adults: Theoretical and practical considerations. *The Behavior Analyst*, 36(1), 35-55.
- Porter, J. F., Spates, R. C., & Smitham, S. (2004). Behavioral activation group therapy in public mental health settings: A pilot investigation. *Professional Psychology: Research and Practice*, 35, 297-301
- Ratcliffe, M (2014). *Experiences of depression. A study in phenomenology*. Oxford: Oxford University Press.
- Rhodes, S., Richards, D. A., Ekers, D., McMillan, D., Byford, S., Farrand, P. A., Gilbody, S., Hollon, S. D., Kuyken, W., Martell, C., O'Mahen, H. A., O'Neill, E., Reed, N., Taylor, R. S., Watkins, E. R., & Wright, K. A. (2014). Cost and outcome of behavioural activation versus cognitive behaviour

- therapy for depression (COBRA): study protocol for a randomised controlled trial. *Trials* 15: 29.
- Ritschel, L. A., Ramirez, C. L., Jones, M., & Craighead, W. E. (2011). Behavioral activation for depressed teens: A pilot study. *Cognitive and Behavioral Practice*, 18(2), 281-299.
- Romero Gamero, R., Poves Oñate, S., Vucinovich, N. (2011). Terapia de activación conductual para la depresión aplicación a un paciente con esquizofrenia paranoide. *Análisis y modificación de conducta*, 37, 65-75.
- Ruggiero, K. J., Morris, T. L., Hopko, D. R., & Lejuez, C. (2007). Application of behavioral activation treatment for depression to an adolescent with a history of child maltreatment. *Clinical Case Studies*, 6 (1), 64-78.
- Ryba, M. M., Lejuez, C. W., & Hopko, D. R. (2014). Behavioral activation for depressed breast cancer patients: The impact of therapeutic compliance and quantity of activities completed on symptom reduction. *Journal of Consulting and Clinical Psychology*, 82(2), 325-335.
- Santiago-Rivera, A.L., Kanter, J., Benson, G., DeRose, T., Illes, R., & Reyes, W. (2008). Behavioral Activation as an Alternative Treatment Approach for Latinos with Depression. *Psychotherapy: Theory, Research, Practice, training*, 45,173-185.
- Segal, Z. V., Williams, J. M. G., & Teasdale, J. D. (2002). *Mindfulness-based cognitive therapy for depression: A new approach to preventing relapse*. New York: Guilford.
- Snarski, M., Scogin, F., DiNapoli, E., Presnell, A., McAlpine, J., & Marcinak, J. (2011). The effects of behavioral activation therapy with inpatient geriatric psychiatry patients. *Behavior Therapy*, 42(1), 100-108.
- Spiegel, D., & Giese-Davis, J. (2003). Depression and cancer: Mechanisms and disease progression. *Biological Psychiatry*, 54, 269-282.
- Tellenbach, H. (1974). *Melancolía*. Madrid: Morata (traducción de 1976).
- Torres Ceja, C., Ortiz Rueda, G., Rangel Bernal, N., & González Becerra, V. (2012). Análisis del concepto de situación en la estructuración de las funciones psicológicas. In M. A. Padilla Vargas y R. Pérez-Almonacid (Eds.), *La función sustitutiva referencial: análisis histórico-crítico/avances y perspectivas* (pp. 149-168). New Orleans: University Press of the South.
- Treynor, W. F. C., González, R., Nolen-Hoeksema, S. (2003) Rumination reconsidered: A psychometric analysis. *Cognitive Therapy and Research*, 27, 247-259.
- Turner, A. P., & Jakupcak, M. (2010). Behavioral activation for treatment of PTSD and depression in an iraqi combat veteran with multiple physical injuries. *Behavioural and Cognitive Psychotherapy*, 38(3), 355-361.
- Turner, J. S., & Leach, D. J. (2010). Experimental evaluation of behavioral activation treatment of anxiety (BATA) in three older adults. *International Journal of Behavioral Consultation and Therapy*, 6(4), 373-394.

Bibliografia

- (2012). Behavioural activation therapy: Philosophy, concepts, and techniques. *Behaviour Change*, 29(2), 77-96.
- Wagner, A. W., Zatzick, D. F., Ghesquiere, A., & Jurkovich, G. J. (2007). Behavioral activation as an early intervention for posttraumatic stress disorder and depression among physically injured trauma survivors. *Cognitive and Behavioral Practice*, 14(4), 341-349.
- Wallis, A., Roeger, L., Milan, S., Walmsley, C., & Allison, S. (2012). Behavioural activation for the treatment of rural adolescents with depression. *The Australian Journal of Rural Health*, 20(2), 95-96.
- Wilson, K. G., Sandoz, E. K., Kitchens, J., & Roberts, M. (2010). The Valued Living Questionnaire: Defining and Measuring Valued Action within a Behavioral Framework. *The Psychological Record*, 60, 249-272.
- Yon, A., & Scogin, F. (2009). Behavioral activation as a treatment for geriatric depression. *Clinical Gerontologist: The Journal of Aging and Mental Health*, 32(1), 91-103.
- Zimmerman, M., Ellison, W., Young, D., Chelminski, I., & Dalrymple, K. (2015). How many different ways do patients meet the diagnostic criteria for major depressive disorder? *Comprehensive Psychiatry*, 56, 29-34.